

ON THE HOMOTOPY TYPE OF THE SPACE OF METRICS OF POSITIVE SCALAR CURVATURE

JOHANNES EBERT, MICHAEL WIEMELER

ABSTRACT. The main result of this paper is that when M_0, M_1 are two simply connected spin manifolds of the same dimension $d \geq 5$ which both admit a metric of positive scalar curvature, the spaces $\mathcal{R}^+(M_0)$ and $\mathcal{R}^+(M_1)$ of such metrics are homotopy equivalent. This supersedes a previous result of Chernysh and Walsh which gives the same conclusion when M_0 and M_1 are also spin cobordant.

We also prove an analogous result for simply connected manifolds which do not admit a spin structure; we need to assume that $d \neq 8$ in that case.

CONTENTS

1. Introduction	2
Acknowledgements	3
2. A cut-and-paste construction	4
2.1. The cobordism theorem	4
2.2. Admissible splittings	5
3. Admissible splittings of projective bundles	7
3.1. Connection metrics	8
3.2. The curvature of connection metrics	9
3.3. The generalized Gromov–Lawson–Chernysh theorem	11
3.4. Construction of the splittings	12
4. The computation in the spin case	17
4.1. The simply connected case	17
4.2. Generalization to some nontrivial fundamental groups	19
5. Two further geometric ingredients for the non-spin case	20
5.1. Stability of product metrics	20
5.2. Splitting a certain 5-manifold	23
6. The computation in the non-spin case	27
6.1. Proof of Theorem B	27
6.2. The structure of the oriented cobordism ring	28
References	31

Date: December 2, 2020.

Both authors were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy EXC 2044 – 390685587, Mathematics Münster: Dynamics–Geometry–Structure.

1. INTRODUCTION

Let $\mathcal{R}^+(M)$ be the space of Riemannian metrics of positive scalar curvature (“psc metrics”) on the closed manifold M . The main result of this paper concerns the homotopy type of $\mathcal{R}^+(M)$, when M is simply connected and of dimension $d \geq 5$. There are two cases to distinguish: M may admit a spin structure or not.

Theorem A. *Let M be a simply connected closed spin manifold of dimension $d \geq 5$. Then if M admits a psc metric, there is a homotopy equivalence*

$$\mathcal{R}^+(M) \simeq \mathcal{R}^+(S^d).$$

- Remark 1.1.*
- (1) It has been known before that the homotopy type of $\mathcal{R}^+(M)$ only depends on the spin cobordism class $[M] \in \Omega_d^{\text{Spin}}$. More precisely, if M, N are simply connected, spin and of dimension $d \geq 5$, and $[M] = [N]$, then $\mathcal{R}^+(M) \simeq \mathcal{R}^+(N)$. This is a consequence of by Chernysh’s refinement [8] of the Gromov–Lawson surgery theorem [18], see [43, Corollary 4.2].
 - (2) There is an index-theoretic necessary condition for a closed spin manifold M^d to admit a psc metric. Namely, the \hat{a} -invariant $\hat{a}(M) \in \text{ko}_d$ has to be trivial ($\hat{a}(M)$ only depends on the spin cobordism class $[M]$ of M). A celebrated result of Stolz [37] states that this condition is also sufficient, when M is simply connected and $d \geq 5$.
 - (3) It was shown before by Kordass [26, Corollary 3.6] that $\mathcal{R}^+(\mathbb{H}\mathbb{P}^2) \simeq \mathcal{R}^+(S^8)$, giving the first example of a homotopy equivalence between spaces of psc metrics on manifolds which are not cobordant. In fact, our proof of Theorem A can be viewed as a generalization of the proof in [26], and the main technical result of [26] (Theorem 3.1 of that paper) is a major ingredient for the present paper.

Let us turn to the non-spin case. If M is simply connected, of dimension $d \geq 5$, and does not admit a spin structure, it admits a psc metric, by a result of Gromov and Lawson [18, Corollary C]. As an example for such a manifold, one can take the unit sphere bundle W^d of the unique nontrivial vector bundle of rank $d - 1$ over S^2 . There is a similar cobordism invariance theorem for the homotopy type of $\mathcal{R}^+(M)$ in that case: namely, if M, N are both simply connected and oriented, do not admit spin structures, and their cobordism classes in the oriented cobordism group Ω_d^{SO} agree, then $\mathcal{R}^+(M) \simeq \mathcal{R}^+(N)$ (see Theorem 2.3 below for a more general statement). We prove that more is true:

Theorem B. *Let M be a simply connected closed manifold of dimension $d \geq 5$ which does not admit a spin structure. Then if $d \neq 8$, there is a homotopy equivalence*

$$\mathcal{R}^+(M) \simeq \mathcal{R}^+(W^d).$$

If $d = 8$, there is either a homotopy equivalence $\mathcal{R}^+(M) \simeq \mathcal{R}^+(W^8)$ or $\mathcal{R}^+(M) \simeq \mathcal{R}^+(\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2)$.

- Remark 1.2.*
- (1) There is no reason to assume that $\mathcal{R}^+(W^d) \simeq \mathcal{R}^+(S^d)$. On the other hand, there are currently no known techniques to disprove such a statement. While $\mathcal{R}^+(S^d)$ is known to have a very rich topology by e.g. [19] and [7], nothing is known about $\mathcal{R}^+(W^d)$, except that it is nonempty.
 - (2) We conjecture that $\mathcal{R}^+(\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2) \simeq \mathcal{R}^+(W^8)$ as well; however our method misses this single case.

Theorems A and B are special cases of the following more general

Conjecture 1.3. Let M and N be two closed d -manifolds, $d \geq 5$, with the same normal 2-type. If both, M and N , admit psc metrics, then $\mathcal{R}^+(M) \simeq \mathcal{R}^+(N)$.

Here two manifolds M_i , $i = 1, 2$, are said to have the same normal 2-type, if there is a fibration $\xi : B \rightarrow BO$ such that the classifying maps $\nu_i : M_i \rightarrow BO$ of the stable normal bundles lift to maps $\bar{\nu}_i : M_i \rightarrow B$ and both $\bar{\nu}_i$ are 2-connected.

Remark 1.4.

It was observed in [14, §9] that Conjecture 1.3 is implied by the concordance–implies–isotopy conjecture for psc metrics (which, however, is completely open). See Remark 2.11 for an alternative explanation.

The method of the proof of Theorem A also proves Conjecture 1.3 for the normal 2-type $B\text{Spin} \times BG$ for some finitely presented groups, e.g. $G = \mathbb{Z}^n$ or $G = F_n$ (free group on n generators). That is, if M, N are spin with fundamental group G , and both admit psc metrics, then $\mathcal{R}^+(M) \simeq \mathcal{R}^+(N)$. See Theorem 4.5 for a more general statement.

Using the methods of this paper and computations in cobordism theory, one can prove partial results for many other normal 2-types such as BO , $B\text{SO} \times BG$ or $BO \times BG$. We refrain from stating them.

The proof of Theorem A has three major steps. The first step, carried out in §2, is a general method to construct homotopy equivalences $\mathcal{R}^+(M) \simeq \mathcal{R}^+(N)$ when M and N are not cobordant. The key result is Theorem 2.6. It asserts that if a d -dimensional spin psc manifold M admits a decomposition of a particular type (the technical name we chose for that is “admissible splitting”), then for each simply connected N in the cobordism class of M , we have $\mathcal{R}^+(N) \simeq \mathcal{R}^+(S^d)$. The second step is to show that total spaces of $\mathbb{H}\mathbb{P}^2$ -bundles with structure group $P(\text{Sp}(2) \times \text{Sp}(1))$ do admit admissible splittings. This follows from an extension of the Gromov–Lawson–Chernysh theorem that was proven by Kordaß [26] and is done in §3. The third step is to show that each simply connected spin manifold with psc metric is cobordant to the total space of an $\mathbb{H}\mathbb{P}^2$ -bundle with structure group $P(\text{Sp}(2) \times \text{Sp}(1))$. This is derived, by homotopy-theoretic methods, from the proof of Stolz’ theorem [37] in §4.

The proof of Theorem B is somewhat more convoluted. The first step applies verbatim, and the geometric argument of §3 also apply to bundles whose fibers are complex projective spaces of (complex) dimension at least 3. However, these examples do not suffice and we need a more explicit understanding of the structure of Ω_*^{SO} . Therefore, some extra arguments and a careful treatment of a particular 5-manifold are necessary. Both is done in §5. We have not been able to do the extra work which would be needed for the 8-dimensional case, which is why Theorem B remains incomplete for $d = 8$.

Acknowledgements. We would like to thank Jan-Bernhard Kordaß on discussions on generalizations of the Gromov–Lawson–Chernysh surgery construction.

J. Ebert wants to thank Michael Joachim, Christoph Böhm and Burkhard Wilking for helpful conversations about various arguments in this paper.

2. A CUT-AND-PASTE CONSTRUCTION

2.1. The cobordism theorem. We have to review the role of cobordism theory for questions about positive scalar curvature and begin with the surgery theorem of Gromov–Lawson, Schoen–Yau and Chernysh. Let us recall that a *torpedo metric* g_{tor}^k of radius δ is an $O(k)$ -invariant Riemannian metric on \mathbb{R}^k such that

- (1) in polar coordinates, g_{tor}^k is of the form $dr^2 \oplus f(r)^2 g_{S^{k-1}}$, where $0 \leq f \leq \delta$ is a smooth function with $f(0) = 0$, $f'(0) = 1$, $0 \leq f' \leq 1$, $f'' \leq 0$ and $f(t) = \delta$ for t in a neighborhood of $[1, \infty)$.
- (2) $\text{scal}(g_{\text{tor}}^k) \geq \text{scal}(\delta^2 g_{S^{k-1}}) = \frac{1}{\delta^2}(k-1)(k-2)$.

Let N be a closed manifold and let $\varphi : N \times \mathbb{R}^k \rightarrow M$ be a codimension 0 embedding. Fix a Riemannian metric g_N on N , let δ be such that $\text{scal}(g_N) + \frac{1}{\delta^2}(k-1)(k-2) > 0$ and fix a torpedo metric g_{tor}^k of radius $\delta > 0$. We denote by $\mathcal{R}^+(M, \varphi) \subset \mathcal{R}^+(M)$ the subspace of those metrics which are of the form $g_N \oplus g_{\text{tor}}^k$ on $N \times D^k$.

Theorem 2.1 (Surgery theorem). [8] *As long as $k \geq 3$, the inclusion*

$$\mathcal{R}^+(M, \varphi) \rightarrow \mathcal{R}^+(M)$$

is a weak homotopy equivalence.

Gromov–Lawson [18, Theorem A] and Schoen–Yau [36, Theorem 3] previously proved that $\mathcal{R}^+(M, \varphi) \neq \emptyset$ if $\mathcal{R}^+(M) \neq \emptyset$. A detailed account of Chernysh’s proof appears also in [12].

To use this result effectively, we have to use structured cobordism. For a manifold M , let ν_M be its normal bundle (a stable vector bundle), and let $\gamma \rightarrow BO$ be the universal stable vector bundle. Let $\xi : B \rightarrow BO$ be a fibration. By a ξ -manifold, we mean a manifold M , together with a map $\ell : M \rightarrow B$ and an isomorphism $(\xi \circ \ell)^* \gamma \cong \nu_M$ of stable vector bundles. The notion of ξ -cobordisms is defined similarly, and Ω_d^ξ is the cobordism group of d -dimensional ξ -manifolds. We say that a ξ -manifold (M, ℓ) is *relatively 2-connected* if ℓ is 2-connected. A necessary condition for the existence of a relatively 2-connected M is that B is of type (F_2) [41], and if $d \geq 4$, this is also sufficient: under those assumptions, each class $x \in \Omega_d^\xi$ contains a relatively 2-connected representative.

The reader should have the following examples in mind; in each case, B is of type (F_2) .

- Examples 2.2.*
- (1) $\xi : B\text{Spin} \rightarrow BO$. In that case, a ξ -manifold is the same as a spin manifold, and it is relatively 2-connected if and only if it is simply connected. The cobordism groups are $\Omega_d^{\text{Spin}} = \pi_d(\text{MSpin})$.
 - (2) $B\text{Spin} \times BG \rightarrow B\text{Spin} \rightarrow BO$, where G is a finitely presented group. A ξ -manifold is the same as a spin manifold equipped with a map $f : M \rightarrow BG$. The relevant cobordism group is $\Omega_d^{\text{Spin}}(BG) = \pi_d(\text{MSpin} \wedge BG_+)$, and a ξ -manifold is relatively 2-connected if and only if it is 0-connected and if f induces an isomorphism on fundamental groups.
 - (3) $B = BSO \rightarrow BO$. In that case, the relevant cobordism group is oriented cobordism $\Omega_d^{\text{SO}} = \pi_d(\text{MSO})$. An oriented manifold is relatively 2-connected if and only if it is simply connected and does not admit a spin structure.

- (4) $B = BO$. The relevant cobordism group is unoriented cobordism $\Omega_d^O = \pi_d(\text{MO})$. A manifold is relatively 2-connected if it is 0-connected, not orientable, has fundamental group $\mathbb{Z}/2$ and the universal cover does not admit a spin structure.

Theorem 2.3 (Cobordism theorem). *Let (M_i, ℓ_i) be two ξ -manifolds of dimension $d \geq 5$ and assume that $[M_0, \ell_0] = [M_1, \ell_1] \in \Omega_d^\xi$.*

- (1) *If $\mathcal{R}^+(M_0) \neq \emptyset$ and (M_1, ℓ_1) is relatively 2-connected, then $\mathcal{R}^+(M_1) \neq \emptyset$.*
- (2) *If both (M_0, ℓ_0) and (M_1, ℓ_1) are relatively 2-connected, there is a weak homotopy equivalence*

$$\mathcal{R}^+(M_0) \simeq \mathcal{R}^+(M_1).$$

This is a consequence of Theorem 2.1 and handle cancellation theory. The first part is due to Gromov and Lawson in the cases $B = B\text{Spin}$ and $B = BSO$ [18]; the general version appears in [20], Theorem 2.1.1. The second part in the case $B = B\text{Spin}$ was done in [43, Corollary 4.2]; the general case requires slightly more elaborate arguments and is carried out in [12, Theorem 1.5].

2.2. Admissible splittings. We now head towards a method which can be used to compare $\mathcal{R}^+(M_0)$ and $\mathcal{R}^+(M_1)$ when M_0 and M_1 are not cobordant. For the formulation, we shall make use of the notions of right stable and left stable psc metrics which were defined in [13, Definition 1.2.1]. To recall the definition, we have to introduce some notation and terminology. If W is a compact manifold with (collared) boundary M , we denote by $\mathcal{R}^+(W)$ the space of all psc metrics on W which are of product form near M . There is a restriction map $\text{res} : \mathcal{R}^+(W) \rightarrow \mathcal{R}^+(M)$ that takes the boundary metric, and for $h \in \mathcal{R}^+(M)$, we let $\mathcal{R}^+(W)_h := \text{res}^{-1}(h)$.

If $W : M_0 \rightsquigarrow M_1$ is a cobordism and $h_i \in \mathcal{R}^+(M_i)$, we let $\mathcal{R}^+(W)_{h_0, h_1} := \mathcal{R}^+(W)_{h_0} \amalg_{h_1}$. For each $g \in \mathcal{R}^+(W)_{h_0, h_1}$, each cobordism $V : M_1 \rightsquigarrow M_2$ and each $h_2 \in \mathcal{R}^+(M_2)$, there is a gluing map

$$\mu(g, -) : \mathcal{R}^+(V)_{h_1, h_2} \rightarrow \mathcal{R}^+(W \cup V)_{h_0, h_2} \quad \mu(g, h) := g \cup h,$$

and we say that g is *right stable* if $\mu(g, -)$ is a weak equivalence for all such V and h_2 . Left stability is formulated with the analogous gluing map $\mu(-, g)$ defined for cobordisms with outgoing boundary M_0 . A metric is *stable* if it is both, left and right stable. We remark that g is right stable if and only if it is left stable when considered as a metric on the reversed cobordism. As an example, consider $N \times D^k : \emptyset \rightsquigarrow N \times S^{k-1}$ for $k \geq 3$. If g_N and g_{tor}^k are as in Theorem 2.1, that result might be restated by saying that $g_N \oplus g_{\text{tor}}^k \in \mathcal{R}^+(N \times D^k)_{g_N \oplus g_0}$ is right stable.

Definition 2.4. Let M be a closed d -manifold. An *admissible splitting* of M is a decomposition of M as the concatenation of two cobordisms

$$\emptyset \xrightarrow{M_0} N \xrightarrow{M_1} \emptyset,$$

such that the following conditions hold:

- (1) there are psc metrics $h_i \in \mathcal{R}^+(N)$ and $g_i \in \mathcal{R}^+(M_i)_{h_i}$, such that h_0 and h_1 lie in the same path component of $\mathcal{R}^+(N)$,
- (2) g_0 is right stable and g_1 is left stable,
- (3) the inclusion maps $N \rightarrow M_i$ are 2-connected.

It is often more convenient to write such a splitting as $M = M_0 \cup_N M_1$. We remark that if M has an admissible splitting, it automatically admits a psc metric (this is because we can find a psc metric in $\mathcal{R}^+(N \times [0, 1])_{h_0, h_1}$ if h_0 and h_1 lie in the same path component).

Examples 2.5. (1) The splitting $S^d = D^d \cup_{S^{d-1}} D^d$ is admissible when $d \geq 3$: take $g_0 = g_1 = g_{\text{tor}}^d$.
 (2) The standard splitting $S^d = (S^{k-1} \times D^{d-k+1}) \cup_{S^{k-1} \times S^{d-k}} (D^k \times S^{d-k})$ is admissible if $3 \leq k \leq d-2$: take $g_0 = g_{S^{k-1}} \oplus g_{\text{tor}}^{d-k+1}$ and $g_1 = g_{\text{tor}}^k \oplus g_{S^{d-k}}$.
 (3) Let P be a d -manifold, $d \geq 6$, such that $\partial P \rightarrow P$ is 2-connected. Then by [13, Theorem D], there is $g_{\partial P} \in \mathcal{R}^+(\partial P)$ and a right stable $g_P \in \mathcal{R}^+(P)_{g_{\partial P}}$. The decomposition $dP = P \cup_{\partial P} P^{op}$ of the double of P is admissible.

All these examples are nullbordant; and in fact the key point of this work is to construct admissible splittings of manifolds which are not nullbordant. Let us first see how admissible splittings are helpful for proving our main theorems.

Theorem 2.6. *Let $d \geq 5$ and let $\xi : B \rightarrow BO$ be a fibration. Let $x \in \Omega_d^\xi$ be a cobordism class which contains a representative M that has an admissible splitting. Let K and L be relatively 2-connected ξ -manifolds such that*

$$[L] = [K] + x \in \Omega_d^\xi.$$

Then there is a weak equivalence

$$\mathcal{R}^+(L) \simeq \mathcal{R}^+(K).$$

It is obvious that the set of cobordism classes $x \in \Omega_d^\xi$ which contain a representative with an admissible splitting is an additive subgroup. For the proof of Theorem 2.6, we need a lemma.

Lemma 2.7. *Let $M : \emptyset \rightsquigarrow N$ be a cobordism, let $g_0 \in \mathcal{R}^+(M)$ be right stable and put $h_0 := g_0|_N$. Let $h_1 \in \mathcal{R}^+(N)$ be any other psc metric in the path component of h_0 . Then there is a right stable $g_1 \in \mathcal{R}^+(M)$ with $g_1|_N = h_1$.*

Proof. The restriction map $\text{res} : \mathcal{R}^+(M) \rightarrow \mathcal{R}^+(N)$ is a Serre fibration by [12, Theorem 1.1] (which is an improvement of the main result of [9]). Therefore, we can lift any path $t \mapsto h_t$ connecting h_0 and h_1 in $\mathcal{R}^+(N)$ to a path $t \mapsto g_t$ in $\mathcal{R}^+(M)$, starting at g_0 . By [14, Lemma 2.3.4], g_1 is right stable. \square

Proof of Theorem 2.6. Because of Lemma 2.7, we may assume that $h_0 = h_1 =: h$.

Without loss of generality, B is of type (F_2) ; otherwise, the Proposition holds vacuously. Let $M = M_0 \cup_N M_1$ be an admissible splitting of M , with psc metrics $g_0 \in \mathcal{R}^+(M_0)_h$ and $g_1 \in \mathcal{R}^+(M_1)_h$ as in Definition 2.4. Denote by M'_0 disjoint union of K with M_0 . Since B is of type (F_2) , we can turn M'_0 by a sequence of ξ -surgeries of index ≤ 2 in its interior to a relatively 2-connected ξ -manifold M''_0 with boundary N . The ξ -manifold $M' := M''_0 \cup_N M_1$ lies in the cobordism class $x + [K]$, and is therefore ξ -cobordant to L .

We show below that M' is relatively 2-connected. Taking this for granted, Theorem 2.3 shows that there is a weak equivalence

$$(2.8) \quad \mathcal{R}^+(L) \simeq \mathcal{R}^+(M').$$

Let further $M'' := M'_0 \cup_N M_0^{op}$. Since g_1 and g_0^{op} are left stable, there are homotopy equivalences

$$(2.9) \quad \mathcal{R}^+(M') \xleftarrow{\mu(\cdot, g_1)} \mathcal{R}^+(M'_0)_h \xrightarrow{\mu(\cdot, g_0^{op})} \mathcal{R}^+(M''),$$

given by gluing in g_1 and g_0^{op} respectively. On the other hand

$$[M''] = [K \# (M_0 \cup_N M_0^{op})] = [K] \in \Omega_d^\xi;$$

the second equality holds since the double of M_0 is nullbordant.

We show below that M'' is relatively 2-connected. Taking this for granted, Theorem 2.3 shows that there is a weak equivalence

$$(2.10) \quad \mathcal{R}^+(M'') \simeq \mathcal{R}^+(K),$$

and combining (2.8), (2.9) and (2.10) will finish the proof.

It remains to be shown that M' and M'' are relatively 2-connected. Consider the commutative diagram

$$\begin{array}{ccc} M'_0 & \xrightarrow{a} & M' \\ \downarrow b & & \downarrow \ell_{M'} \\ M'' & \xrightarrow{\ell_{M''}} & B. \end{array}$$

The map $\ell_{M''} = \ell_{M'} \circ a = \ell_{M''} \circ b$ is 2-connected. Furthermore, since $N \rightarrow M_i$ is 2-connected, it follows by excision (or general position) that a and b are 2-connected. Therefore $\ell_{M'}$ and $\ell_{M''}$ are 2-connected as well. \square

Remark 2.11. Let M be an arbitrary closed d -manifold with $d \geq 5$. Using Morse theory, we can write M as composition $\emptyset \xrightarrow{M_0} N \xrightarrow{M_1} \emptyset$, where both inclusions $N \rightarrow M_i$ are 2-connected. Using [13, Theorem D], there are $h_0, h_1 \in \mathcal{R}^+(N)$, a right stable $g_0 \in \mathcal{R}^+(M_0)_{h_0}$ and a left stable $g_1 \in \mathcal{R}^+(M_1)_{h_1}$. If M admits a psc metric, the metrics h_0, h_1 are concordant. If we could infer from that information that h_0 and h_1 are isotopic (i.e. in the same path component), we would get an admissible splitting.

In other words, an affirmative solution of the concordance–implies–isotopy conjecture (which is wide open of course) would imply immediately the truth of Conjecture 1.3.

On the other hand, this observation means that when constructing admissible splittings, the key point is always to ensure that h_0 and h_1 are in the same path component.

3. ADMISSIBLE SPLITTINGS OF PROJECTIVE BUNDLES

To apply Theorem 2.6, we need a sufficient supply of manifolds which have admissible splittings (note that all of the examples listed in (2.5) are nullbordant). The foremost goal of this section is the following result.

Theorem 3.1. *Let $E \rightarrow B$ be a smooth fibre bundle with fibre $\mathbb{K}\mathbb{P}^{m+n-1}$ over a closed manifold B (here $\mathbb{K} = \mathbb{R}, \mathbb{C}$ or \mathbb{H}). Then E has an admissible splitting in each of the following cases.*

- (1) $\mathbb{K} = \mathbb{R}$, $m, n \geq 3$, and the structure group is $P(O(m) \times O(n)) = (O(m) \times O(n)) / \{\pm 1\}$.
- (2) $\mathbb{K} = \mathbb{C}$, $m, n \geq 2$, and the structure group is $U(m) \times U(n)$.

- (3) $\mathbb{K} = \mathbb{C}$, $m + n - 1 \geq 3$, and the structure group is C_2 , acting by complex conjugation.
- (4) $\mathbb{K} = \mathbb{H}$, $m, n \geq 1$, and the structure group is $P(\mathrm{Sp}(m) \times \mathrm{Sp}(n)) = (\mathrm{Sp}(m) \times \mathrm{Sp}(n))/\{\pm 1\}$.

This goal is achieved in §3.4, where we also introduce more uniform notation. The key to this result is a generalization of the surgery theorem to embeddings with nontrivial normal bundle, which was proven by Kordas in [26]. Precise statements are the key, and to that end, we need to review some basic material from Riemannian geometry.

3.1. Connection metrics.

Assumption 3.2. Let G be a Lie group with Lie algebra \mathfrak{g} and let $p : P \rightarrow M$ be a smooth G -principal bundle, equipped with a connection θ . Assume further that F is a manifold with a left G -action. Finally, let g_M be a Riemannian metric on M and let g_F be a G -invariant Riemannian metric on F .

Out of these data, we produce a Riemannian metric $g_M \oplus_\theta g_F$ on the Borel construction $P \times_G F$. The construction is as follows.

The datum of θ can be given by a certain 1-form $\theta \in \mathcal{A}^1(P; \mathfrak{g})$, or more geometrically by one of the following equivalent gadgets:

- (1) a G -equivariant left inverse V of the inclusion map $T_v P := \ker(dp) \rightarrow TP$ of the vertical tangent bundle,
- (2) or by a G -invariant complementary subbundle $T_h P \subset TP$ to $T_v P$, the *horizontal tangent bundle*.

Given V , one puts $T_h P := \ker(V)$ (this is also the same as $\ker(\theta)$). Given a decomposition $TP = T_v P \oplus T_h P$, one lets V be the projection onto the first summand. There is also the horizontal projection $H : TP \rightarrow T_h P$; both V and H are G -equivariant. The differential of p induces a G -equivariant isomorphism $T_h P \cong p^*TM$.

Let $q : P \times F \rightarrow P \times_G F$ be the quotient map, and let $\pi : P \times_G F \rightarrow M$ be the projection to M . The subbundle $T_h P \times 0 \subset T(P \times F)$ projects to a subbundle

$$q_*(T_h P \times 0) =: T_h(P \times_G F) \subset T(P \times_G F).$$

Furthermore $d\pi$ restricts to a bundle isomorphism

$$T_h(P \times_G F) \rightarrow \pi^*TM,$$

and the kernel of $d\pi$ is the vertical tangent bundle $T_v(P \times_G F) = P \times_G TF$. Together, we obtain an isomorphism

$$(3.3) \quad \pi^*TM \oplus T_v(P \times_G F) \cong T(P \times_G F)$$

which only depends on θ .

Definition 3.4. Let G, P, M, F, θ, g_M and g_F as in 3.2. The *connection metric* $g_M \oplus_\theta g_F$ is the Riemannian metric on $P \times_G F$ which under the isomorphism (3.3) corresponds to the direct sum of π^*g_M and the bundle metric on $T_v(P \times_G F) = P \times_G TF$ induced from the G -equivariant metric g_F .

Lemma 3.5. [17, Proposition 2.7.1] *The map*

$$\pi : (P \times_G F, g_M \oplus_\theta g_F) \rightarrow (M, g_M)$$

is a Riemannian submersion with totally geodesic fibres.

The most common case arises from a rank r Riemannian vector bundle $V \rightarrow M$ with a metric connection ω and an $O(r)$ -invariant metric g^r on \mathbb{R}^r : the metric connection induces a connection on the $O(r)$ -principal bundle $\text{Fr}(V)$ of orthonormal frames of V .

Let us record an obvious, but useful naturality property of connection metrics.

Lemma 3.6. *Let G, P, M, F, θ, g_M and g_F as in 3.2.*

- (1) *Let H be a further Lie group which acts from the left on P and M , compatible with the G -action on P , the map p , the Riemannian metric g_M and the connection θ . Then the connection metric $g_M \oplus_\theta g_F$ is H -invariant.*
- (2) *If K is a Lie group that acts on F , commuting with the G -action and leaving g_F invariant. Then $g_M \oplus_\theta g_F$ is K -invariant.*

Assume the situation of (1), let $q : Q \rightarrow N$ be a smooth H -principal bundle with a connection ω . In this situation, we can form the associated bundle

$$Q \times_H (P \times_G F) = (Q \times_H P) \times_G F \rightarrow N$$

with fibre $P \times_G F$. Note that $Q \times_H P$ is a G -principal bundle over $Q \times_H M$, and it has an induced connection $Q \times_H \theta$, given by the following recipe. The subbundle $T_h Q \oplus T_h P \subset T(P \times Q)$ is $H \times G$ -invariant and projects to an G -invariant subbundle $T_h(Q \times_H P) \subset T(Q \times_H P)$ which is complementary to $T_v(Q \times_H P)$.

Lemma 3.7. *In the situation just described, the two metrics $(g_N \oplus_\omega g_M) \oplus_{Q \times_H \theta} g_F$ and $g_N \oplus_\omega (g_M \oplus_\theta g_F)$ on $(Q \times_H P) \times_G F$ agree.*

Proof. Let $r : Q \times P \times F \rightarrow Z := Q \times_H P \times_G F$ be the quotient map. It induces $G \times H$ -equivariant isomorphisms

$$q^*TN \times p^*TM \times TF \cong T_h Q \times T_h P \times TF \cong r^*TZ.$$

Under this isomorphism, both connection metrics on Z correspond to $g_N \oplus g_M \oplus g_F$. \square

3.2. The curvature of connection metrics. We assume the notation introduced in §3.1, and compute the scalar curvature of the connection metric $g_M \oplus_\theta g_F$ with the aid of O'Neill's formulas [32], see also [6], Chapter 9. For a general Riemannian submersion $\pi : E \rightarrow M$, O'Neill defines two tensor fields $A, T \in \Gamma(E, T^*E^{\otimes 2} \otimes TE)$. Because the fibres of π are totally geodesic in our situation, the tensor field T vanishes [6, §9.26] and can be ignored. The A -tensor is defined by the formula (here e, f are vector fields on E)

$$A_e f = V \nabla_{H_e}(Hf) + H \nabla_{H_e}(Vf).$$

Lemma 3.8. (1) *If x, y are horizontal, $A_x y = \frac{1}{2}V[x, y]$. In particular, if $E = P \times_G F$, $A_x y$ only depends on the connection θ and not on the involved metrics.*

(2) *If x is horizontal and v is vertical, $A_x v$ is horizontal.*

(3) *If x, y are horizontal and v is vertical, we have*

$$(3.9) \quad \langle A_x v, y \rangle + \langle v, A_x y \rangle = 0.$$

Proof. The first formula is [32, Lemma 2], and the second claim is clear. The third follows from

$$\langle A_x v, y \rangle + \langle v, A_x y \rangle = \langle H(\nabla_x v), y \rangle + \langle v, V(\nabla_x y) \rangle = \langle \nabla_x v, y \rangle + \langle v, \nabla_x y \rangle = x \langle v, y \rangle = 0.$$

\square

For two horizontal vectors, we can express $A_x y$ in terms of the curvature of θ , as follows. For $X \in \mathfrak{g}$, let $\rho(X)$ be the vector field on F given by differentiating the G -action, and let $\Omega \in \mathcal{A}^2(P; \mathfrak{g})$ be the curvature form of the connection θ . For $q \in F$, let $\varphi : P \rightarrow P \times_G F$ be the map $p \mapsto [(p, q)]$.

Let $x, y \in T_{[(p, q)]}(P \times_G F)$ be two horizontal vectors. By the definition of the horizontal distribution, we find (unique) horizontal tangent vectors $x_*, y_* \in T_p P$, such that $(T\varphi)x_* = x$ and $(T\varphi)y_* = y$.

Using the formula [11, p. 51] for the evaluation of Ω on horizontal vectors and [32, Lemma 2], we obtain the formula

$$(3.10) \quad A_x y = -\rho(\Omega(x_*, y_*))_q.$$

Now let \sec_M, \sec_F and \sec be the sectional curvature of g_M, g_F and $g_M \oplus_\theta g_F$, respectively. If v, w are orthonormal vertical vectors and x, y are orthonormal horizontal vectors, we have the following identities, by [32, Corollary 1]:

$$(3.11) \quad \begin{aligned} \sec(v, w) &= \sec_F(v, w), \\ \sec(x, v) &= \|A_x v\|^2, \\ \sec(x, y) &= \sec_M(x, y) - 3\|A_x y\|^2. \end{aligned}$$

Now we can calculate the scalar curvature of $g_M \oplus_\theta g_F$. Let (x_i) be an orthonormal basis for the horizontal vectors, and (f_j) an orthonormal basis for the vertical vectors, both at a fixed point of $P \times_G F$. Then at this point

$$\begin{aligned} \text{scal}(g_M \oplus_\theta g_F) &= 2 \sum_{i < j} \sec(x_i, x_j) + 2 \sum_{i, j} \sec(x_i, f_j) + 2 \sum_{i < j} \sec(f_i, f_j) = \\ &= \text{scal}(g_M) - 6 \sum_{i < j} \|A_{x_i} x_j\|^2 + 2 \sum_{i, j} \|A_{x_i} f_j\|^2 + \text{scal}(g_F). \end{aligned}$$

Since $A_{x_i} f_j$ is horizontal, its norm square is the same as

$$\begin{aligned} \sum_{i, j} \|A_{x_i} f_j\|^2 &= \sum_{i, j, k} \langle A_{x_i} f_j, x_k \rangle^2 \stackrel{3.9}{=} \\ &= \sum_{i, j, k} \langle f_j, A_{x_i} x_k \rangle^2 = \text{(since } A_{x_i} x_j \text{ is vertical)} \\ &= \sum_{i, k} \|A_{x_i} x_k\|^2 = \text{(by Lemma 3.8 (1))} \\ &= 2 \sum_{i < k} \|A_{x_i} x_k\|^2. \end{aligned}$$

Altogether, we obtain

$$(3.12) \quad \text{scal}(g_M \oplus_\theta g_F) = \text{scal}(g_M) + \text{scal}(g_F) - |A|^2,$$

where we denote

$$|A|^2 := 2 \sum_{i < j} \|A_{x_i} x_j\|^2 \geq 0.$$

For $t > 0$, we can scale the metric g_F to $t g_F$. This does not change the orthonormal basis (x_i) , since g_M is unchanged; so $A_{x_i} x_j$ is leaved unchanged, but its norm square gets scaled by t . Moreover, $\text{scal}(t g_F) = \frac{1}{t} \text{scal}(g_F)$. Altogether, we get

$$(3.13) \quad \text{scal}(g_M \oplus_\theta t g_F) = \text{scal}(g_M) + \frac{1}{t} \text{scal}(g_F) - t|A|^2.$$

More generally, with an additional scaling parameter, we have

$$(3.14) \quad \text{scal}(sg_M \oplus_\theta tg_F) = \frac{1}{s} \text{scal}(g_M \oplus_\theta \frac{t}{s} g_F) = \frac{1}{s} \text{scal}(g_M) + \frac{1}{t} \text{scal}(g_F) - \frac{t}{s^2} |A|^2.$$

3.3. The generalized Gromov–Lawson–Chernysh theorem. We can now state the generalized Gromov–Lawson–Chernysh surgery theorem [26, Theorem 3.1] (see [36, Theorem 3] for a precursor). Using the notion of right stability, the result might be stated as follows.

Theorem 3.15. [26, Theorem 3.1] *Let $V \rightarrow N$ be a Riemannian vector bundle of rank $r \geq 3$ over a closed manifold N with unit disc bundle $D(V)$ and unit sphere bundle $S(V)$, let ω be a metric connection on V and let h be a Riemannian metric on N . If the connection metric $g = h \oplus_\omega g_{\text{tor}}^r$ has positive scalar curvature, where g_{tor}^r is a torpedo metric, then g is right stable on $D(V): \emptyset \rightsquigarrow S(V)$.*

The following consequence of Theorem 3.15 is our main tool to construct admissible splittings.

Proposition 3.16. *Assume that*

- (1) $V_i \rightarrow N_i$, $i = 0, 1$, are Riemannian vector bundles over closed manifolds, of rank $r_i := \text{rank}(V_i) \geq 3$,
- (2) $\psi : S(V_0) \cong S(V_1)$ is a diffeomorphism of the sphere bundles, and $M := D(V_0) \cup_\psi D(V_1)^{\text{op}}$,
- (3) G is a Lie group which acts smoothly on N_i , and on V_i by bundle isometries, such that ψ is G -equivariant (hence G acts on M),
- (4) ω_i , $i = 0, 1$, are G -equivariant metric connections on V_i and h_i are G -invariant Riemannian metric on N_i , and the (G -invariant) connection metrics $g_i := h_i \oplus_{\omega_i} g_{\text{tor}}^{r_i}$ have positive scalar curvature (for suitable torpedo metrics $g_{\text{tor}}^{r_i}$).
- (5) $\psi^* g_1|_{S(V_1)}$ and $g_0|_{S(V_0)}$ lie in the same path component of $\mathcal{R}^+(S(V_0))^G$, the space of G -invariant psc metrics on $S(V_0)$.
- (6) Let finally $P \rightarrow B$ be a smooth G -principal bundle over a closed manifold.

Then the manifold $P \times_G M$ has an admissible splitting.

Proof. We can write

$$(3.17) \quad P \times_G M = (P \times_G D(V_0)) \cup_{P \times_G \psi} (P \times_G D(V_1)^{\text{op}}).$$

The inclusions $S(V_i) \hookrightarrow D(V_i)$ are 2-connected because $r_i \geq 3$, and hence so are the inclusions $P \times_G S(V_i) \rightarrow P \times_G D(V_i)$.

We can write

$$P \times_G V_i = P \times_G (\text{Fr}(V_i) \times_{O(r_i)} \mathbb{R}^{r_i}) = (P \times_G \text{Fr}(V_i)) \times_{O(r_i)} \mathbb{R}^{r_i}.$$

Choose a Riemannian metric g_B on B and a connection θ on $P \rightarrow B$. We consider the metrics

$$(3.18) \quad m_{i,u} := (g_B \oplus_\theta u h_i) \oplus_{P \times_\theta \omega_i} u g_{\text{tor}}^{r_i} \stackrel{3.7}{=} g_B \oplus_\theta (u h_i \oplus_{\omega_i} u g_{\text{tor}}^{r_i}) = g_B \oplus_\theta u g_i.$$

on $P \times_G V_i$ for a scaling parameter $u > 0$. By the scaling identity (3.13), there is u_0 , such that for all $0 < u \leq u_0$, $\text{scal}(m_{i,u}) > 0$ (look at the right hand side of (3.13)).

For such u , $m_{i,u}$ lies in the same path component of $\mathcal{R}^+(P \times_G D(V))$ as

$$\frac{1}{u} m_{i,u} = \left(\frac{1}{u} g_B \oplus_\theta h_i \right) \oplus_{P \times_\theta \omega_i} g_{\text{tor}}^{r_i},$$

which is right stable by Theorem 3.15 (where $P \times_G D(V)$ is viewed as a cobordism from \emptyset to $P \times_G S(V)$). Using [14, Lemma 2.3.4], it follows that $m_{i,u}$ is right stable.

Let $k_i := g_i|_{S(V_i)}$ and let $\gamma : [0, 1] \rightarrow \mathcal{R}^+(S(V_0))^G$ be a path connecting k_0 and ψ^*k_1 . Without loss of generality, we can assume that u_0 is small enough so that $\text{scal}(g_B \oplus u\gamma(t)) > 0$ for all $0 < u \leq u_0$ and $t \in [0, 1]$.

For such a choice of u , we have a right stable metric $m_{0,u}$ on $Q_0 := P \times_G D(V_0)$, a left stable metric $m_{1,u}^{op}$ on $Q_1 := P \times_G D(V_0)$ and a path $\gamma' : [0, 1] \rightarrow \mathcal{R}^+(P \times_G S(V_0))$ from $m_{0,u}|_{P \times_G S(V_0)}$ to $(P \times_G \psi)^*m_{1,u}$. This completes the proof. \square

3.4. Construction of the splittings. In this subsection, we finish the proof of Theorem 3.1. Our proof will also cover bundles of real projective space. Let us first introduce uniform notation for the real, complex and quaternionic case.

- (1) \mathbb{K} is either \mathbb{R} , \mathbb{C} or \mathbb{H} , and $k := \dim_{\mathbb{R}}(\mathbb{K})$. We denote by $\mathfrak{S}(\mathbb{K}) \subset \mathbb{K}$ the imaginary part, which we identify with the Lie algebra of the group $S^{k-1} \subset \mathbb{K}^\times$.
- (2) We consider \mathbb{K}^n as a *right* \mathbb{K} -vector space, equipped with the standard inner product. The group S^{k-1} acts on \mathbb{K}^n and $S(\mathbb{K}^n) = S^{kn-1}$ by right multiplication; let R_z denote the right multiplication by $z \in S^{k-1}$.
- (3) We denote by

$$G(n) := N_{O(kn)}(S^{k-1})$$

the normalizer subgroup of S^{k-1} in $O(kn)$. The centralizer

$$H(n) := C_{O(kn)}(S^{k-1}) \subset G(n)$$

is equal to $O(n), U(n), \text{Sp}(n)$, depending on the case at hand. The action of $H(n)$ on \mathbb{K}^n by left matrix multiplication commutes with the S^{k-1} -action, and we let L_A be the left multiplication by $A \in H(n)$.

- (4) The quotient map $p : S^{kn-1} \rightarrow \mathbb{K}\mathbb{P}^{n-1} := S^{kn-1}/S^{k-1}$ to the projective space is an S^{k-1} -principal bundle. The projective space has an induced action of the normalizer $G(n)$, and the kernel of that $G(n)$ -action is S^{k-1} , so that the group

$$PG(n) := G(n)/S^{k-1}$$

acts (faithfully) on $\mathbb{K}\mathbb{P}^{n-1}$.

- (5) More generally, we consider the subgroup $O(kn) \times O(km) \subset O(k(m+n))$, and define

$$G(n, m) := N_{O(kn) \times O(km)}(S^{k-1}) = G(n+m) \cap (O(kn) \times O(km)),$$

$$H(n, m) := C_{O(kn) \times O(km)}(S^{k-1}) = H(n) \times H(m) \subset G(n, m)$$

and

$$PG(n, m) := G(n, m)/S^{k-1} \subset PG(n+m).$$

- (6) The action of $G(n)$ on S^{kn-1} is effective and transitive with stabilizer $\text{Stab}_{G(n)}(e_n) = G(n-1)$. The action of $PG(n)$ on $\mathbb{K}\mathbb{P}^{n-1}$ is transitive, and the stabilizer of $[e_n] \in \mathbb{K}\mathbb{P}^{n-1}$ is equal to $PG(n-1, 1)$.
- (7) The structure of the group $G(n)$ is determined as follows. There is an exact sequence

$$1 \rightarrow H(n) \rightarrow G(n) \rightarrow \text{Aut}(S^{k-1}) \rightarrow 1$$

in each case. It follows that $G(n) = O(n)$ if $\mathbb{K} = \mathbb{R}$, that $G(n) = C_2 \times U(n)$ in the complex case, where the cyclic group C_2 acts by complex conjugation, and that in the quaternionic case $G(n) = \text{Sp}(n) \times_{\pm 1} \text{Sp}(1)$, where (A, z) act

by $L_A R_z$. The quotient groups are $PG(n) = O(n)/\pm 1$ in the real case, $PG(n) = C_2 \times U(n)/S^1$ in the complex case and $PG(n) = Sp(n)/\pm 1 = PSp(n)$ in the quaternionic case.

The goal of this section is the proof of the following result.

Proposition 3.19. *Let $E \rightarrow B$ be a smooth fibre bundle with fibre $\mathbb{K}\mathbb{P}^{m+n-1}$ and structure group $PG(n, m)$ over a closed manifold B . Then E has an admissible splitting in each of the following cases.*

- (1) $\mathbb{K} = \mathbb{R}$, $n, m \geq 3$,
- (2) $\mathbb{K} = \mathbb{C}$, $n, m \geq 2$,
- (3) $\mathbb{K} = \mathbb{H}$, $n, m \geq 1$.

Proposition 3.19 implies Theorem 3.1, because each of the structure groups mentioned in that theorem act through its image in $PG(n, m)$.

We start the proof of Proposition 3.19 with a suitable decomposition of $\mathbb{K}\mathbb{P}^{n+m-1}$ into disc bundles which is invariant under $PG(n, m)$. Let

$L_0 := \{(v, w) \in S(\mathbb{K}^n \times \mathbb{K}^m) \mid \|v\| \geq \|w\|\}$, $L_1 := \{(v, w) \in S(\mathbb{K}^n \times \mathbb{K}^m) \mid \|v\| \leq \|w\|\}$ with intersection

$$L_{01} := L_0 \cap L_1 = \{(v, w) \mid \|v\|^2 = \|w\|^2 = \frac{1}{2}\} \cong S(\mathbb{K}^n) \times S(\mathbb{K}^m).$$

The decomposition $S^{kn+km-1} = L_0 \cup L_1$ is invariant under the action of $O(kn) \times O(km)$ and of S^{k-1} , and hence in particular under $G(n, m)$. It therefore induces a decomposition $\mathbb{K}\mathbb{P}^{n+m-1} = P_0 \cup_{P_{01}} P_1$, where $P_i := L_i/S^{k-1}$, which is invariant under $PG(n, m)$. The map

$$\varphi_0 : S(\mathbb{K}^n) \times_{S^{k-1}} D(\mathbb{K}^m) \rightarrow \mathbb{K}\mathbb{P}^{n+m-1}; [v, w] \mapsto \left[\frac{1}{\sqrt{2}}(v, w)\right]$$

is a diffeomorphism onto P_0 , and there is a similar diffeomorphism

$$\varphi_1 : S(\mathbb{K}^m) \times_{S^{k-1}} D(\mathbb{K}^n) \cong P_1.$$

The diffeomorphisms φ_i are $PG(n, m)$ -equivariant. Altogether, we have found a $PG(n, m)$ -invariant decomposition

$$(3.20) \quad \mathbb{K}\mathbb{P}^{n+m-1} = D(S(\mathbb{K}^n) \times_{S^{k-1}} \mathbb{K}^m) \cup_{\varphi_1^{-1} \circ \varphi_0} D(S(\mathbb{K}^m) \times_{S^{k-1}} \mathbb{K}^n)$$

into disc bundles. If n and m are as in Proposition 3.19, these disc bundles have rank at least 3. The action of $PG(n, m)$ on the sphere bundle $S(S(\mathbb{K}^n) \times_{S^{k-1}} \mathbb{K}^m)$ is transitive. This sphere bundle is more conveniently written as

$$M_{k,n,m} := S^{kn-1} \times_{S^{k-1}} S^{km-1}.$$

Note that $M_{k,n,m}$ is both, an S^{km-1} -bundle over $\mathbb{K}\mathbb{P}^{n-1}$ and an S^{kn-1} -bundle over $\mathbb{K}\mathbb{P}^{m-1}$. The action of the group $PG(n, m)$ on $M_{k,n,m}$ is transitive, because the action of $H(n, m)$ on $S^{kn-1} \times S^{km-1}$ is already transitive.

Let $g_{S^{kn-1}}$ be the round metric on S^{kn-1} . The action of $O(kn)$ and hence of $G(n)$ is by isometries. The action of the subgroup $S^{k-1} \subset G(n)$ is geodesic in the sense that orbits of 1-parameter subgroups of S^{k-1} are geodesics in S^{kn-1} . We let $T_v S^{kn-1}$ be the $(k-1)$ -dimensional distribution of tangent vectors to orbits. The distribution $T_v S^{kn-1}$ is preserved by $G(n)$, and so is its orthogonal complement $T_h S^{kn-1} := (T_v S^{kn-1})^\perp$. Moreover $T_h S^{kn-1}$ is the horizontal distribution for a

principal connection ω_n on the S^{k-1} -principal bundle $p : S^{kn-1} \rightarrow \mathbb{K}\mathbb{P}^{n-1}$. In fact, ω_n is $G(n)$ -equivariant, and it is the only connection with this property.

The *Fubini-Study metric* on $\mathbb{K}\mathbb{P}^{n-1}$ is the quotient metric $g_{\mathbb{K}\mathbb{P}^{n-1}}$ of the round metric. The map $\pi : (S^{kn-1}, g_{S^{kn-1}}) \rightarrow (\mathbb{K}\mathbb{P}^{n-1}, g_{\mathbb{K}\mathbb{P}^{n-1}})$ is a Riemannian submersion with totally geodesic fibres, and $g_{S^{kn-1}}$ can be identified with the connection metric

$$g_{S^{kn-1}} = g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{S^{k-1}}$$

(since both, vertical and horizontal distributions are orthogonal for both metrics, and both metrics agree on horizontal and vertical vectors).

The metric $g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} tg_{\text{tor}}^{km}$ on $S^{kn-1} \times_{S^{k-1}} D^{km} = P_0$ is $PG(n, m)$ -invariant, and similarly is the metric $g_{\mathbb{K}\mathbb{P}^{m-1}} \oplus_{\omega_m} sg_{\text{tor}}^{kn}$ on $S^{km-1} \times_{S^{k-1}} D^{kn} = P_1$. For t, s sufficiently small, these have positive scalar curvature (provided n and m are as in Proposition 3.19). By Proposition 3.16, the proof of Proposition 3.19 is complete, once we can show that for suitable s and t , the two metrics

$$g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} tg_{S^{km-1}}; \quad g_{\mathbb{K}\mathbb{P}^{m-1}} \oplus_{\omega_m} sg_{S^{kn-1}} \in \mathcal{R}^+(M_{k,n,m})^{PG(n,m)}$$

can be connected by a path of $PG(n, m)$ -invariant psc metrics. Therefore, all that remains to be shown is:

Lemma 3.21. *The space of $PG(n, m)$ -invariant psc metrics on $M_{k,n,m}$ is contractible, if $k = 1$ and $m, n \geq 3$, or $k = 2$ and $m, n \geq 2$, or $k = 4$ and $m, n \geq 1$.*

The proof of Lemma 3.21 will be quite direct, but a bit lengthy, and we begin with some preliminaries.

Definition 3.22. The *Berger metric* $g_{s,t}^{k,n}$ on S^{kn-1} with parameters $s, t > 0$ is

$$g_{s,t}^{k,n} := sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} tg_{S^{k-1}}.$$

The Berger metrics are $G(n)$ -invariant, and are the only such metrics (we shall not need this fact explicitly). The metric

$$sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$$

on $M_{k,n,m}$ is $PG(n, m)$ -invariant, and similarly is

$$sg_{\mathbb{K}\mathbb{P}^{m-1}} \oplus_{\omega_m} g_{u,t}^{k,n}.$$

In fact, it is not hard to see that

$$sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m} = ug_{\mathbb{K}\mathbb{P}^{m-1}} \oplus_{\omega_m} g_{s,t}^{k,n},$$

but we do not make use of this fact.

Lemma 3.23. *All $PG(n, m)$ -invariant metrics on $M_{k,n,m}$ are of the form $sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$.*

Proof. The map

$$q : M_{k,n,m} \rightarrow \mathbb{K}\mathbb{P}^{n-1} \times \mathbb{K}\mathbb{P}^{m-1}$$

is an S^{k-1} -principal bundle and $PG(n, m)$ -equivariant. The tangent bundle of $M_{k,n,m}$ can be decomposed into three summands:

$$M_{k,n,m} = T_v M_{k,n,m} \oplus q^*(T\mathbb{K}\mathbb{P}^{n-1} \oplus 0) \oplus q^*(0 \oplus T\mathbb{K}\mathbb{P}^{m-1}),$$

and this decomposition is $PG(n, m)$ -equivariant. We shall now argue that any $PG(n, m)$ -invariant symmetric bilinear form on $TM_{k,n,m}$ is the direct sum of invariant bilinear forms on the three summands, and that each summand carries a

unique (up to scalar multiple) such invariant symmetric bilinear form. From this, the claim follows by counting parameters (for that, note that the construction of $sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$ can be carried out for arbitrary real s, u, t and gives an invariant bilinear form).

For that goal, pick the basepoint $[e_n, e_m] \in M_{k,n,m}$ and let $K(n, m) \subset PG(n, m)$ be the stabilizer subgroup at $[e_n, e_m]$. Because the action of $PG(n, m)$ on $M_{k,n,m}$ is transitive, $PG(n, m)$ -invariant bilinear forms on $M_{k,n,m}$ correspond bijectively to $K(n, m)$ -invariant bilinear forms on the tangent space $T_{[e_n, e_m]}M_{k,n,m}$. The stabilizer $K(n, m)$ is equal to

$$\begin{aligned} & \{(A, z, B, z) \in O(n-1) \times O(1) \times O(m-1) \times O(1)\} / \{\pm 1\} & \mathbb{K} = \mathbb{R}, \\ C_2 \times & \{(A, z, B, z) \in U(n-1) \times U(1) \times U(m-1) \times U(1)\} / U(1) & \mathbb{K} = \mathbb{C}, \\ & \{(A, z, B, z) \in \text{Sp}(n-1) \times \text{Sp}(1) \times \text{Sp}(m-1) \times \text{Sp}(1)\} / \{\pm 1\} & \mathbb{K} = \mathbb{H}. \end{aligned}$$

The tangent space $T_{(e_n, e_m)}(S^{kn-1} \times S^{km-1})$ can be identified with

$$\mathbb{K}^{n-1} \oplus \mathfrak{S}(\mathbb{K}) \oplus \mathbb{K}^{m-1} \oplus \mathfrak{S}(\mathbb{K}),$$

and similarly, we have

$$(3.24) \quad T_{[e_n, e_m]}(S^{kn-1} \times_{S^{k-1}} S^{km-1}) = \mathbb{K}^{n-1} \oplus \mathbb{K}^{m-1} \oplus \mathfrak{S}(\mathbb{K}).$$

This identification is $K(n, m)$ -equivariant, where the action on the right hand side is as follows.

(1) If $\mathbb{K} = \mathbb{R}$, we have

$$T_{[e_n, e_m]}(S^{n-1} \times_{S^0} S^{m-1}) = \mathbb{R}^{n-1} \oplus \mathbb{R}^{m-1},$$

and the action of $[A, z, B, z] \in K(n, m)$ is by $L_A R_z$ in the first summand, and by $L_B R_z$ in the second one. The two summands are irreducible inequivalent representations of $O(n) \times O(m) \subset K(n, m)$.

(2) If $\mathbb{K} = \mathbb{C}$, we have

$$T_{[e_n, e_m]}(S^{2n-1} \times_{S^1} S^{2m-1}) = \mathbb{C}^{n-1} \oplus \mathbb{C}^{m-1} \oplus \mathfrak{S}(\mathbb{C}).$$

The action of $[A, z, B, z] \in K(n, m)$ is by $L_A R_{\bar{z}}$ in the first summand, by $L_B R_{\bar{z}}$ in the second one, and by the identity on the third. The action of the complex conjugation $\kappa \in K(n, m)$ is by complex conjugation in all three summands. Again, the decomposition (3.24) is a decomposition into irreducible, pairwise inequivalent $K(n, m)$ -representations.

(3) If $\mathbb{K} = \mathbb{H}$, we have

$$T_{[e_n, e_m]}(S^{4n-1} \times_{S^3} S^{4m-1}) = \mathbb{H}^{n-1} \oplus \mathbb{H}^{m-1} \oplus \mathfrak{S}(\mathbb{H}).$$

The action of $[A, z, B, z] \in K(n, m)$ is by $L_A R_{\bar{z}}$ in the first summand, by $L_B R_{\bar{z}}$ in the second summand, and by $\text{Ad}(z)$ in the third one. Again, this is a decomposition into pairwise inequivalent representations.

Since the summands in the decomposition (3.24) are irreducible and inequivalent $K(n, m)$ -representations, it follows that all invariant bilinear forms are given by direct sums of invariant forms on the three summands, and the forms on the summands are unique up to scalar multiplication. Hence the (real) dimension $d_{k,n,m}$ of

the space of $PG(n, m)$ -invariant bilinear forms on $M_{k,n,m}$ is equal to

$$d_{k,n,m} = \begin{cases} 0 & \mathbb{K} = \mathbb{R}, n = m = 1 \\ 1 & \mathbb{K} = \mathbb{R}, \text{ either } n = 1 \text{ or } m = 1 \\ 2 & \mathbb{K} = \mathbb{R}, n, m \geq 2 \\ 1 & \mathbb{K} = \mathbb{C}, \mathbb{H}, n = m = 1 \\ 2 & \mathbb{K} = \mathbb{C}, \mathbb{H} \text{ either } n = 1 \text{ or } m = 1 \\ 3 & \mathbb{K} = \mathbb{C}, \mathbb{H}, n, m \geq 2. \end{cases}$$

A parameter count shows that this is the same as the number of parameters in $sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$. \square

What remains to be proven is that

$$\{(s, u, t) \in (0, \infty)^3 \mid \text{scal}(sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}) > 0\} \subset \mathbb{R}^3$$

is contractible. To write down a formula for $\text{scal}(sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m})$, we fix some more notation. We define

$$(3.25) \quad c_n := \text{scal}(g_{S^{n-1}}) = (n-1)(n-2)$$

and

$$(3.26) \quad b_{k,n} := \text{scal}(g_{\mathbb{K}\mathbb{P}^{n-1}}) = \begin{cases} c_n & \mathbb{K} = \mathbb{R}, \\ 4n(n-1) & \mathbb{K} = \mathbb{C}, \\ 16(n^2-1) & \mathbb{K} = \mathbb{H} \end{cases}$$

(for the last one, see [16]). Furthermore, we let $a_{k,n}$ be the (constant) value of the quantity $|A|^2$ (see (3.12)) for the Riemannian submersion $(S^{kn-1}, g_{S^{kn-1}}) \rightarrow (\mathbb{K}\mathbb{P}^{n-1}, g_{\mathbb{K}\mathbb{P}^{n-1}})$. Using (3.12), we find

$$c_{kn} = \text{scal}(g_{S^{kn-1}}) = \text{scal}(g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{S^{k-1}}) = b_{k,n} + c_k - a_{k,n},$$

from which we may evaluate

$$(3.27) \quad a_{k,n} = \begin{cases} 0 & \mathbb{K} = \mathbb{R}, \\ 2(n-1) & \mathbb{K} = \mathbb{C}, \\ 12(n-1) & \mathbb{K} = \mathbb{H}. \end{cases}$$

Lemma 3.28. *With these notations, we have*

$$\text{scal}(g_{s,t}^{k,n}) = \frac{1}{s}b_{k,n} + \frac{1}{t}c_k - \frac{t}{s^2}a_{k,n}$$

and

$$\text{scal}(sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}) = \frac{1}{s}b_{k,n} + \frac{1}{u}b_{k,m} + \frac{1}{t}c_k - \frac{t}{u^2}a_{k,m} - \frac{t}{s^2}a_{k,n}.$$

Proof. The formula for the scalar curvature of the Berger metric is a straightforward consequence of (3.13). To get the second formula, let $a_{k,n,m,u,t}$ be the (constant) value of $|A|^2$ for the connection metric $g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$. This can be computed with the aid of the formula (3.10). Because the action vector field of the S^{k-1} -action on S^{km-1} lies in the vertical tangent bundle $T_v S^{km-1}$ which is scaled by the factor t , the quantity $a_{k,n,m,u,t}$ does not depend on u ; the dependence on t is expressed by the formula

$$a_{k,n,m,u,t} = ta_{k,n,m,1,1}.$$

Moreover, if S^{km-1} carries the round metric, the norm of the action vector field by an element in the Lie algebra of S^{k-1} agrees with the norm of the action vector field on S^{k-1} (the curve $t \mapsto p \exp(tX)$ is a closed geodesic with length independent of m). Therefore, $a_{k,n,m,1,1}$ does not depend on m and is given by $a_{k,n,m,1,1} = a_{k,n}$. This shows

$$a_{k,n,m,u,t} = ta_{k,n},$$

and we conclude

$$\text{scal}(g_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}) = b_{k,n} + \frac{1}{u}b_{k,m} + \frac{1}{t}c_k - \frac{t}{u^2}a_{k,m} - ta_{k,n}.$$

By global scaling with $\frac{1}{s}$, we obtain the formula for $\text{scal}(sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m})$. \square

Proof of Lemma 3.21. Each $PG(n, m)$ -invariant metric can be written as $sg_{\mathbb{K}\mathbb{P}^{n-1}} \oplus_{\omega_n} g_{u,t}^{k,m}$, by Lemma 3.23. By global scaling and Lemma 3.28, the space of invariant psc metrics is homotopy equivalent to the space of all $(s, u) \in (0, \infty)^2$ such that

$$(3.29) \quad \frac{1}{s}b_{k,n} + \frac{1}{u}b_{k,m} + c_k - \frac{1}{u^2}a_{k,m} - \frac{1}{s^2}a_{k,n} > 0.$$

In the exceptional cases, the left hand side of (3.29) is always 0. If $\mathbb{K} = \mathbb{R}$, we have $c_1 = a_{1,n} = 0$, and hence the left hand side of (3.29) is equal to $\frac{1}{s}b_{k,n} + \frac{1}{u}b_{k,m}$, which is always positive.

For the remaining cases, change coordinates to $x := \frac{1}{s}$ and $y := \frac{1}{u}$, and let

$$q(x, y) := a_{k,n}x^2 + a_{k,m}y^2,$$

$$c := c_k \geq 0$$

and

$$\ell(x, y) := b_{k,n}x + b_{k,m}y.$$

We have to look at the set of all $(x, y) \in (0, \infty)^2$ such that

$$(3.30) \quad q(x, y) - \ell(x, y) < c.$$

Now q is a positive semidefinite quadratic form and ℓ is a linear form. Hence $q - \ell$ is a convex function, and so the set of solutions to the inequality (3.30) is convex. It is also nonempty: if $\mathbb{K} = \mathbb{H}$, $c > 0$ and so (3.30) is satisfied for sufficiently small (x, y) . If $\mathbb{K} = \mathbb{C}$, $c = 0$, we find solutions of (3.30), provided that one of $b_{k,n}$, $b_{k,m}$ is positive, which is the case unless $n = m = 1$. \square

This concludes the proof of Proposition 3.19.

4. THE COMPUTATION IN THE SPIN CASE

4.1. The simply connected case. In this section, we finish the proof of Theorem A.

Proposition 4.1. *The map $\pi : BP(\text{Sp}(2) \times \text{Sp}(1)) \rightarrow BPS\text{p}(3)$ induces a surjective map*

$$\pi_* : \Omega_*^{\text{Spin}}(BP(\text{Sp}(2) \times \text{Sp}(1))) \rightarrow \Omega_*^{\text{Spin}}(BPS\text{p}(3)).$$

Let us postpone the proof for a moment.

Proof of Theorem A, assuming Proposition 4.1. We first have to recall some aspects of the proof of Stolz' theorem [37] (that a simply connected spin manifold of dimension $d \geq 5$ admits a psc metric if $\hat{a}(M) = 0$). The Atiyah–Bott–Shapiro orientation of spin vector bundles gives rise to a map

$$\hat{a} : \text{MSpin} \rightarrow \text{ko}$$

of ring spectra, from the spin cobordism spectrum to the connective real K -theory spectrum. On the level of homotopy groups, \hat{a} induces

$$\Omega_d^{\text{Spin}} \rightarrow \text{ko}_d$$

which assigns to a spin manifold the index of the spin Dirac operator, by the Atiyah–Singer index theorem. The Lichnerowicz formula implies that $\hat{a}(M) = 0$ is a necessary condition for the existence of a psc metric on M . Now let

$$T : \Omega_{d-8}^{\text{Spin}}(B\text{PSp}(3)) \rightarrow \Omega_d^{\text{Spin}}$$

be the map that assigns to a map $f : B \rightarrow \text{PSp}(3)$ the total space of the $\mathbb{H}\mathbb{P}^2$ -bundle classified by f . The composition $\hat{a} \circ T$ is zero, since total spaces of $\mathbb{H}\mathbb{P}^2$ -bundles admit psc metrics. The key result of [37] is that the sequence

$$(4.2) \quad \Omega_{d-8}^{\text{Spin}}(B\text{PSp}(3)) \xrightarrow{T} \Omega_d^{\text{Spin}} \xrightarrow{\hat{a}} \text{ko}_d$$

is exact (it is *not* part of a long exact sequence). To give credits properly, Stolz [37, Theorem 1.2] proved exactness of (4.2) after localization at the prime 2. After inverting 2, this was shown later by Kreck and Stolz [27, Proposition 3.3].

Now let M^d be a spin manifold which admits a psc metric, $d \geq 5$. Then $\hat{a}(M) = 0$, and so by the exactness of (4.2), $[M] \in \Omega_d^{\text{Spin}}$ contains the total space of an $\mathbb{H}\mathbb{P}^2$ -bundle with structure group $\text{PSp}(3)$. By Proposition 4.1, $[M]$ contains the total space of an $\mathbb{H}\mathbb{P}^2$ -bundle with structure group $P(\text{Sp}(2) \times \text{Sp}(1))$. By Theorems 3.1 and 2.6, it follows that $\mathcal{R}^+(M) \simeq \mathcal{R}^+(S^d)$, as claimed, if in addition M is simply connected. \square

Proof of Proposition 4.1. We shall prove separately that $\pi_* \otimes \mathbb{Z}_{(2)}$ and $\pi_* \otimes \mathbb{Z}[\frac{1}{2}]$ are surjective. We first turn to the 2-local case. Because

$$(4.3) \quad \mathbb{H}\mathbb{P}^2 = \text{PSp}(3)/P(\text{Sp}(2) \times \text{Sp}(1)),$$

π is homotopy equivalent to a smooth fibre bundle with fibre $\mathbb{H}\mathbb{P}^2$. In this situation, Becker and Gottlieb [5] constructed a transfer map

$$\text{trf}_\pi : \Sigma^\infty B\text{PSp}(3)_+ \rightarrow \Sigma^\infty BP(\text{Sp}(2) \times \text{Sp}(1))_+$$

on the level of suspension spectra, and they showed that the composition $\pi \circ \text{trf}_\pi : \Sigma^\infty B\text{PSp}(3)_+ \rightarrow \Sigma^\infty B\text{PSp}(3)_+$ induces multiplication by the Euler number $\chi(\mathbb{H}\mathbb{P}^2) = 3 \in \mathbb{Z}_{(2)}^\times$ on (ordinary) homology. If A_* is a generalized homology theory whose coefficient groups are $\mathbb{Z}_{(2)}$ -modules, a straightforward argument with the Atiyah–Hirzebruch spectral sequence proves from this fact that $\pi_* : A_*(BP(\text{Sp}(2) \times \text{Sp}(1))) \rightarrow A_*(B\text{PSp}(3))$ is (split) surjective. Apply that statement to $A_* := \Omega_*^{\text{Spin}}(-) \otimes \mathbb{Z}_{(2)}$. There was nothing special about spin cobordism in this argument.

For the consideration after inverting 2, some more ingredients are needed. Firstly, the map $B\text{Sp}(3) \rightarrow B\text{PSp}(3)$ induces an isomorphism in homology with coefficients in $\mathbb{Z}[\frac{1}{2}]$, and hence also in spin bordism tensored with $\mathbb{Z}[\frac{1}{2}]$. The same argument

applies to $B(\mathrm{Sp}(2) \times \mathrm{Sp}(1)) \rightarrow BP(\mathrm{Sp}(2) \times \mathrm{Sp}(1))$. Therefore, we only need to prove that

$$(4.4) \quad \Omega_*^{\mathrm{Spin}}(B(\mathrm{Sp}(2) \times \mathrm{Sp}(1))) \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow \Omega_*^{\mathrm{Spin}}(B\mathrm{Sp}(3)) \otimes \mathbb{Z}[\frac{1}{2}]$$

is surjective. It is well-known that the integral homology of $B\mathrm{Sp}(n)$ is torsionfree and concentrated in degrees divisible by 4, see e.g. [31, Theorem III.5.6]. Moreover, the map $B(\mathrm{Sp}(2) \times \mathrm{Sp}(1)) \rightarrow B\mathrm{Sp}(3)$ induces an epimorphism in homology with arbitrary coefficients. This is because the homotopy fibre is $\mathbb{H}\mathbb{P}^2$; hence the Leray–Serre spectral sequence collapses (for degree reasons) at the E^2 -page.

We also need to know a fact about spin cobordism from Anderson, Brown and Peterson’s work [2]. Namely, Theorem 2.2 of that paper implies that $\Omega_*^{\mathrm{Spin}} \otimes \mathbb{Z}[\frac{1}{2}]$ is concentrated in degrees divisible by 4 (this can also be deduced from the structure of Ω_*^{SO} described in Theorem 6.4 below, together with the fact that the forgetful map $\Omega_*^{\mathrm{Spin}} \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow \Omega_*^{\mathrm{SO}} \otimes \mathbb{Z}[\frac{1}{2}]$ is an isomorphism, which follows from Serre class theory). Therefore, the Atiyah–Hirzebruch spectral sequence

$$E_{p,q}^2 = H_p(B\mathrm{Sp}(3); \Omega_q^{\mathrm{Spin}} \otimes \mathbb{Z}[\frac{1}{2}]) \Rightarrow \Omega_{p+q}^{\mathrm{Spin}}(B\mathrm{Sp}(3)) \otimes \mathbb{Z}[\frac{1}{2}]$$

and its analogue $\bar{E}_{p,q}^2$ for $B(\mathrm{Sp}(2) \times \mathrm{Sp}(1))$ collapse at the E^2 -page. We have argued that the comparison map $\bar{E}_{p,q}^2 \rightarrow E_{p,q}^2$ is surjective, and since the spectral sequences collapse, surjectivity on the E^∞ -page follows. Finally, an argument with the 5-lemma proves that (4.4) is surjective, as claimed. \square

4.2. Generalization to some nontrivial fundamental groups. If one tries to carry over the above argument for Theorem A to the case of spin manifolds with a given fundamental group Γ , one runs into formidable difficulties. These difficulties are very similar to that which are met when studying the Gromov–Lawson–Rosenberg conjecture, which has counterexamples for rather innocent groups [34]. But at least in some interesting cases, our argument goes through. We have not attempted a more systematic study, though.

Theorem 4.5. *Let Γ be a finitely presented torsion-free group, and assume the following:*

- (1) *the classifying space $B\Gamma$ splits stably as a wedge of sphere spectra, in other words, $\Sigma^\infty B\Gamma_+ \simeq \bigvee_{i \in I} \Sigma^{n_i} \mathbb{S}$,*
- (2) *Γ satisfies the real strong Novikov conjecture, in other words, the assembly map $\mu : \mathrm{KO}_*(B\Gamma) \rightarrow \mathrm{KO}_*(C_r^*(\Gamma))$ is injective.*

Then if M_0, M_1 are two closed spin manifolds of dimension $d \geq 5$ with fundamental group Γ and $\mathcal{R}^+(M_i) \neq \emptyset$, we have

$$\mathcal{R}^+(M_0) \simeq \mathcal{R}^+(M_1).$$

- Examples 4.6.*
- (1) The free abelian group \mathbb{Z}^n satisfies all requirements. The stable splitting condition is clear, and free abelian groups even satisfy the Baum–Connes conjecture (in the complex case), which implies the real strong Novikov conjecture by [35].
 - (2) Similarly, the free group F_n satisfies the hypotheses of the Theorem (free groups satisfy the Baum–Connes conjecture by [21] because they have the Haagerup property).

Proof. We have to prove that the cobordism class $[M_i] \in \Omega_d^{\text{Spin}}(B\Gamma)$ contains a representative which is an $\mathbb{H}\mathbb{P}^2$ -bundle with structure group $H := P(\text{Sp}(2) \times \text{Sp}(1))$.

The hypothesis that $\Sigma^\infty B\Gamma_+ \simeq \bigvee_{i \in I} \Sigma^{n_i} \mathbb{S}$ has the effect that

$$A_k(B\Gamma) = \bigoplus_{i \in I} A_{k-n_i}(*).$$

for each homology theory A_* , and this sum decomposition is natural in A_* . Kreck–Stolz' sequence (4.2), together with Proposition 4.1 and the (obvious) injectivity of $\text{ko}_* \rightarrow \text{KO}_*$ yields an exact sequence

$$\Omega_{*-8}^{\text{Spin}}(BP(\text{Sp}(2) \times \text{Sp}(1))) \rightarrow \Omega_*^{\text{Spin}} \rightarrow \text{KO}_*$$

whose maps are induced by map of spectra. The induced sequence

$$\Omega_{*-8}^{\text{Spin}}(BP(\text{Sp}(2) \times \text{Sp}(1)) \times B\Gamma) \rightarrow \Omega_*^{\text{Spin}}(B\Gamma) \rightarrow \text{KO}_*(B\Gamma)$$

is therefore exact, by the above observation. Assuming that Γ satisfies the real strong Novikov conjecture gives an exact sequence

$$(4.7) \quad \Omega_{*-8}^{\text{Spin}}(BP(\text{Sp}(2) \times \text{Sp}(1)) \times B\Gamma) \xrightarrow{T} \Omega_*^{\text{Spin}}(B\Gamma) \xrightarrow{\text{ind}_\Gamma} \text{KO}_*(C_r^*(\Gamma)).$$

The map T again takes the total space of the classified $\mathbb{H}\mathbb{P}^2$ -bundle, and ind_Γ associates to a cobordism class (M, f) the index of the Rosenberg–Dirac operator. The Lichnerowicz vanishing argument carries over to this case and shows that if M admits a psc metric, then $\text{ind}_\Gamma(M, f) = 0$. Using the exactness of (4.7), we conclude that (M, f) is cobordant to the total space of an $\mathbb{H}\mathbb{P}^2$ -bundle with structure group $P(\text{Sp}(2) \times \text{Sp}(1))$. \square

5. TWO FURTHER GEOMETRIC INGREDIENTS FOR THE NON-SPIN CASE

For the proof of Theorem A that we just gave, Theorem 3.1 provided enough examples for admissible splittings. For the proof of Theorem B, these do not quite suffice, and we need some more geometric ingredients, which will be developed in this section. The main results are Propositions 5.1, 5.2 and 5.8.

5.1. Stability of product metrics. The goal of this subsection is the proof of the following two propositions.

Proposition 5.1. *Let $x \in \Omega_*^{\text{SO}}$ be a cobordism class which can be written as $x = yz$, with y of dimension at least 6 and z of positive dimension. Then x contains a representative with an admissible splitting.*

Proposition 5.2. *Let $V_i \rightarrow N_i$ be vector bundles of rank $r_i \geq 3$, let $\psi : S(V_0) \cong S(V_1)$ be a diffeomorphism and let $M := D(V_0) \cup_\psi D(V_1)^{\text{op}}$. Let P be a closed manifold which admits a psc metric. Then $M \times P$ has an admissible splitting.*

An important step in the proof of Proposition 5.1 is the following result.

Lemma 5.3. *Let $W : M_0 \rightsquigarrow M_1$ be a cobordism such that $\dim(W) = d \geq 6$ and such that both inclusions $M_i \rightarrow W$ are 2-connected. Let (N, g_N) be a closed psc manifold, and let g_W be a Riemannian metric on W which has product form near the boundary, so that the product metric $g_W \oplus g_N$ on $W \times N$ has positive scalar curvature. Then $g_W \oplus g_N$ is stable.*

The proof of Lemma 5.3 begins with two lemmas.

Lemma 5.4. *Let W be a compact manifold with boundary. Then for each $\epsilon > 0$, the space*

$$\mathcal{R}^{>-\epsilon}(W) := \{h \in \mathcal{R}(W) \mid \inf \text{scal}(h) > -\epsilon\}$$

is (weakly) contractible. If (N, g_N) is a closed psc manifold, the space of product psc metrics (that is, psc metrics of the form $g_W \oplus g_N$) on $W \times N$ is contractible.

Proof. The space $\mathcal{R}^{>-\epsilon}(W)$ is nonempty: if $h \in \mathcal{R}(W)$, then $th \in \mathcal{R}^{>-\epsilon}(W)$ if t is sufficiently large. Pick $h_0 \in \mathcal{R}^{>-\epsilon}(W)$. Let X be a compact space and let $h : X \rightarrow \mathcal{R}^{>-\epsilon}(W)$ be a map. Let

$$H : K \times I \rightarrow \mathcal{R}(W), \quad H(x, t) := (1-t)h(x) + th_0,$$

and

$$\lambda := \inf_{(x,t) \in K \times [0,1]} \text{scal}(H(x, t)).$$

Choose $a \geq 1$ such that $\frac{1}{a}\lambda > -\epsilon$, and define a homotopy $P : K \times [0, 3] \rightarrow \mathcal{R}^{>-\epsilon}(W)$ by

$$P(x, t) := \begin{cases} (1-t+ta)h(x) & t \in [0, 1] \\ (2-t)ah(x) + (t-1)ah_0 & t \in [1, 2] \\ ((3-t)a + (t-2))h_0 & t \in [2, 3]. \end{cases}$$

Hence $\mathcal{R}^{>-\epsilon}(W) \simeq *$. The second claim of the Lemma follows from the first and the observation that

$$\text{scal}(g_W \oplus g_N) > 0 \Leftrightarrow \inf \text{scal}(g_W) > -\inf \text{scal}(g_N).$$

□

Lemma 5.5. *Let N be a closed manifold which admits a psc metric, let $W^d : M_0 \rightsquigarrow M_1$ be a cobordism, and assume that each product psc metric on $W \times N$ is right stable. Let $\varphi : S^{k-1} \times D^{d-k+1} \rightarrow W$ be a surgery datum in the interior of W with $3 \leq k \leq d-2$, and let W_φ be the result of performing a surgery along φ . Then each product psc metric on $W_\varphi \times N$ is right stable. The same is true for left stable metrics.*

Proof. Let $\varphi' : S^{d-k} \times D^k \rightarrow W_\varphi$ be the dual surgery datum. We define

$$\phi := \varphi \times \text{Id}_N : S^{k-1} \times D^{d-k+1} \times N \rightarrow W \times N$$

and

$$\phi' := \varphi' \times \text{Id}_N : S^{d-k} \times D^k \rightarrow W_\varphi \times N.$$

Then

$$W_\varphi \times N = ((W \times N) \setminus \text{Im}(\phi)) \cup_{S^{k-1} \times S^{d-k} \times N} (D^k \times S^{d-k} \times N).$$

Fix a psc metric g_N on N and let $\mathcal{R}^+(W \times N, \phi) \subset \mathcal{R}^+(W \times N)$ be the subspace of those metrics which are equal to $g_{S^{k-1}} \oplus g_{\text{tor}}^{d-k} \oplus g_N$ on $\text{Im}(\phi)$, and define $\mathcal{R}^+(W_\varphi \times N, \phi') \subset \mathcal{R}^+(W_\varphi \times N)$ in an analogous way. The inclusions

$$\mathcal{R}^+(W \times N, \phi) \rightarrow \mathcal{R}^+(W \times N), \quad \mathcal{R}^+(W_\varphi \times N, \phi') \rightarrow \mathcal{R}^+(W_\varphi \times N)$$

are weak equivalences by Theorem 2.1 (or rather a variant of that for manifolds with boundary which is covered by the proof in [12]). There is a homeomorphism

$$(5.6) \quad \mathcal{R}^+(W \times N, \phi) \cong \mathcal{R}^+(W_\varphi \times N, \phi'),$$

and hence a weak equivalence

$$S : \mathcal{R}^+(W \times N) \simeq \mathcal{R}^+(W_\varphi \times N).$$

By the argument given in the proof of Theorem 2.6 of [7], the path components which contain right stable metric correspond to each other under the bijection $\pi_0(S)$. Furthermore, the homeomorphism (5.6) and its inverse preserve product metrics. By Lemma 5.4, there are *unique* components of $\mathcal{R}^+(W \times N)$ and $\mathcal{R}^+(W_\varphi \times N)$ which contain metrics of the form $g_W \oplus g_N$, and so these two components correspond to each other under $\pi_0(S)$. Hence one is right stable if and only if the other one is. \square

Proof of Lemma 5.3. This is similar to the proof of [13, Theorem E = Theorem 3.1.2(ii)], and we refer to that paper for more details. First note that any psc metric of the form $g_M \oplus dt^2 \oplus g_N$ on $M \times [0, 1] \times N$ is stable. Now let W be as in the statement of the proposition. By the proof of the s -cobordism theorem [24] or [42], W has a handlebody decomposition (relative to M_0) in which all handles have index in $\{3, \dots, d-2\}$. As in explained in the proof of [13, Theorem 3.1.2(ii)], it follows that $W \cup W^{op}$ and $M_0 \times [0, 2]$ are related by a sequence of surgeries of index in $\{3, \dots, d-2\}$. By Lemma 5.5, each product psc metric on $(W \cup W^{op}) \times N$ is stable. Now let $h \in \mathcal{R}^+(W \times N)$ be a product psc metric. Then $h \cup h^{op}$ is a product psc metric, therefore stable, thus the gluing map $\mu(-, h)$ has a left inverse in the homotopy category, namely $\mu(-, h^{op})$.

The same argument can be applied with the roles of W and W^{op} reversed, and leads to the conclusion that $\mu(-, h^{op})$ also has a left inverse in the homotopy category. Therefore $\mu(-, h^{op})$ is a weak equivalence, and so is $\mu(-, h)$. This means that h is left stable. By Theorem E of [13], h is also right stable. \square

Proof of Proposition 5.1. Let M^d be a representative for y which is relatively 3-connected. In particular, M is simply connected, $H_2(M; \mathbb{Z}) \cong H_2(BSO; \mathbb{Z}) \cong \mathbb{Z}/2$, and M does not have a spin structure.

Let $S^2 \rightarrow M$ be an embedding which represents a generator of $\pi_2(M) \cong \mathbb{Z}/2$. The normal bundle is the nontrivial vector bundle over S^2 , and so we obtain a 2-connected embedding $P \rightarrow M$ of the disc bundle of the nontrivial vector bundle over S^2 , which leads to a decomposition $M = P \cup_{\partial P} (M \setminus P)$. By general position, the inclusion map $M \setminus P \rightarrow M$ is $(d-3) \geq 3$ -connected, and so the manifold $M \setminus P$ is relatively 3-connected.

The argument so far only used that M was relatively 3-connected and not that it was closed. Hence we can apply the same argument to $M \setminus P$. The result is a decomposition

$$M = P \cup W \cup P^{op}$$

where $W : \partial P \rightsquigarrow \partial P$ is a cobordism such that both inclusions $\partial P \rightarrow W$ are 2-connected. Let g_P be a connection torpedo metric on P , and choose $g_W \in \mathcal{R}(W)_{g_{\partial P}, g_{\partial P}}$, not necessarily of positive scalar curvature.

The cobordism class $z \in \Omega_*^{\text{SO}}$ contains a representative N which carries a psc metric g_N . This follows from the proof of [18, Corollary C] (a more elegant way to see this is to invoke the main result of [15]). We can assume that $\text{scal}(g_W \oplus g_N) > 0$.

By Lemma 5.3, the metric $g_W \oplus g_N$ is stable, and the metric $g_P \oplus g_N$ is a connection torpedo metric, so it is right stable by Theorem 3.15. Likewise, $g_P^{op} \oplus g_N$

is left stable, and we have found an admissible splitting

$$M \times N = (P \times N) \cup_{\partial P \times N} ((W \cup P^{op}) \times N)$$

with suitable psc metric $(g_P \oplus g_N) \cup ((g_W \cup g_P^{op}) \oplus g_N)$. \square

For the proof of Proposition 5.2, we need another lemma.

Lemma 5.7. *Let (M, g_M) and (N, g_N) be two psc manifolds. Then the product map $\nu : \mathcal{R}^+(M) \times \mathcal{R}^+(N) \rightarrow \mathcal{R}^+(M \times N)$, $(g_0, g_1) \mapsto g_0 \oplus g_1$ is homotopic to the constant map.*

Proof. We only show that the map is weakly homotopic, i.e. that the restriction to a compact subset $K \subset \mathcal{R}^+(M) \times \mathcal{R}^+(N)$ is nullhomotopic. The technique to write down the homotopies is very similar to the proof of Lemma 5.4, hence we shall be very brief. The first step is a homotopy from $\nu|_K$ to the map $(g_0, g_1) \mapsto g_M \oplus g_1$, which is very similar to the homotopy used in the proof of Lemma 5.4. The second step is a homotopy from that map to the constant map $g_M \oplus g_N$. \square

Proof of Proposition 5.2. Pick Riemannian metrics h_i on N_i such that $h_i \oplus g_P$ is a psc metric on $N_i \times P$. Pick metric connections ω_i on V_i and suitable torpedo metrics $g_{\text{tor}}^{r_i}$ on \mathbb{R}^{r_i} such that $h_i \oplus_{\omega_i} g_{\text{tor}}^{r_i}$ is a psc metric on $D(V_i)$.

The product metrics $(h_i \oplus_{\omega_i} g_{\text{tor}}^{r_i}) \oplus g_P$ are connection torpedo metrics. Lemma 5.7 implies that the metrics $(h_0 \oplus_{\omega_0} g_{\text{tor}}^{r_0})|_{S(V_0)} \oplus g_P$ and $(\psi^*(h_1 \oplus_{\omega_1} g_{\text{tor}}^{r_1})|_{S(V_1)}) \oplus g_P$ are isotopic; this finishes the proof because of Proposition 3.16. \square

5.2. Splitting a certain 5-manifold. Here, we prove the following.

Proposition 5.8. *There is a closed oriented 5-manifold M with the following properties:*

- (1) *We can write $M = W_0 \cup W_1$ as a union of two disc bundles of rank 3 over S^2 ,*
- (2) *this is an admissible splitting, and*
- (3) *$[M] \neq 0 \in \Omega_5^{\text{SO}}$.*

Construction 5.9. Let

$$V := S^3 \times_{S^1} (\mathbb{C} \oplus \mathbb{R}) \rightarrow \mathbb{C}\mathbb{P}^1,$$

where S^1 acts on \mathbb{C} by rotations and trivially on \mathbb{R} . This is the unique nontrivial real vector bundle of rank 3 on $\mathbb{C}\mathbb{P}^1$, and V does not have a spin structure. We write $W := D(V)$ and $N := S(V)$ for the unit disc bundle and unit sphere bundle. Note that

$$N = S^3 \times_{S^1} S^2$$

is the *Hirzebruch surface*; in [22], it is shown that $N \cong \mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$. Let us write

$$(5.10) \quad N = (S^3 \times_{S^1} D^2) \cup (S^3 \times [0, 1]) \cup_{\varphi} (S^3 \times_{S^1} D^2),$$

using the diffeomorphism $\varphi : S^3 \times_{S^1} S^1 \rightarrow S^3$ given by $\varphi([u, x]) := ux$. We now define a diffeomorphism $F : N \rightarrow N$ as follows. On the first copy of $S^3 \times_{S^1} D^2$, F is the identity. On the second copy of $S^3 \times_{S^1} D^2$, F is given by complex conjugation $[u, x] \mapsto [\bar{u}, \bar{x}]$. Via the gluing diffeomorphisms, these maps induce the identity on $S^3 \times \{0\}$ and the complex conjugation κ on $S^3 \times \{1\}$. The complex conjugation $S^3 \subset \mathbb{C}^2$ lies in $\text{SO}(4)$, and hence there is a path $\gamma : [0, 1] \rightarrow \text{SO}(4)$ with $\gamma(0) = 1$ and $\gamma(1) = \kappa$. We furthermore assume that γ is constant near 0 and constant

near 1. On the middle $S^3 \times [0, 1]$, we define F by the formula $F(t, u) := (t, \gamma(t)u)$. Altogether, this defines $F : N \rightarrow N$. We now define

$$M := W \cup_F W,$$

which is a closed 5-manifold which by construction is the union of two disc bundles of rank 3, as claimed in Proposition 5.8. Since W is simply connected and N is connected, M is simply connected, by Seifert–Van-Kampen. Therefore, M is orientable, and we chose an orientation on M .

Proof that M is not nullbordant. We will show that

$$(5.11) \quad H_k(M; \mathbb{Z}) = \begin{cases} \mathbb{Z} & k = 0, 5, \\ \mathbb{Z}/2 & k = 2, \\ 0 & k \neq 0, 2, 5. \end{cases}$$

This will imply the lemma, by the following argument. It is enough to prove that $\langle w_2(TM)w_3(TM), [M] \rangle \neq 0$. To that end, recall that the *semi-characteristic* $\kappa(X; \mathbb{F}) \in \mathbb{Z}/2$ of a closed $(2m+1)$ -manifold X with respect to a field \mathbb{F} is defined as

$$\kappa(X; \mathbb{F}) := \sum_{k=0}^m \dim_{\mathbb{F}}(H_k(X; \mathbb{F})) \in \mathbb{Z}/2.$$

Lusztig, Milnor and Peterson showed in [28] that

$$\kappa(X; \mathbb{Q}) - \kappa(X; \mathbb{F}_2) = \langle w_2(TX)w_{2m-1}(TX), [X] \rangle \in \mathbb{Z}/2.$$

Together with (5.11), this shows that $\langle w_2(TM)w_3(TM), [M] \rangle \neq 0$.

It remains to compute $H_2(M; \mathbb{Z})$; the other groups are determined by the simple connectivity of M and by Poincaré duality. The S^1 -action on $S^2 \subset \mathbb{C} \oplus \mathbb{R}$ has the two fixed points $(0, \pm 1)$. Those two fixed points determine sections s_{\pm} of the bundle $N = S^3 \times_{S^1} S^2 \rightarrow \mathbb{C}\mathbb{P}^1$ and hence elements $b_{\pm} = \text{hur}(s_{\pm}) \in H_2(N; \mathbb{Z})$. It is easily verified (e.g. by the Mayer-Vietoris sequence for the decomposition (5.10)) that (b_+, b_-) is a \mathbb{Z} -basis of $H_2(N) \cong \mathbb{Z}^2$.

The two S^1 -fixed points $(0, \pm 1)$ can be connected by a path of fixed points in D^3 ; therefore, the two sections s_{\pm} becomes homotopic in $S^3 \times_{S^1} D^3 = W$. Hence $j_*(b_-) = j_*(b_+) \in H^2(W)$, where $j : N \rightarrow W$ denotes the inclusion map. Since $N \rightarrow W$ is 2-connected, j_* is surjective on H_2 . Hence $j_*(b_{\pm})$ must be a generator $u \in H_2(W) \cong \mathbb{Z}$.

To compute the effect of the diffeomorphism F on homology, let us agree that the signs are chosen in such a way that s_- lies in the first part of the decomposition (5.10) and s_+ in the third one. With these choices

$$F_*(b_-) = b_-; \quad F_*(b_+) = -b_+.$$

The first is clear. For the second, consider the commutative diagram

$$\begin{array}{ccc} S^3 & \xrightarrow{\kappa} & S^3 \\ \downarrow \eta & & \downarrow \eta \\ \mathbb{C}\mathbb{P}^1 & \xrightarrow{q} & \mathbb{C}\mathbb{P}^1 \end{array}$$

(q and κ are the complex conjugation maps and η the Hopf map). Using that $\text{deg}(q) = -1$, we get that the complex conjugation induces -1 on $H_2(S^3 \times_{S^1} 0) \cong$

$H_2(S^3 \times_{S^1} D^2)$. To compute $H_2(M)$, consider the piece of the Mayer-Vietoris sequence for the decomposition $M = W \cup_F W$:

$$H_2(N; \mathbb{Z}) \xrightarrow{(j_*, j_* \circ F_*)} H_2(W; \mathbb{Z}) \oplus H_2(W; \mathbb{Z}) \rightarrow H_2(M; \mathbb{Z}) \rightarrow 0$$

With respect to the bases (b_+, b_-) and (u, u) , the first map is represented by the matrix

$$\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$$

with determinant 2. It follows that $H_2(M) \cong \mathbb{Z}/2$, as claimed. \square

Remark 5.12. It follows from Barden's classification theorem for simply connected 5-manifolds [4] that the manifold M is diffeomorphic to $SU(3)/SO(3)$, the Wu manifold.

It remains to be proven that $M = W \cup_F W$ is an admissible splitting. This is accomplished by Lemmas 5.13 and 5.14 below. To prove them, we have to do some geometry.

Lemma 5.13. *Let ω be the $U(2)$ -invariant connection on the Hopf bundle $\eta : S^3 \rightarrow \mathbb{C}P^1$ which was introduced in §3.4, and let g_{tor}^3 be a torpedo metric on D^3 of radius 1. Then the metric*

$$g_{\mathbb{C}P^1} \oplus_{\omega} g_{\text{tor}}^3$$

on $S^3 \times_{S^1} D^3 = W$ has positive scalar curvature and is right stable.

Proof. By Theorem 3.15, it is enough to verify that $\text{scal}(g_{\mathbb{C}P^1} \oplus_{\omega} g_{\text{tor}}^3) > 0$. Using (3.12) and (3.26), we get

$$\text{scal}(g_{\mathbb{C}P^1} \oplus_{\omega} g_{\text{tor}}^3) = 8 + \text{scal}(g_{\text{tor}}^3) - |A_{g_{\text{tor}}^3}|^2.$$

By definition of a torpedo metric, $\text{scal}(g_{\text{tor}}^3) \geq \text{scal}(g_{S^2}) = 2$. The closed orbits of the S^1 -action on D^3 , in the torpedo metric g_{tor}^3 , have length at most 2π and are therefore at most as long as the orbits of the translation action of S^1 on itself. It follows, using (3.10), that

$$|A_{g_{\text{tor}}^3}|^2 \leq |A|^2 = 2.$$

where $|A|^2$ is the norm of the A -tensor for the translation action, which was computed in (3.27). \square

Lemma 5.14. *The psc metrics $g_{\mathbb{C}P^1} \oplus_{\omega} g_{S^2}$ and $F^*(g_{\mathbb{C}P^1} \oplus_{\omega} g_{S^2})$ lie in the same path component of $\mathcal{R}^+(N)$.*

For the proof, we need another auxiliary result:

Lemma 5.15. *Let (N, g) be a closed Riemannian manifold, $\text{scal}(g) > 0$, and let $\gamma : [0, 1] \rightarrow \text{Isom}(N, g)$ be a smooth path which is constant near 0 and 1. Let $\Gamma : N \times [0, 1] \rightarrow N \times [0, 1]$ be the diffeomorphism $(x, t) \mapsto (\gamma(t)x, t)$. Then the two metrics*

$$g \oplus dt^2, \Gamma^*(g \oplus dt^2) \in \mathcal{R}^+(N \times [0, 1])_{g, g}$$

lie in the same path component.

Proof. Extend γ to a smooth function $\gamma : \mathbb{R} \rightarrow \text{Isom}(N, g)$, constant on $(-\infty, 0]$ and on $[1, \infty)$. For $r > 0$, let $\gamma_r(t) := \gamma(\frac{1}{r}t)$ and define $\Gamma_r : N \times \mathbb{R} \rightarrow N \times \mathbb{R}$ accordingly. It will be enough to prove that for sufficiently large r , the metric

$$(1-s)g \oplus dt^2 + s\Gamma_r^*(g \oplus dt^2)$$

on $N \times \mathbb{R}$ has positive scalar curvature for all $s \in [0, 1]$. Let X_r be the vector field on $N \times \mathbb{R}$ given by

$$X_r(p, t) := \left. \frac{d}{du} \right|_{u=0} (\gamma_r(t+u)p, t).$$

Note that

$$X_r(p, t) = \frac{1}{r} X_1(p, \frac{t}{r}).$$

Using that $\gamma(t)$ is an isometry, one calculates that the metric $\Gamma^*(g \oplus dt^2)$ is given (using the splitting $T_{(p,t)}(N \times \mathbb{R}) = T_p N \oplus \mathbb{R}$) by the matrix

$$\begin{pmatrix} g & X_r \\ X_r^T & 1 + \|X_r\|^2 \end{pmatrix}.$$

For $r \rightarrow \infty$, this converges uniformly to $g \oplus dt^2$, together with all derivatives. From that, the claim follows. \square

Proof of Lemma 5.14. Let g_{tor}^2 be a torpedo metric on D^2 of radius 1. This does not have positive scalar curvature, but at least we know that $\text{scal}(g_{\text{tor}}^2) \geq 0$. Let $g_{\text{d\text{tor}}}^2$ be the metric on $D^2 \cup (S^1 \times [0, 1]) \cup D^2 = S^2$ which is given by g_{tor}^2 on the two discs and by $g_{S^1} \oplus dt^2$ on the cylinder. There is a smooth path $u \mapsto g(u)$ of Riemannian metrics on S^2 with $g(0) = g_{S^2}$, $g(1) = g_{\text{d\text{tor}}}^2$, such that each $g(u)$ is S^1 -invariant, has nonnegative scalar curvature and such that the lengths of the S^1 -orbits for each $g(u)$ are at most 2π . This can be seen from the formula for the scalar curvature of a warped product metric on $S^1 \times \mathbb{R}$ which can be found in e.g. [44, §2.2].

Consider the metric $g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g(u)$ on $S^3 \times_{S^1} S^2 = N$. It follows that

$$\text{scal}(g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g(u)) = 8 + \text{scal}(g(u)) - |A_{g(u)}|^2 \geq 8 - |A|^2 = 6$$

for all u . Hence $g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{S^2}$ and $g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2$ lie in the same path component of $\mathcal{R}^+(N)$, and therefore, it suffices for our aim to prove that

$$[g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2] = [F^*(g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2)] \in \pi_0(\mathcal{R}^+(N)).$$

By the construction of F (and because the torpedo metric is $O(2)$ -invariant), $F^*(g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2)$ and $g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2$ agree on the two copies of $S^3 \times_{S^1} D^2$ inside N . On the middle piece $S^3 \times [0, 1]$, we have

$$F^*(g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega g_{\text{d\text{tor}}}^2) = F^*(g_{\mathbb{C}\mathbb{P}^1} \oplus_\omega (g_{S^1} \oplus dx^2)) = F^*(g_{S^3} \oplus dx^2) \in \mathcal{R}^+(S^3 \times [0, 1])_{g_{S^3}, g_{S^3}}.$$

By Lemma 5.15, it is isotopic within $\mathcal{R}^+(S^3 \times [0, 1])_{g_{S^3}, g_{S^3}}$ to the product metric. This finishes the proof. \square

6. THE COMPUTATION IN THE NON-SPIN CASE

The exact sequence (4.2) has an analogue for the oriented cobordism, namely the exact sequence

$$\Omega_{d-4}^{\text{SO}}(B(C_2 \times PU(3))) \xrightarrow{T} \Omega_d^{\text{SO}} \rightarrow \pi_d(H\mathbb{Z}),$$

where T takes the total space of $\mathbb{C}\mathbb{P}^2$ -bundles and the second map is given by the Thom class, see [15]. Unfortunately, this is of no use for the proof of Theorem B, since $\mathbb{C}\mathbb{P}^2$ has too small dimension. Therefore, we have to take a more pedestrian route and need rather explicit generators for the oriented bordism ring Ω_*^{SO} . The following summarizes the information which is relevant for us; we show in §6.2 how to extract this from the classical literature.

Theorem 6.1. *The oriented cobordism ring Ω_*^{SO} is generated (multiplicatively) by elements of the following type.*

- (1) Classes $v_{4k} \in \Omega_{4k}^{\text{SO}}$, one for each $k \geq 2$ and each v_{4k} is a sum of classes which are represented by $\mathbb{C}\mathbb{P}^n$ -bundles with structure group $U(n+1)$ and $n \geq 3$,
- (2) $v_4 := [\mathbb{C}\mathbb{P}^2]$,
- (3) an infinite set Z of classes in degrees ≥ 9 which are represented by $\mathbb{C}\mathbb{P}^n$ -bundles with structure group C_2 , $n \geq 3$,
- (4) a single element z_5 in degree 5.

The low-dimensional cobordism groups are as follows.

- (1) $\Omega_k^{\text{SO}} = 0$ for $k = 1, 2, 3, 6, 7$,
- (2) $\Omega_4^{\text{SO}} = \mathbb{Z}$, generated by $v_4 = [\mathbb{C}\mathbb{P}^2]$,
- (3) $\Omega_5^{\text{SO}} \cong \mathbb{Z}/2$, generated by z_5
- (4) $\Omega_8^{\text{SO}} \cong \mathbb{Z}^2$, generated by $v_8 = [\mathbb{C}\mathbb{P}^4]$ and $v_4^2 = [\mathbb{C}\mathbb{P}^2]^2$, and the subgroup generated by $[\mathbb{C}\mathbb{P}^4]$ and $[\mathbb{H}\mathbb{P}^2]$ has index 3.

6.1. Proof of Theorem B. Let us first show how to use it to finish the proof of Theorem B. Besides Theorem 6.1, we need one more computation from homotopy theory.

Proposition 6.2. *The map*

$$(6.3) \quad \Omega_*^{\text{SO}}(BU(1)^n) \rightarrow \Omega_*^{\text{SO}}(BU(n)),$$

induced by the inclusion of a maximal torus $U(1)^n \rightarrow U(n)$, is surjective. Hence each $\mathbb{C}\mathbb{P}^{n-1}$ -bundle $E \rightarrow M$ over a closed oriented manifold with structure group $U(n)$ is oriented cobordant to a $\mathbb{C}\mathbb{P}^{n-1}$ -bundle with structure group $U(1)^n$.

Proof. The ring spectrum MSO is complex oriented; in fact the forgetful map $\text{MU} \rightarrow \text{MSO}$ defines a complex orientation, using Lemma II.4.6. of [1]. By [25, Proposition 4.3.3] or [1, Lemma II.4.1], surjectivity of (6.3) follows. \square

Proof of Theorem B. Let $A_d \subset \Omega_d^{\text{SO}}$ be the subgroup of cobordism classes which contain a representative that has an admissible splitting. We first consider the case $d \neq 8$, $d \geq 5$, and have to prove that $A_d = \Omega_d^{\text{SO}}$.

Let $k \geq 2$ and let v_{4k} be the generator described in Theorem 6.1. For an arbitrary $x \in \Omega_*^{\text{SO}}$, the element xv_{4k} can be represented by a $\mathbb{C}\mathbb{P}^{n-1}$ -bundle with structure group $U(2) \times U(n-2)$, and $n \geq 4$, by Proposition 6.2. Therefore, using Theorem 3.1, the ideal $(v_8, v_{12}, \dots) \subset \Omega_*^{\text{SO}}$ is contained in A_* . An analogous argument shows

that the ideal (Z) (Z as in Theorem 6.1) is contained in A_* . The only additive generators of Ω_*^{SO} in dimensions ≥ 5 which are missed by that argument are the elements of the form

$$z_5^a v_4^b, \quad 5a + 4b \geq 5.$$

The manifold M in Proposition 5.8 has an admissible splitting and since $[M] \neq 0$ and $\Omega_5^{\text{SO}} \cong \mathbb{Z}/2$, we have $[M] = z_5$ and so $z_5 \in A_5$.

Propositions 5.8, together with Proposition 5.1, implies that $z_5 x \in A_*$ when $x \in \Omega_*^{\text{SO}}$ is of positive dimension.

For $b \geq 3$, we write $v_4^b = [M \times \mathbb{C}\mathbb{P}^2]$ with $\dim(M) \geq 6$ and use Proposition 5.1 to verify that $v_4^b \in A_*$.

These arguments cover all cases, except $d = 8$. In that case, we certainly have $[\mathbb{H}\mathbb{P}^2], [\mathbb{C}\mathbb{P}^4] \in A_8$, and therefore the index of A_8 in Ω_8^{SO} divides 3, by Theorem 6.1. Hence if $[M] \in \Omega_8^{\text{SO}}$, M is cobordant modulo A_8 to either W^8 , $\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2$, or $\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2$ with the reverse orientation, and so if M is simply connected and not spinnable, $\mathcal{R}^+(M)$ has the homotopy type of either $\mathcal{R}^+(W^8)$ or $\mathcal{R}^+(\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2)$. \square

6.2. The structure of the oriented cobordism ring. This section is a guide through the literature on oriented cobordism. For a commutative ring R , a class $c \in H^*(BO; R)$ and $x \in \Omega_*^{\text{SO}}$, we denote by $c(x) \in R$ the characteristic number $c(x) = \langle c(TM), [M] \rangle \in R$, where $[M] = x$. Recall furthermore the classes $s_n \in H^{4n}(BO; \mathbb{Z})$ from [30, §16].

Theorem 6.4. *Let $T_* \subset \Omega_*^{\text{SO}}$ be the ideal of torsion elements. Assume that $v_{4k} \in \Omega_{4k}^{\text{SO}}$, $k \geq 1$, are classes such that*

$$(6.5) \quad s_k(v_{4k}) = \begin{cases} 1 & 2k+1 \text{ not a prime power} \\ p & 2k+1 = p^s, s > 0, p \text{ prime,} \end{cases}$$

and let $\overline{v_{4k}} \in \Omega_{4k}^{\text{SO}}/T_{4k}$ be the residue class in the torsionfree quotient. Then

$$\Omega_*^{\text{SO}}/T_* = \mathbb{Z}[\overline{v_{4k}} | k > 0].$$

Furthermore, $2T_* = 0$.

The computation of Ω_*^{SO}/T_* is apparently due to Milnor (unpublished, but see [38], p. 207 for an account). Milnor [29, Theorem 5] also proved that there is no odd primary torsion in Ω_*^{SO} , and Wall [40, Theorem 2] showed that there is no element of order 4.

Proposition 6.6. *One can choose $v_4 = [\mathbb{C}\mathbb{P}^2]$ and $v_8 = [\mathbb{C}\mathbb{P}^4]$. For $k \geq 3$, one can choose v_{4k} as a linear combination of total spaces of $\mathbb{C}\mathbb{P}^n$ -bundles with structure group $U(n+1)$, with $n \geq 3$.*

Proof. If $2k+1$ is a prime, we can take $v_{4k} = [\mathbb{C}\mathbb{P}^{2k}]$, because

$$s_k(\mathbb{C}\mathbb{P}^{2k}) = 2k+1,$$

see [23], p. 42. This applies to $k = 1, 2$. If $k \geq 3$ and $2k+1$ is not a prime, we have to use further manifolds, the *Milnor manifolds*. To define them, let $i \leq j$. The tautological line bundle $L_i \rightarrow \mathbb{C}\mathbb{P}^i$ embeds canonically into the trivial vector bundle $\underline{\mathbb{C}}^{j+1}$, and we let L_{ij}^\perp be the orthogonal complement of L_i in $\underline{\mathbb{C}}^{j+1}$. The Milnor manifold is

$$H_{ij} := \mathbb{P}(L_{ij}^\perp).$$

By construction, H_{ij} is a $\mathbb{C}\mathbb{P}^{j-1}$ -bundle over $\mathbb{C}\mathbb{P}^i$ with structure group $U(j)$. Note that $H_{0j} = \mathbb{C}\mathbb{P}^{j-1}$. The manifolds H_{1j} are nullbordant. If $i \geq 2$ and $i + j - 1 = 2k$, we have

$$s_k(H_{ij}) = -\binom{i+j}{i} = -\binom{2k+1}{i},$$

see [23], p. 43.

For $k \geq 3$, $\mathbb{C}\mathbb{P}^{2k}$ as well as H_{ij} with $i + j = 2k + 1$, $i \geq 2$ and $j \geq 4$, are bundles of projective space as asserted. Therefore, by taking a suitable linear combination of these manifolds, we get a class $v_{4k} \in \Omega_{4k}^{\text{SO}}$ with

$$s_k(v_{4k}) = \gcd\{2k+1, \binom{2k+1}{2k+1-j}, 4 \leq j \leq 2k-1\}.$$

If $k \geq 3$, this number is the same as

$$d_{2k+1} := \gcd\{\binom{2k+1}{n} \mid 1 \leq n \leq 2k\}.$$

It is a rather unknown, but elementary fact [33] that

$$\gcd\{\binom{n}{j} \mid 1 \leq j \leq n-1\} = \begin{cases} 1 & n \text{ not a prime power} \\ p & n = p^s, s > 0, p \text{ prime.} \end{cases}$$

From that, the claim follows. \square

Wall also determined the structure of T_* to which we turn now. We denote by $[M]^O \in \Omega_n^O$ the unoriented cobordism class of a closed n -manifold M . Following [40], let $\mathfrak{W}_* \subset \Omega_*^O$ be the set of all (unoriented) cobordism classes which have representatives M such that the Stiefel-Whitney class $w_1(TM) \in H^1(M; \mathbb{Z}/2)$ are the (mod 2) reduction of an integral class $w \in H^1(M; \mathbb{Z}) = [M; S^1]$. This is a subalgebra of Ω_*^O [40, Lemma 3], and the forgetful map $\rho : \Omega_*^{\text{SO}} \rightarrow \Omega_*^O$ factors through \mathfrak{W}_* .

Let $u : M^n \rightarrow S^1$ be a smooth map representing w and pick a regular value a of u . The manifold $V := u^{-1}(a)$ is orientable, and $2[V] = 0 \in \Omega_{n-1}^{\text{SO}}$ [40, Lemma 1]. Theorem 3 of [40] says that the assignment $M \mapsto V$ gives a well-defined map

$$\partial : \mathfrak{W}_* \rightarrow \Omega_{*-1}^{\text{SO}},$$

which is a derivation and fits into a long exact sequence

$$\dots \rightarrow \Omega_n^{\text{SO}} \xrightarrow{2} \Omega_n^{\text{SO}} \xrightarrow{\rho} \mathfrak{W}_n \xrightarrow{\partial} \Omega_{n-1}^{\text{SO}} \rightarrow \dots$$

This and the fact that $2T_* = 0$ implies $T_* = \text{Im}(\partial)$. For the description of the structure of \mathfrak{W}_* , we need the following manifolds. Let $P(m, n)$ be the Dold manifold $S^m \times_{C_2} \mathbb{C}\mathbb{P}^n \rightarrow \mathbb{R}\mathbb{P}^m$, where C_2 acts on $\mathbb{C}\mathbb{P}^n$ by complex conjugation. Furthermore, let $A : P(m, n) \rightarrow P(m, n)$ be the map induced by $(x, z) \mapsto (Rx, z)$, where R is a reflection along a hyperplane. We define $Q(m, n)$ as the mapping torus of A ; there is an obvious fibre bundle $u : Q(m, n) \rightarrow S^1$ with fibre $P(m, n)$. If m is odd and n is even, $P(m, n)$ is orientable, and A reverses orientation, and u defines an integral lift of $w_1(TQ(m, n))$ in this situation. Hence $[P(m, n)]^O$ and $[Q(m, n)]^O$ are elements of \mathfrak{W}_* , if m is odd and n is even.

Theorem 6.7. *If k is not a power of 2, write $k = 2^{r-1}(2s + 1)$, $s \neq 0$, and define*

$$y_{2k-1} := [P(2^r - 1, 2^r s)]^O \in \mathfrak{W}_{2k-1}$$

and

$$y_{2k} := [Q(2^r - 1, 2^r s)]^O \in \mathfrak{W}_{2k}.$$

Let furthermore $y_{2^r} := [\mathbb{C}\mathbb{P}^{2^r-1}]^O \in \mathfrak{W}_{2^r}$. Then \mathfrak{W}_* is a polynomial algebra over \mathbb{F}_2 , with generators the elements y_{2k}, y_{2k-1} (k not a power of 2) and y_{2^r} , $r \geq 2$.

Short guide through the proof. This is shown in [40, §5-6], but not explicitly stated there, hence we recall the argument, following the streamlined version [3] of Wall's proof by Atiyah. Recall that Thom [39, Théorème IV.12] proved that Ω_*^O is a polynomial algebra over \mathbb{F}_2 , with one generator x_i for each i which is not of the form $2^r - 1$, see also [38, p. 96]. Combining Thom's theorem with [10, Satz 3] and the proof of [40, Lemma 6], the elements in the theorem are algebraically independent, and hence generate a polynomial subalgebra $\mathfrak{V}_* \subset \mathfrak{W}_*$. The Hilbert–Poincaré series $HS_{\mathfrak{W}_*}(t) = \sum_{n=0}^{\infty} \dim_{\mathbb{F}_2}(\mathfrak{W}_n) t^n \in \mathbb{Z}[[t]]$ is easily calculated.

Using formula (9) of [3] and the structure of Ω_*^O , one computes the Hilbert–Poincaré series $HS_{\mathfrak{W}_*}(t)$ and verifies that both series agree, whence $\mathfrak{V}_* = \mathfrak{W}_*$. \square

The following finishes the description of the multiplicative generators of Ω_*^{SO} .

Corollary 6.8. *Let $x \in T_*$ be a torsion element in Ω_*^{SO} . Then x is a linear combination of manifolds, which are either*

- (1) $\mathbb{C}\mathbb{P}^n$ -bundles with structure group C_2 and $n \geq 3$,
- (2) or products of $P(1, 2)$ with some other oriented manifold.

Proof. Since $T_* = \text{Im}(\partial)$, it is enough to check that all monomials in the generators of \mathfrak{W}_* , are mapped to cobordism classes of the claimed form under ∂ .

Let k, r and s as in Theorem 6.7. The elements y_i introduced there have the preimages

$$z_{2k-1} := [P(2^r - 1, 2^r s)] \in \Omega_{2k-1}^{\text{SO}}$$

and

$$z_{2^r} := [\mathbb{C}\mathbb{P}^{2^r-1}] \in \Omega_{2^r}^{\text{SO}}$$

under $\rho : \Omega_*^{\text{SO}} \rightarrow \mathfrak{W}_*$. Therefore $\partial(y_{2k-1}) = 0$ and $\partial(y_{2^r}) = 0$.

Since $\partial : \mathfrak{W}_* \rightarrow \Omega_*^{\text{SO}}$ is a derivation

$$\partial\left(\prod_i y_{2k_i-1}^{a_i} \prod_i y_{2^r_i}^{b_i} \prod_i y_{2k_i}^{c_i}\right) = \left(\prod_i y_{2k_i-1}^{a_i}\right) \left(\prod_i y_{2^r_i}^{b_i}\right) \partial\left(\prod_i y_{2k_i}^{c_i}\right).$$

The Dold manifold $P(2^r - 1, 2^r s)$ is a $\mathbb{C}\mathbb{P}^{2^r s}$ -bundle, and $2^r s \geq 2$, with equality only if $r = s = 1$, i.e. $k = 3$. The element z_5 is represented by $P(1, 2)$. For all other values of k , z_{2k-1} is represented by a bundle of complex projective spaces of dimension at least 3.

Furthermore, $\partial(\prod_i y_{2k_i}^{c_i})$ is represented by a manifold V of the following type. Take a product $Q(m_1, n_1) \times \dots \times Q(m_p, n_p) \rightarrow (S^1)^i$, which is a fibre bundle with fibre a product of Dold manifolds, restrict this fibre bundle to the kernel $\Delta \subset (S^1)^i$ of the multiplication map $\mu : (S^1)^p \rightarrow S^1$, and take the total space.

Now there is a fibre bundle $Q(m, n) \rightarrow Q(m, 0)$ with fibre $\mathbb{C}\mathbb{P}^n$ and structure group C_2 (complex conjugation). This is compatible with the map to S^1 . Therefore, V is the total space of a bundle of projective spaces with structure group C_2 . The complex dimension of the fibre can be chosen to be at least 3, except in the case of y_6^c . If c is even, then $\partial(y_6^c) = 0$, and furthermore

$$\partial(y_6^{2m+1}) = z_5 z_6^{2m}.$$

The claim follows. \square

From the description of the generators, it is also apparent that $\Omega_5^{\text{SO}} \cong \mathbb{Z}/2$, that $\Omega_6^{\text{SO}} = \Omega_7^{\text{SO}} = 0$ and that Ω_8^{SO} has the \mathbb{Z} -basis $[\mathbb{C}\mathbb{P}^4]$ and $[\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2]$. Moreover

$$[\text{HP}^2] = 3[\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2] - 2[\mathbb{C}\mathbb{P}^4] :$$

both sides of the equation have the same signature; furthermore $s_2([\mathbb{C}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^2]) = 0$, $s_2([\mathbb{C}\mathbb{P}^4]) = 5$ and $s_2([\text{HP}^2]) = -10$, which is easily derived from the computations in [23], p.5 f. Therefore, $[\mathbb{C}\mathbb{P}^4]$ and $[\text{HP}^2]$ span a subgroup of index 3 in Ω_8^{SO} .

REFERENCES

- [1] J. F. Adams, *Stable homotopy and generalised homology*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1995. Reprint of the 1974 original. MR1324104
- [2] D. W. Anderson, E. H. Brown Jr., and F. P. Peterson, *The structure of the Spin cobordism ring*, Ann. of Math. (2) **86** (1967), 271–298. MR219077
- [3] M. F. Atiyah, *Bordism and cobordism*, Proc. Cambridge Philos. Soc. **57** (1961), 200–208. MR126856
- [4] D. Barden, *Simply connected five-manifolds*, Ann. of Math. (2) **82** (1965), 365–385. MR184241
- [5] J. C. Becker and D. H. Gottlieb, *The transfer map and fiber bundles*, Topology **14** (1975), 1–12. MR0377873
- [6] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 10, Springer-Verlag, Berlin, 1987. MR867684
- [7] B. Botvinnik, J. Ebert, and O. Randal-Williams, *Infinite loop spaces and positive scalar curvature*, Invent. Math. **209** (2017), no. 3, 749–835.
- [8] V. Chernysh, *On the homotopy type of the space $\mathcal{R}^+(M)$* , 2004. arXiv:math/0405235.
- [9] ———, *A quasifibration of spaces of positive scalar curvature metrics*, Proc. Amer. Math. Soc. **134** (2006), no. 9, 2771–2777 (electronic).
- [10] A. Dold, *Erzeugende der Thomschen Algebra \mathfrak{A}* , Math. Z. **65** (1956), 25–35. MR79269
- [11] J. L. Dupont, *Curvature and characteristic classes*, Lecture Notes in Mathematics, Vol. 640, Springer-Verlag, Berlin-New York, 1978. MR0500997
- [12] J. Ebert and G. Frenck, *The Gromov-Lawson-Chernysh surgery theorem*, 2018. ArXiv:1807.06311, to appear in Boletín de la Sociedad Matemática Mexicana.
- [13] J. Ebert and O. Randal-Williams, *Infinite loop spaces and positive scalar curvature in the presence of a fundamental group*, Geom. Topol. **23** (2019), no. 3, 1549–1610. MR3956897
- [14] J. Ebert and O. Randal-Williams, *The positive scalar curvature cobordism category*, 2019. arXiv:1904.12951.
- [15] S. Fühling, *Bordism und projective space bundles*, 2020. arXiv:2006.15394.
- [16] A. Gray, *Pseudo-Riemannian almost product manifolds and submersions*, J. Math. Mech. **16** (1967), 715–737. MR0205184
- [17] D. Gromoll and G. Walschap, *Metric foliations and curvature*, Progress in Mathematics, vol. 268, Birkhäuser Verlag, Basel, 2009. MR2500106
- [18] M. Gromov and H. B. Lawson Jr., *The classification of simply connected manifolds of positive scalar curvature*, Ann. of Math. (2) **111** (1980), no. 3, 423–434.
- [19] B. Hanke, T. Schick, and W. Steimle, *The space of metrics of positive scalar curvature*, Publ. Math. Inst. Hautes Études Sci. **120** (2014), 335–367. MR3270591
- [20] F. Hebestreit and M. Joachim, *Twisted spin cobordism and positive scalar curvature*, J. Topol. **13** (2020), no. 1, 1–58.
- [21] N. Higson and G. Kasparov, *E-theory and KK-theory for groups which act properly and isometrically on Hilbert space*, Invent. Math. **144** (2001), no. 1, 23–74. MR1821144
- [22] F. Hirzebruch, *Über eine Klasse von einfachzusammenhängenden komplexen Mannigfaltigkeiten*, Math. Ann. **124** (1951), 77–86. MR45384

- [23] F. Hirzebruch, T. Berger, and R. Jung, *Manifolds and modular forms*, Aspects of Mathematics, E20, Friedr. Vieweg & Sohn, Braunschweig, 1992. With appendices by Nils-Peter Skoruppa and by Paul Baum. MR1189136
- [24] M. A. Kervaire, *Le théorème de Barden-Mazur-Stallings*, Comment. Math. Helv. **40** (1965), 31–42. MR0189048
- [25] S. O. Kochman, *Bordism, stable homotopy and Adams spectral sequences*, Fields Institute Monographs, vol. 7, American Mathematical Society, Providence, RI, 1996. MR1407034
- [26] J.-B. Kordaß, *On the space of riemannian metrics satisfying surgery stable curvature conditions*, 2018. ArXiv 1808.00581.
- [27] M. Kreck and S. Stolz, *HP²-bundles and elliptic homology*, Acta Math. **171** (1993), no. 2, 231–261. MR1251581
- [28] G. Lusztig, J. Milnor, and F. P. Peterson, *Semi-characteristics and cobordism*, Topology **8** (1969), 357–359. MR246308
- [29] J. Milnor, *On the cobordism ring Ω^* and a complex analogue. I*, Amer. J. Math. **82** (1960), 505–521. MR119209
- [30] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J., 1974. Annals of Mathematics Studies, No. 76. MR0440554 (55 #13428)
- [31] M. Mimura and H. Toda, *Topology of Lie groups. I, II*, Translations of Mathematical Monographs, vol. 91, American Mathematical Society, Providence, RI, 1991. Translated from the 1978 Japanese edition by the authors. MR1122592
- [32] B. O’Neill, *The fundamental equations of a submersion*, Michigan Math. J. **13** (1966), 459–469. MR200865
- [33] B. Ram, *Common factors of $\frac{n!}{m!(n-m)!}$, ($m = 1, 2, \dots, n-1$)*, J. Indian Math. Club **1** (1909), no. 2, 39–43. MR3821621
- [34] T. Schick, *A counterexample to the (unstable) Gromov-Lawson-Rosenberg conjecture*, Topology **37** (1998), no. 6, 1165–1168. MR1632971
- [35] ———, *Real versus complex K-theory using Kasparov’s bivariant KK-theory*, Algebr. Geom. Topol. **4** (2004), 333–346. MR2077669
- [36] R. Schoen and S. T. Yau, *On the structure of manifolds with positive scalar curvature*, Manuscripta Math. **28** (1979), no. 1-3, 159–183. MR535700
- [37] S. Stolz, *Simply connected manifolds of positive scalar curvature*, Ann. of Math. (2) **136** (1992), no. 3, 511–540. MR1189863
- [38] R. E. Stong, *Notes on cobordism theory*, Mathematical notes, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1968. MR0248858
- [39] R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment. Math. Helv. **28** (1954), 17–86. MR61823
- [40] C. T. C. Wall, *Determination of the cobordism ring*, Ann. of Math. (2) **72** (1960), 292–311. MR120654
- [41] ———, *Finiteness conditions for CW-complexes*, Ann. of Math. (2) **81** (1965), 56–69. MR0171284
- [42] ———, *Geometrical connectivity. I*, J. London Math. Soc. (2) **3** (1971), 597–604.
- [43] M. Walsh, *Cobordism invariance of the homotopy type of the space of positive scalar curvature metrics*, Proc. Amer. Math. Soc. **141** (2013), no. 7, 2475–2484.
- [44] M. Walsh, *Metrics of positive scalar curvature and generalised Morse functions, Part I*, Mem. Amer. Math. Soc. **209** (2011), no. 983, xviii+80. MR2789750

Email address: johannes.ebert@uni-muenster.de

Email address: wiemelerm@uni-muenster.de