

A note on an alleged proof of the relative consistency of $P = NP$ with PA

Ralf-Dieter Schindler

Institut für formale Logik, Universität Wien, 1090 Wien, Austria

rds@logic.univie.ac.at

<http://www.logic.univie.ac.at/~rds/>

N.C.A. da Costa and F.A. Doria claim to have shown in [1] that $P = NP$ is relatively consistent with PA . The purpose of the present note is to argue that there is a mistake in that paper. Specifically, we want to point out that Corollary 5.14 of [1] – which is used in the proof of their main result – is probably false.

We are first going to reconstruct their argument. According to that reconstruction, the argument of [1] would in fact show that PA proves $P = NP$. We'll then discuss Cor. 5.14 of [1]. However, rather than talking about provability in PA or stronger theories, we'll stick to a different attitude and argue internally¹: using Cor. 5.14 of [1] we'll derive a contradiction from the assumption that $P < NP$ (and we implicitly assume our argument goes thru in PA).

We'll follow the notation of [1] (with the exception of f_{-A}).

Some Turing machines. $\bigvee(z) = 1$ iff $\pi_1(z)$ codes a cnf-Boolean expression and $\pi_2(z)$ codes an assignment which satisfies it (o.w. $\bigvee(z) = 0$). E is a fixed exponential Turing machine that solves any instance of the satisfiability problem (in particular, $\bigvee(\langle z, E(z) \rangle) = 1$ for any z coding a satisfiable cnf-Boolean expression). Let G be a partial recursive function; for $n < \omega$ such that $G(n) \downarrow$ we let $Q^{G(n)}$ be that Turing machine which, given an input z , first computes $G(n)$ and then computes $E(z)$ in case $z \leq G(n)$ and 0 in case $z > G(n)$ (in particular, $Q^{G(n)}(z) = E(z)$ for $z \leq G(n)$ and $Q^{G(n)}(z) = 0$ for $z > G(n)$; cf. p. 10 of [1]). Notice \bigvee and all $Q^{G(n)}$ (for $n < \omega$ and $G(n) \downarrow$) are polynomial time Turing machines (in fact the $Q^{G(n)}$'s are "finite"). By the Baker-Gill-Solovay trick there is a recursive enumeration $(P_m: m < \omega)$ of all polynomial Turing machines; we may think of (the "Gödel number" of) P_m as an ordered pair $\langle g, c \rangle$ where g codes P_m 's program and c is a code for a polynomial clock.

¹Argumenting externally might be more in line with the paper itself. However, such a reconstruction typically starts from a listing of Turing machines which is such that PA won't be able to prove " $P < NP \Leftrightarrow f$ is total." (The definition of f is given in the main text.)

Some functions. We let $f(m)$ be the least z such that $\bigvee(z) = 1$, whereas $\bigvee(\langle \pi_1(z), P_m(\pi_1(z)) \rangle) = 0$ (i.e., $f(m)$ witnesses that P_m doesn't prove $P = NP$). We have: f is recursive, and f is total iff $P < NP$. (f is written $f_{\neg A}$ in [1]; cf. [1] p. 4.) Let $(\psi_i: i < \omega)$ be a recursive enumeration of all polynomial functions from ω to ω . We let $F(m) = \max\{f \circ \psi_i(m): m \leq i \wedge f \circ \psi_i(m) \downarrow\} + 1$. (We understand that $\max \emptyset = 0$.) Note that F dominates $f \circ \psi$ (in the sense that $F(m) > f \circ \psi(m)$ for all sufficiently large m with $f \circ \psi_i(m) \downarrow$) for any polynomial ψ . If f is total then F is recursive.

Corollary 5.14 of [1] now reads as follows: **Main Lemma.** If F is recursive, then there is a *linear* $\psi: \omega \rightarrow \omega$ s.t. for all m and n do we have that $Q^{F(m)}(n) = P_{\psi(m)}(n)$. Let us also consider the following, which trivially follows from this Main Lemma: **Main Lemma'.** If F is recursive, then there is a *polynomial* $\psi: \omega \rightarrow \omega$ s.t. for all m and n do we have that $Q^{F(m)}(n) = P_{\psi(m)}(n)$.

Given the Main Lemma' we may now prove $P = NP$ as follows. Suppose not. Then f is total recursive, and hence so is F . If ψ is as in the Main Lemma' then $F(m) > f \circ \psi(m)$ for all sufficiently large m . On the other hand, $f(\psi(m))$ is the least z such that $\bigvee(z) = 1$, whereas $\bigvee(\langle \pi_1(z), Q^{F(m)}(\pi_1(z)) \rangle) = 0$. For $\pi_1(z) \leq F(m)$ we'll have $Q^{F(m)}(\pi_1(z)) = E(\pi_1(z))$, so that $f(\psi(m)) \geq \langle F(m) + 1, 0 \rangle \geq F(m) + 1$. We'll thus have $F(m) > f \circ \psi(m) \geq F(m) + 1$ for all sufficiently large m . Contradiction! We have shown that $P = NP$.

Have we? Not so, I claim. Let's discuss the Main Lemma. The Baker-Gill-Solovay trick uses the device of "clocks" in order to arrive at a recursive enumeration $(P_m: m < \omega)$ of all polynomial Turing machines. I.e. (cf. above), the Gödel number of $Q^{F(m)}$, viewed as a machine with a clock attached to it, will be – typically – the ordered pair $\langle g, c \rangle = \langle g(m), c(m) \rangle$ of a code g for $Q^{F(m)}$'s program (without a clock) and a code c for a polynomial clock. Now the clock is not supposed to shut down $Q^{F(m)}$'s operation before $F(m)$ gets known. That is, c depends on the length of the computation of $F(m)$; in fact, if F is "complicated," $c \approx$ the length of the computation of $F(m)$. Hence the function $m \mapsto$ Gödel number $\langle g(m), c(m) \rangle$ of $Q^{F(m)}$ will be at least as "complex" as $m \mapsto$ the length of the computation of $F(m)$. There is no reason to believe that it should be linear (or, polynomial, for that matter). In other words, any function ψ s.t. $Q^{F(m)}(n) = P_{\psi(m)}(n)$ for all m and n will be as "complex" as F is; which is, I think, as it should be. But this then poses a serious problem. Suppose a ψ as above can only be as "complex" as F . The above argument breaks down without an F s.t. F dominates $f \circ \psi$.

It is hard to believe that there should be a proof of the Main Lemma' which doesn't actually prove (without assuming $P < NP$) the following. (*) For any total recursive G there is a polynomial $\psi: \omega \rightarrow \omega$ s.t. for all m and n do we have that $Q^{G(m)}(n) = P_{\psi(m)}(n)$. But (*) is false, as we shall now show. Recall that $Q^{G(m)}$

always first computes $G(m)$; in this sense, the value $G(m)$ “shows up (on the tape) during the calculation of $Q^{G(m)}(n)$,”² for every m and n . Now let us consider

$$G(m) = \max\{t : t \text{ shows up during the calculation of } P_i(m) \text{ for } i \leq m^m\} + 1.$$

Clearly, G is total recursive. Suppose that ψ is as in $(*)$ for G . In particular, for every m , $G(m)$ shows up during the calculation of $Q^{G(m)}(m)$. However, pick m s.t. $\psi(m) \leq m^m$. Then by construction, $G(m) > t$ for all t which show up in the calculation of $P_{\psi(m)}(m) = Q^{G(m)}(m)$. Contradiction!³

I don't understand the alleged proof of Cor. 5.14 of [1] which appears on pp. 13 ff. of [1]. Specifically, I don't understand Remark 5.11 on p. 14 of [1]. I think it contains a statement (“Again we have, [...]”) which has not been verified. The statement is reminiscent to $(*)$ above.

We believe that at some point s.o. will prove $P < NP$ by some form of “Galois theory.”

I cordially thank N.C.A. da Costa and Chico Doria for their interest in my e-mail messages concerning [1].

References

- [1] N.C.A. da Costa and F.A. Doria, *On the consistency of $P = NP$ with fragments of ZFC whose own consistency strength can be measured by an ordinal assignment*, <http://arXiv.org/abs/math/0006079>.

²We leave it to the reader to make this precise.

³Another way to look at [1] is the following. If the argument of [1] worked, we could re-run it by talking about finite time machines rather than polynomial time machines (by finite time I mean no matter how long the input is the machine will stop after c steps, where c is a constant being independent from the length of the input). All $Q^{F(m)}$'s are finite time. So by starting from an enumeration of finite time machines rather than of polynomial time machines the argument should really prove (the relative consistency of) “finite” = NP rather than just of $P = NP$, which truly is absurd.