

# $\Pi_2$ Consequences of $\mathbf{BMM} + \mathbf{NS}_{\omega_1}$ is precipitous and the semiproperness of stationary set preserving forcings.

Philipp Doebler, Ralf Schindler\*  
Institut für mathematische Logik und Grundlagenforschung  
Universität Münster  
Einsteinstr. 62, 48149 Münster, Germany

February 16, 2009

## Abstract

We investigate which  $\Pi_2$  sentences (over  $H_{\omega_2}$ ) that are consequences of  $\mathbf{MM}$  also follow from  $\mathbf{BMM} + \mathbf{NS}_{\omega_1}$  is precipitous. It turns out that admissible club guessing ( $\mathbf{acg}$ ),  $\delta_2^1 = \omega_2$ , the club bounding principle ( $\mathbf{CBP}$ ), and  $\psi_{AC}$  as well as  $\phi_{AC}$  follow from this weaker theory. This was known for  $\delta_2^1 = \omega_2$  and  $\psi_{AC}$  but not for  $\phi_{AC}$  and  $\mathbf{acg}$ . Additionally we show that if for all regular  $\theta \geq \omega_2$  there is a semiproper partial ordering that adds a generic iteration of length  $\omega_1$  with last model  $H_\theta$ , then all stationary set preserving forcings are semiproper.

## 1 Introduction

By  $\mathbf{NS}_{\omega_1}$  we denote the nonstationary ideal on  $\omega_1$ . A  $V$ -generic  $G$  for the forcing  $(\mathcal{P}(\omega_1) \setminus \mathbf{NS}_{\omega_1}, \subset)$  is an ultrafilter on  $V$  that extends the club filter. Hence we can form the ultrapower  $j : V \rightarrow \text{Ult}(V, G)$  in  $V[G]$ . We will always assume the well-founded part of such an ultrapower to be transitive. Clearly  $j$  has critical point  $\omega_1$ . If every condition  $S \in \mathcal{P}(\omega_1) \setminus \mathbf{NS}_{\omega_1}$  forces that  $\text{Ult}(V, G)$  is well-founded, then we call  $\mathbf{NS}_{\omega_1}$  precipitous. Since the precipitousness of an ideal can be recast as a first order statement, the model  $\text{Ult}(V, G)$  has a precipitous nonstationary ideal if  $V$  has one. One can now pick a  $\text{Ult}(V, G)$ -generic for  $(\mathcal{P}(\omega_1) \setminus \mathbf{NS}_{\omega_1}, \subset)^{\text{Ult}(V, G)}$  and form another ultrapower. This leads to the notion of generic iterations.

**Definition 1.1** Let  $M$  be a transitive model of  $\mathbf{ZFC}^- + \text{“}\omega_1 \text{ exists,“}$  and let  $I \subseteq \mathcal{P}(\omega_1^M)$  be such that  $\langle M; \in, I \rangle \models \text{“}I \text{ is a uniform and normal ideal on } \omega_1^M \text{.”}$  Let  $\gamma \leq \omega_1$ . Then

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \gamma \rangle, \langle G_i; i < \gamma \rangle \rangle \in V$$

is called a *putative generic iteration of  $\langle M; \in, I \rangle$  (of length  $\gamma + 1$ )* iff the following hold true.

1.  $M_0 = M$  and  $I_0 = I$ .
2. For all  $i \leq j \leq \gamma$ ,  $\pi_{i,j} : \langle M_i; \in, I_i \rangle \rightarrow \langle M_j; \in, I_j \rangle$  is elementary,  $I_i = \pi_{0,i}(I)$ , and  $\kappa_i = \pi_{0,i}(\omega_1^M) = \omega_1^{M_i}$ .

---

\*Both authors gratefully acknowledge support by DFG grant no. SCHI 484/3-1.

3. For all  $i < \gamma$ ,  $M_i$  is transitive and  $G_i$  is  $(\mathcal{P}(\kappa_i) \setminus I_i, \subset)$ -generic over  $M_i$ .
4. For all  $i + 1 \leq \gamma$ ,  $M_{i+1} = \text{Ult}(M_i; G_i)$  and  $\pi_{i,i+1}$  is the associated ultrapower map.
5.  $\pi_{j,k} \circ \pi_{i,j} = \pi_{i,k}$  for  $i \leq j \leq \gamma$ .
6. If  $\lambda \leq \gamma$  is a limit ordinal, then  $\langle M_\lambda, \pi_{i,\lambda}, i < \lambda \rangle$  is the direct limit of  $\langle M_i, \pi_{i,j}, i \leq j < \lambda \rangle$ .

We call

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \gamma \rangle, \langle G_i; i < \gamma \rangle \rangle$$

a *generic iteration* of  $\langle M; \in, I \rangle$  (of length  $\gamma + 1$ ) iff it is a putative generic iteration of  $\langle M; \in, I \rangle$  of length  $\gamma + 1$  and  $M_\gamma$  is transitive.  $\langle M; \in, I \rangle$  is *generically  $\gamma$  iterable* iff for any  $\gamma \leq \omega_1$  every putative generic iteration of  $\langle M; \in, I \rangle$  of length  $\gamma + 1$  is an iteration.

Notice that we want (putative) iterations of a given model  $\langle M; \in, I \rangle$  to exist in  $V$ , which amounts to requiring that the relevant generics  $G_i$  may be found in  $V$ .

In [CS09] the notion of forcing  $\mathbb{P}(\theta, \text{NS}_{\omega_1})$  was defined for regular  $\theta \geq \omega_2$ . Granted the precipitousness of nonstationary ideal  $\text{NS}_{\omega_1}$  the forcing is nonempty and preserves stationary subsets of  $\omega_1$ . Forcing with  $\mathbb{P}(\theta, \text{NS}_{\omega_1})$  adds a generic iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

such that all  $M_i$  with countable index are countable and the last model  $M_{\omega_1}$  equals  $H_\theta$ . Here  $I_i$  is  $M_i$ 's nonstationary ideal and the  $\kappa_i = \omega_1^{M_i}$  are the critical points of the generic ultrapowers  $\pi_{i,i+1} : M_i \rightarrow M_{i+1} \simeq \text{Ult}(M_i, G_i)$ . It is also possible to produce iterations as above with generically iterable  $M_0$ . This fact is used in [CS09] to show that  $\text{BMM} + \text{NS}_{\omega_1}$  is precipitous implies  $\delta_2^1 = \omega_2$ . Note that  $\delta_2^1 = \omega_2$  is a  $\Pi_2$  statement in  $H_{\omega_2}$ . In this paper we use generic iterations as above to analyse which  $\Pi_2$  sentences in  $H_{\omega_2}$  that are consequences of  $\text{ZFC} + \text{MM}$  are also consequences of the weaker theory  $\text{ZFC} + \text{BMM} + \text{NS}_{\omega_1}$  is precipitous. Note that  $\text{MM}$  implies that  $\text{NS}_{\omega_1}$  is  $\omega_2$ -saturated [FMS88] but by [Woo99, 10.103, 10.99]  $\text{BMM} + \text{NS}_{\omega_1}$  is precipitous does not<sup>1</sup>. We consider two  $\Pi_2$  statements in  $H_{\omega_2}$ . Both are known to hold in  $H_{\omega_2}$  if  $\text{MM}$  holds.

**Definition 1.2** 1. We call the following principle *admissible club guessing* (**acg**). For all clubs  $C \subseteq \omega_1$  there exists a real  $x$  such that

$$A_x := \{ \alpha < \omega_1; L_\alpha[x] \text{ is admissible} \} \subset C.$$

2. Let  $S \subset \omega_1$ . Then we set

$$\tilde{S} := \{ \alpha < \omega_2; \omega_1 \leq \alpha \wedge \mathbf{1}_{\mathbb{B}} \Vdash \check{\alpha} \in j(\check{S}) \},$$

where  $\mathbb{B} = \text{ro}(\mathcal{P}(\omega_1) \setminus \text{NS}_{\omega_1})$  and  $j$  is a name for the corresponding generic elementary embedding  $V \rightarrow (M, E) \subset V^{\mathbb{B}}$ . Note that  $\alpha \in \tilde{S}$  if and only if for all (one) canonical function(s)  $f_\alpha$  for  $\alpha$ , there is a club  $C$  such that if  $\beta \in C$  then  $f_\alpha(\beta) \in S$ .

Let  $\vec{S} = \langle S_i; i \in \omega \rangle$ ,  $\vec{T} = \langle T_i; i \in \omega \rangle$  be sequences of pairwise disjoint subsets of  $\omega_1$ , such that all  $S_i$  are stationary and

$$\omega_1 = \bigcup \{ T_i; i \in \omega \}.$$

$\varphi_{AC}(\vec{S}, \vec{T})$  is the conjunction of the following two statements:

<sup>1</sup>In the situation of [Woo99, 10.103] one considers a  ${}^2\mathbb{P}_{\max}$  extension; there  $\text{NS}_{\omega_1}$  is not saturated but one can check that it is precipitous using the  ${}^2\mathbb{P}_{\max}$  analysis in [Woo99, 6.14].

- (a) There is an  $\omega_1$  sequence of distinct reals.<sup>2</sup>
- (b) There is  $\gamma < \omega_2$  and a continuous increasing function  $F : \omega_1 \rightarrow \gamma$  with range cofinal in  $\gamma$  such that for all  $i \in \omega$

$$F \text{``} T_i \subset \tilde{S}_i.$$

$\varphi_{AC}(\vec{S}, \vec{T})$  is clearly  $\Sigma_1(\{\vec{S}, \vec{T}\})$  in  $\langle H_{\omega_2}; \in \rangle$ . We set

$$\phi_{AC} \equiv \forall \vec{S} \forall \vec{T} \varphi_{AC}(\vec{S}, \vec{T}).$$


Note that  $\phi_{AC}$  is equivalent to a  $\Pi_2$  statement in  $\langle H_{\omega_2}; \in \rangle$ .

**Remark 1.3** The principle **acg** was isolated by Woodin. If **MM** holds, then the universe is closed under the sharp operation (this is already a consequence of **BMM**). So by [Woo99, 3.17]  $\delta_2^1 = \omega_2$  and hence by [Woo99, 3.16, 3.19] **acg** holds. The axiom  $\phi_{AC}$  is due to Woodin. By [Woo99, 5.9] **MM** implies  $\phi_{AC}$ . Note that by an observation of Larson **MM(c)** already suffices, see [Woo99, p.200].

We now state our results.

**Theorem 1.4** *If **BMM** holds and additionally  $\text{NS}_{\omega_1}$  is precipitous, then **acg** and  $\phi_{AC}$  hold.*

We will prove the above theorem using (a variant of)  $\mathbb{P}(\theta, \text{NS}_{\omega_1})$ . The technology developed to show  $\phi_{AC}$  can also be used to yield  $\psi_{AC}$ . We sketch such a construction only since Woodin has shown that **BMM** +  $\text{NS}_{\omega_1}$  is precipitous implies  $\psi_{AC}$  using more straightforward methods, see [Woo99, 10.95]. The following diagram illustrates the logical structure of the various statements:



Here **CBP** is the club bounding principle, i.e. the statement that every function  $f : \omega_1 \rightarrow \omega_1$  is bounded by a canonical function for some ordinal  $< \omega_2$  on a club. The implication from  $\psi_{AC}$  to **CBP** is due to Aspero and Welch, see [AW02]. All implications from **acg** are due to Woodin, see [Woo99, (proof of) 3.19].

The second part of this paper deals with the semiproperness of  $\mathbb{P}(\theta, \text{NS}_{\omega_1})$  for all regular  $\theta \geq \omega_2$  (or more general the semiproperness of any class of forcings that adds generic iterations like above). We will show:

**Theorem 1.5** *The following are equivalent:*

1. For arbitrarily large  $\theta \geq \omega_2$  there is a semiproper partial order  $\mathbb{P}$  that adds a generic iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle.$$

<sup>2</sup>We are working in models of **ZFC** so this will trivially hold. It is more interesting if working in models of **ZF** + **DC**.

such that  $H_\theta \subset M_{\omega_1}$  and all  $M_i$  are countable.

2. All stationary set preserving forcings are semiproper

## 2 The principle acg

In this section, we shall clean up [CS09] by showing the following.

**Lemma 2.1**  $\text{BMM} + \text{NS}_{\omega_1}$  is precipitous  $\implies$  acg.

*Proof.* Fix some club  $C$ . We show that admissible club guessing holds under BMM if the nonstationary ideal is precipitous. The forcing  $\mathbb{P}'(\omega_2, \text{NS}_{\omega_1})$  from [CS09] adds a countable generically iterable  $M_0$  generically iterating in  $\omega_1^V$  many steps to  $\langle (H_{\omega_2}^V)^\sharp, \in, \text{NS}_{\omega_1} \rangle$ , i.e. an iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle.$$

For brevity we write  $\pi_\alpha$  instead of  $\pi_{\alpha, \omega_1}$ . So there is some  $\alpha_0 < \omega_1$  such that  $C \cap \omega_1^{M_{\alpha_0}} \in M_{\alpha_0}$  and  $\pi_{\alpha_0}(C \cap \omega_1^{M_{\alpha_0}}) = C$ . We can assume w.l.o.g. by changing some indices that  $0 = \alpha_0$ . We now show that in the extension by  $\mathbb{P}'(\omega_2, \text{NS}_{\omega_1})$  there is a real  $y$  such that  $A_y \subset C$ . Let  $x$  be a real that codes  $M_0$  and let  $y$  code  $x^\sharp$ .

Writing  $C_\alpha = C \cap \omega_1^{M_\alpha}$  we have  $C_\alpha \in M_\alpha$  and  $\pi_\alpha(C_\alpha) = C$  for all  $\alpha < \omega_1$ . By elementarity,  $C_\alpha$  is unbounded in  $\omega_1^{M_\alpha}$ . So by the closedness of  $C$  we have  $\omega_1^{M_\alpha} \in C$ .

**Claim 1.** If  $\alpha$  is an  $x$ -indiscernible and

$$\langle \langle M'_i, \pi'_{i,j}, I'_i, \kappa'_i; i \leq j \leq \alpha \rangle, \langle G'_i; i < \alpha \rangle \rangle$$

is an arbitrary generic iteration of  $M = M'_0$  then  $\alpha = \omega_1^{M'_\alpha}$ .

*Proof of Claim 1.* First note that  $M$  is generically  $\omega_1 + 1$  iterable, by Theorem 18 of [CS09]. Fix an  $x$ -indiscernible  $\alpha$  and an iteration as above. Every  $x$ -indiscernible is inaccessible in  $L[x]$ , so for all  $\beta < \alpha$

$$L[x]^{\text{Col}(\omega, \beta)} \models \alpha \text{ is inaccessible.}$$

Let  $g \subset \text{Col}(\omega, \beta)$  be  $L[x]$ -generic. Assume w.l.o.g. that  $g$  is a real. Then, by [Woo99, 3.15] (compare Lemma 19 in [CS09]),  $M'_\beta \cap OR < \omega_1^{L[x, g]}$ . Hence  $\omega_1^{M'_\beta} < \alpha$ . This implies  $\omega_1^{M'_\alpha} \leq \alpha$ . So it follows easily that  $\omega_1^{M'_\alpha} = \alpha$ .  $\square$ (Claim 1)

If  $\alpha$  is  $x^\sharp$ -admissible, then  $\alpha$  is  $x$ -indiscernible. Hence by the above claim it follows that each  $y$ -admissible  $< \omega_1$  is in  $C$ . Hence  $A_{x^\sharp} \subset C$ . Since the existence of a real  $y$  such that  $A_y \subset C$  can be recast as a  $\Sigma_1$ -statement over  $H_{\omega_2}$  with  $C$  as a parameter, BMM implies that it is already true in  $V$ .  $\square$

## 3 Obtaining $\phi_{AC}$

We modify the forcing  $\mathbb{P}'(\omega_2, \text{NS}_{\omega_1})$  from [CS09] to show an arbitrary instance of  $\phi_{AC}$  in the generic extension. An application of BMM will then give us the desired result.

### 3.1 Hitting many regular cardinals

The following lemma states that for a generically iterable  $\langle M, I \rangle$  there is a generic iteration that realizes many regular cardinals.

**Lemma 3.1** (Hitting many regular cardinals lemma) *Let  $\langle M, I \rangle$  be a countable model of  $\text{ZFC}^-$  and let  $I$  be a precipitous ideal on  $\omega_1^M$ . Assume that  $\mathcal{P}(\mathcal{P}(\omega_1))$  exists in  $M$ . Let  $\theta, \alpha \in M$  be such that*

$$M \models (2^{2^{\omega_1}})^+ = \theta = \aleph_\alpha,$$

furthermore assume that

$$M \models (\aleph_{\alpha+\omega_1})^M \text{ exists.}$$

Let  $\theta' := (\aleph_{\alpha+\omega_1})^M$ . Then a genericity iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \theta' \rangle, \langle G_i; i < \theta' \rangle \rangle$$

of  $M_0 = M$  exists such that for all  $\beta < \omega_1^M$

$$\pi_{0, \aleph_{\alpha+\beta+1}^M}(\omega_1^M) = \aleph_{\alpha+\beta+1}^M.$$

*Proof.* Let  $g \subset \text{Col}(\omega, < \theta')$  be generic over  $M$ . Since  $M$  is countable in  $V$  the generic  $g$  can be chosen in  $V$ . Let  $\mathbb{P} := \mathcal{P}(\omega_1^M)^M \setminus I$ . For  $\beta < \omega_1^M$  we set

$$g_{\alpha+\beta+1} := g \cap \text{Col}(\omega, < \aleph_{\alpha+\beta+1}^M).$$

Clearly all the  $g_i$  defined in this fashion are generic over  $M$ . Recursively we construct a generic iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \theta' \rangle, \langle G_i; i < \theta' \rangle \rangle$$

such that for  $\beta < \omega_1^M$  the sequence  $\langle G_i; i < \aleph_{\alpha+\beta+1}^M \rangle$  is in  $M[g_{\alpha+\beta+1}]$ . We inductively maintain the following:

- For  $\beta < \omega_1^M$  and  $i < \aleph_{\alpha+\beta+1}^M$  the set

$$D_i = \{d \in M_i; d \subset \pi_{0,i}(\mathbb{P}) \wedge M_i \models d \text{ is a dense in } \pi_{0,i}(\mathbb{P})\}$$

is countable in  $M[g_{\alpha+\beta+1}]$ .

Set  $M_0 = M$ ,  $I_0 = I$  and  $\kappa_0 = \omega_1^M$ . Assume we are at stage  $i < \theta'$  of the construction. Let  $\beta < \omega_1^M$  be least such that  $i < \aleph_{\alpha+\beta+1}^M$ . Inductively we have that  $D_i$  is countable in  $M[g_{\alpha+\beta+1}]$ . Choose a  $D_i$  generic  $G_i$  in  $M[g_{\alpha+\beta+1}]$ . At limit stages form direct limits.

Let us check our inductive hypotheses in the successor case, the limit case being an easy consequence of the fact that the sequence  $\langle G_i; i < \aleph_{\alpha+\beta+1}^M \rangle$  is in  $M[g_{\alpha+\beta+1}]$ . For the successor case note that an appropriate hull of

$$\pi_{0,i+1}{}^{\text{cc}}(H_\theta)^{M_0} \cup \{\kappa_j; j < i+1\}$$

is  $(H_{\theta_{i+1}})^{M_{i+1}}$  where  $\theta_{i+1} = \pi_{0,i+1}(\theta)$ . This hull can be calculated in  $M[g_{\alpha+\beta+1}]$ . Hence  $D_{i+1} \subset (H_{\theta_{i+1}})^{M_{i+1}}$  is also countable in  $M[g_{\alpha+\beta+1}]$ . It is trivial to maintain that the sequence  $\langle G_j; j < i+1 \rangle$  is in  $M[g_{\alpha+\beta+1}]$ .

Now we need that  $\aleph_{\alpha+\beta+1}^M$  is regular in  $M$ . Hence

$$\omega_1^{M[g_{\alpha+\beta+1}]} = \aleph_{\alpha+\beta+1}^M.$$

So an easy calculation shows that for all  $\beta < \omega_1^M$

$$\pi_{0, \aleph_{\alpha+\beta+1}^M}(\omega_1^M) = \aleph_{\alpha+\beta+1}^M.$$

□

Clearly the previous lemma can be generalized further. Since we only need the case above, we refrained to state it in a more general fashion. Note that we have a lot of freedom when choosing the generics of the iteration; the only true restriction is that they come from small generic extensions. We will make use of this later. We define a set of ordinals relative to a generic iteration. This set will come in handy in the proof of the main result of this section.

**Definition 3.2** Let  $\langle M, I \rangle$  be a model of  $\text{ZFC}^- + \text{“}\omega_1 \text{ exists,“}$  such that  $M \models I$  is precipitous. Let  $\theta$  be a cardinal in  $M$ . Let

$$\mathcal{J} := \langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \rho \rangle, \langle G_i; i < \rho \rangle \rangle$$

be a generic iteration of  $\langle M_0, I_0 \rangle = \langle M, I \rangle$ . We inductively define the *important ordinals of  $\mathcal{J}$  relative to  $\theta$* .

1. 0 is an important ordinal.
2. If  $\alpha$  is an important ordinal then the least ordinal  $\gamma$  such that  $\pi_{0,\alpha}(\theta) \leq \gamma = \kappa_\gamma$  is the next important ordinal.
3. Limits of important ordinals are important.

**Remark 3.3** Let  $\langle M, I \rangle$  be countable and as in the previous definition and let  $\mathcal{J}$  as in the previous definition and  $\rho = \omega_1$ . Then clearly the set of important ordinals of  $\mathcal{J}$  relative to  $\theta$  is a club in  $\omega_1$ . Also, if  $\alpha$  is important, then  $\kappa_\alpha = \alpha$ .

### 3.2 Forcing $\phi_{AC}$

We will show the following theorem:

**Theorem 3.4** Let  $\aleph_\alpha = 2^{2^{\omega_1}}$ . Let  $\theta := \aleph_{\alpha+\omega_1}$ . Let  $\text{NS}_{\omega_1}$  be precipitous and suppose  $H_\theta^\sharp$  exists. Let  $F : \omega_1 \rightarrow \theta$  defined by

$$F(\beta) = \aleph_{\alpha+\beta+1}.$$

Let  $\vec{S} = \langle S_k; k \in \omega \rangle$ ,  $\vec{T} = \langle T_k; k \in \omega \rangle$  be sequences of pairwise disjoint subsets of  $\omega_1$ , such that all  $S_k$  are stationary and  $\omega_1 = \bigcup \{T_k; k \in \omega\}$ . There exists a forcing construction  $\mathbb{P} = \mathbb{P}'(\theta, \text{NS}_{\omega_1}, \vec{S}, \vec{T})$  that preserves stationary subsets such that if  $G$  is  $\mathbb{P}$ -generic over  $V$ , then in  $V[G]$  there is generic iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

such that if  $i < \omega_1$ , then  $M_i$  is countable and  $M_{\omega_1} = \langle H_\theta^\sharp; \in, \text{NS}_{\omega_1} \rangle$ . In particular,  $M_0$  is generically  $\omega_1$ -iterable. Additionally the following holds in  $V[G]$  for all  $k \in \omega$ :

$$F \text{``} T_k \subset \tilde{S}_k.$$

We use a similar setup as [CS09], i.e. we assume:

$$\theta = 2^{<\theta} < 2^\theta < \rho = 2^{<\rho},$$

for some cardinal  $\rho$ . For reasons of convenience we like to think of  $\aleph_\alpha = 2^{2^{\omega_1}}$  as  $\aleph_3$ . This eases notation considerably. Note that we can force  $\aleph_3 = 2^{2^{\omega_1}}$  with stationary set preserving forcing. If  $2^{\omega_1} = \aleph_2$ , then the precipitousness of  $\text{NS}_{\omega_1}$  is preserved by forcing with  $\text{Col}(\omega_3, 2^{2^{\omega_1}})$ , since no new subsets of  $2^{\omega_1}$  are added, see [Jec03, 22.19]. Nevertheless the reader will gladly verify that all of the following arguments

go through for an arbitrary  $\aleph_\alpha$  instead of  $\aleph_3$ . If  $\aleph_\alpha = \aleph_3$ , then clearly  $\theta = \aleph_{\omega_1}$ . At this point a remark is in order. In [CS09]  $\theta$  is supposed to be regular. Nevertheless it is straightforward to check that if one can add generic iterations like in in [CS09] with last model  $H_\eta$  for arbitrarily large regular  $\eta$  you can also add generic iterations with last model  $H_\theta$ . We can hence work with a singular  $\theta$  and use the theory of [CS09].

Fix a well-order  $<$  of  $H_\rho$  as in [CS09]. We now fix  $\vec{S} = \langle S_k ; k \in \omega \rangle$ ,  $\vec{T} = \langle T_k ; k \in \omega \rangle$  sequences of pairwise disjoint subsets of  $\omega_1$ , such that all  $S_k$  are stationary and  $\omega_1 = \bigcup \{T_k ; k \in \omega\}$ . We use

$$\mathcal{H} = \langle H_\rho ; \in, H_\theta^\sharp, \text{NS}_{\omega_1}, < \rangle$$

and

$$\mathcal{M} = \langle H_\theta^\sharp ; \in, \text{NS}_{\omega_1}, < \rangle$$

since we are defining a variant of  $\mathbb{P}'(\theta, \text{NS}_{\omega_1})$ . We will now define our modified forcing construction  $\mathbb{P}'(\theta, \text{NS}_{\omega_1}, \vec{S}, \vec{T})$ .

**Definition 3.5** Conditions  $p$  in  $\mathbb{P}'(\theta, \text{NS}_{\omega_1}, \vec{S}, \vec{T})$  are triples

$$p = \langle \langle \kappa_i^p ; i \in \text{dom}(p) \rangle, \langle \pi_i^p ; i \in \text{dom}(p) \rangle, \langle \tau_i^p ; i \in \text{dom}_-(p) \rangle \rangle$$

such that the following conditions hold:

1. Both  $\text{dom}(p)$  and  $\text{dom}_-(p)$  are finite, and  $\text{dom}_-(p) \subset \text{dom}(p) \subset \omega_1$ .
2.  $\langle \kappa_i^p ; i \in \text{dom}(p) \rangle$  is a sequence of countable ordinals.
3.  $\langle \pi_i^p ; i \in \text{dom}(p) \rangle$  is a sequence of finite partial maps from  $\omega_1$  to  $H_\theta^\sharp \cap \text{OR}$ .
4.  $\langle \tau_i^p ; i \in \text{dom}_-(p) \rangle$  is a sequence of complete  $\mathcal{H}$ -types over  $H_\theta$ , i.e., for each  $i \in \text{dom}_-(p)$  there is some  $x \in H_\rho$  such that, having  $\varphi$  range over  $\mathcal{H}$ -formulae with free variables  $u, \vec{v}$ ,

$$\tau_i^p = \{ \langle \ulcorner \varphi \urcorner, \vec{z} \rangle ; \vec{z} \in H_\theta \wedge \mathcal{H} \models \varphi[x, \vec{z}] \}.$$

5. If  $i, j \in \text{dom}_-(p)$ , where  $i < j$ , then there is some  $n < \omega$  and some  $\vec{u} \in \text{ran}(\pi_j^p)$  such that

$$\tau_i^p = \{ (m, \vec{z}) ; (n, \vec{u} \frown m \frown \vec{z}) \in \tau_j^p \}.$$

6. In  $V^{\text{Col}(\omega, \theta)}$ , there is a model which certifies  $p$  with respect to  $\mathcal{M}$ , i.e. a model  $\mathfrak{A}$  such that  $H_\theta^\sharp \in \text{wfp}(\mathfrak{A})$ ,  $\mathfrak{A} \models \text{ZFC}^-$ , for all stationary  $S$ ,  $\mathfrak{A} \models$  “ $S$  is stationary”, and inside  $\mathfrak{A}$  there is a generic iteration

$$\mathcal{J}^\mathfrak{A} := \langle \langle M_i^\mathfrak{A}, \pi_{i,j}^\mathfrak{A}, I_i^\mathfrak{A}, \kappa_i^\mathfrak{A} ; i \leq j \leq \omega_1 \rangle, \langle G_i^\mathfrak{A} ; i < \omega_1 \rangle \rangle$$

such that

- (a) if  $i < \omega_1$ , then  $M_i^\mathfrak{A}$  is countable,
- (b) if  $i < \omega_1$  and if  $\xi < \theta$  is definable over  $\mathcal{M}$  from parameters in  $\text{ran}(\pi_{i, \omega_1}^\mathfrak{A})$ , then  $\xi \in \text{ran}(\pi_{i, \omega_1}^\mathfrak{A})$ ,
- (c)  $M_{\omega_1} = \langle H_\theta^\sharp ; \in, \text{NS}_{\omega_1} \rangle$ ,
- (d) if  $i \in \text{dom}(p)$ , then  $\kappa_i^p = \kappa_i^\mathfrak{A}$  and  $\pi_i^p \subset \pi_{i, \omega_1}^\mathfrak{A}$ ,
- (e) if  $i \in \text{dom}_-(p)$ , then for all  $n < \omega$  and for all  $\vec{z} \in \text{ran}(\pi_{i, \omega_1}^\mathfrak{A})$ ,

$$\exists y \in H_\theta (n, y \frown \vec{z}) \in \tau_i^p \implies \exists y \in \text{ran}(\pi_{i, \omega_1}^\mathfrak{A}) (n, y \frown \vec{z}) \in \tau_i^p.$$

- (f) Let  $D^{\mathfrak{A}}$  be the set of important ordinals of  $\mathcal{J}^{\mathfrak{A}}$  relative to  $(\pi_{0,\omega_1}^{\mathfrak{A}})^{-1}(\theta)$ .  
 If  $\gamma \in D^{\mathfrak{A}}$  then for all  $\beta < \gamma = \kappa_\gamma^{\mathfrak{A}}$  and all  $k \in \omega$ .

$$\aleph_{3+\beta+1}^{M_\gamma^{\mathfrak{A}}} \in S_k \iff \beta \in T_k.$$

If  $p, q \in \mathbb{P}'(\theta, \mathbf{NS}_{\omega_1}, \vec{S}, \vec{T})$ , then we write  $p \leq q$  iff  $\text{dom}(q) \subset \text{dom}(p)$ ,  $\text{dom}_-(q) \subset \text{dom}_-(p)$ , for all  $i \in \text{dom}(q)$ ,  $\kappa_i^p = \kappa_i^q$  and  $\pi_i^q \subset \pi_i^p$ , and for all  $i \in \text{dom}_-(q)$ ,  $\tau_i^q = \tau_i^p$ .

We now show theorem 3.4. First we show that  $\mathbb{P} := \mathbb{P}'(\theta, \mathbf{NS}_{\omega_1}, \vec{S}, \vec{T}) \neq \emptyset$ , i.e. the analog of Lemma 5 in [CS09]. Then we proceed as in [CS09] but we will skip all lemmata and theorems that are literally the same and have literally the same proof.

**Lemma 3.6**  $\mathbb{P} \neq \emptyset$ .

*Proof.* We need to verify, that in  $V^{\text{Col}(\omega, \theta)}$  there is a model which certifies the trivial condition with respect to  $\mathcal{M}$ . Let  $g$  be  $\text{Col}(\omega, < \rho)$ -generic over  $V$ . We work in  $V[g]$  until further notice. So  $\langle V; \in, \mathbf{NS}_{\omega_1} \rangle$  is  $\rho + 1$  iterable, by Lemma 2 of [CS09]. Hence  $\langle H_\theta^\#; \in, \mathbf{NS}_{\omega_1} \rangle$  is also  $\rho + 1$  iterable. We prepare a book-keeping device: pick a bijection  $g : [\rho]^{<\rho} \rightarrow \rho$  and a family  $\langle S_\nu, \nu < \rho \rangle$  of pairwise disjoint stationary subsets of  $\rho$ . Now define  $f : \rho \rightarrow [\rho]^{<\rho}$  by

$$f(i) = s \iff i \in S_{g(s)}.$$

Note that each  $s$  is enumerated stationarily often. We recursively construct a generic iteration

$$\mathcal{J} := \langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \rho \rangle, \langle G_i; i < \rho \rangle \rangle$$

of  $M_0 = \langle V; \in, \mathbf{NS}_{\omega_1} \rangle$  together with a set of local generics  $g_i$ . Later the restriction of this iteration to  $\langle H_\theta^\#; \in, \mathbf{NS}_{\omega_1} \rangle$  will be of interest. For each important ordinal of the iteration a local generic  $g_i$  will be picked. Suppose we have already constructed  $\mathcal{J}$  to some  $i < \rho$ . Note that we can calculate the important ordinals of  $\mathcal{J}$  relative to  $\theta$  while we construct  $\mathcal{J}$ . The following three clauses define the iteration.

1. If  $i$  is an important ordinal of  $\mathcal{J}$  relative to  $\theta$ , then pick some  $g_i \subset \text{Col}(\omega, < \pi_{0,i}(\theta))$  in  $V[g]$  that is generic over  $M_i$ . Then pick  $G_i$  in  $M_i[g_i]$  such that if for a (unique)  $j$  the set  $\pi_{j,i}(f(i))$  is stationary in  $M_i$  then  $\pi_{j,i}(f(i)) \in G_i$ . Note that  $j$  is unique because  $f(i)$  can only be stationary in  $M_j$  if  $\sup f(i) = \omega_1^{M_j}$ .
2. If  $i$  is not important and  $\gamma$  is the largest important ordinal below  $i$ , then we already have chosen some  $g_\gamma \subset \text{Col}(\omega, < \pi_{0,\gamma}(\theta))$  in  $V[g]$  that is generic over  $M_\gamma$ . In the case that  $i = \omega_{3+\beta+1}^{M_\gamma}$  for some  $\beta < \kappa_\gamma = \gamma$  we pick some  $G_i$  in  $M_\gamma[g_\gamma \cap \text{Col}(\omega, < \omega_{3+\beta+2}^{M_\gamma})]$  such that

$$\beta \in \pi_{0,\gamma}(T_k) \iff \pi_{0,i}(S_k) \in G_i.$$

Note that since  $\vec{T}$  is a partition of  $\omega_1$ , there is a unique  $k$  such that  $\beta \in \pi_{0,i}(T_k)$ .

3. If the first and second clause do not hold and  $\gamma$  is the largest important ordinal below  $i$ , then we already have chosen some  $g_\gamma \subset \text{Col}(\omega, < \pi_{0,\gamma}(\theta))$  in  $V[g]$  that is generic over  $M_\gamma$ . In the case that  $i$  is not a successor cardinal  $< \pi_{0,\gamma}(\theta)$  in  $M_\gamma$  there is a least  $\beta < \kappa_\gamma$  such that  $i < \omega_{3+\beta+1}^{M_\gamma}$ . We pick some arbitrary  $G_i$  in  $M_\gamma[g_\gamma \cap \text{Col}(\omega, < \omega_{3+\beta+1}^{M_\gamma})]$ . Else we pick a completely arbitrary generic.



Fix some important  $\gamma > 0$ . So  $\mathcal{J}$  restricted to  $[\gamma, \pi_{0,\gamma}(\theta)[$  is an iteration like in the Hitting many regular cardinals lemma 3.1. Hence we know that the iteration is well defined and additionally we have for  $\beta < \kappa_\gamma = \gamma$  and  $i := \aleph_{3+\beta+1}^{M_\gamma}$

$$i = \pi_{\gamma,i}(\kappa_\gamma) = \kappa_i.$$

By the second clause of the iteration we hence have for  $i$  as above and  $k \in \omega$ :

$$\beta \in \pi_{0,\gamma}(T_k) \iff \pi_{0,i}(S_k) \in G_i \iff \kappa_i \in \pi_{0,i+1}(S_k) \iff i \in \pi_{0,\rho}(S_k).$$

Let  $D$  denote the club of important ordinals and let  $S \in \text{NS}_{\omega_1}^{M_\rho}$ . Let  $j < \rho$  and  $s$  be such that  $\pi_{j,\rho}(s) = S$ . If  $i \in D \setminus j$  and  $f(i) = s$ , then  $\pi_{j,i}(s) \in G_i$ . This shows that

$$D \cap S_{g(s)} \setminus j \subset \{i < \rho; \kappa_i \in S\},$$

so that in fact  $S$  is stationary in  $V[g]$ .

Hence in  $M_\rho^{\text{Col}(\omega, \pi_{0,\rho}(\theta))}$  there is a model that certifies the empty condition with respect to  $\pi_{0,\rho}(\langle H_\theta^\sharp; \in, \text{NS}_{\omega_1} \rangle)$ . Now we can literally complete our proof by following the last paragraph of the proof Lemma 5 in [CS09].  $\square$

We can now literally adopt lemmata 6 through 15 of [CS09]. So we have, using the notation of [CS09]:

**Lemma 3.7** *Let  $G \subset \mathbb{P}$  is  $V$ -generic. Let  $\kappa_i = \kappa_i^p$  for some  $p \in G$ . Then in  $V[G]$*

$$H_\theta^\sharp \cap \text{OR} = \cup \{\text{ran}(\pi_i); i < \omega_1\}$$

and

$$\mathcal{J}_G := \langle \langle M_i, \tilde{\pi}_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1^V \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

is a generic iteration of  $M_0$  such that if  $i < \omega_1$ , then  $M_i$  is countable, and  $M_{\omega_1} = \langle H_\theta^\sharp; \in, I \rangle$ .

Let  $D_G$  denote the important ordinals of  $\mathcal{J}_G$ . We can assume without loss of generality that there are  $\vec{s}, \vec{t} \in M_0$  such that  $\tilde{\pi}_{0,\omega_1}(\langle \vec{s}, \vec{t} \rangle) = \langle \vec{S}, \vec{T} \rangle$ .

**Lemma 3.8**  *$D_G$  is club and for all  $\gamma \in D_G$  the following holds: if  $\beta < \kappa_\gamma$  then for all  $k \in \omega$*

$$\beta \in \pi_{0,\gamma}(t_k) \iff \aleph_{3+\beta+1}^{M_\gamma} \in \pi_{0,\omega_1}(s_k),$$

which by the choice of  $\vec{s}$  and  $\vec{t}$  means

$$\beta \in T_k \iff \aleph_{3+\beta+1}^{M_\gamma} \in S_k$$

*Proof.* That  $D_G$  is club is obvious.

**Claim 1.**  $p \Vdash \check{\gamma} \in D_{\dot{G}}$  if and only if for all  $\mathfrak{A}$  which certify  $p$ ,  $\gamma \in D^{\mathfrak{A}}$ .

*Proof of Claim 1.* Fix  $p$  such that  $p \Vdash \check{\gamma} \in D_{\dot{G}}$  and some structure  $\mathfrak{A}$  which certifies  $p$ . Towards a contradiction suppose  $\gamma \notin D^{\mathfrak{A}}$ . Then there is some  $\gamma' < \gamma$ ,  $\gamma' \in D^{\mathfrak{A}}$  with

$$(\pi_{\gamma',\omega_1}^{\mathfrak{A}})^{-1}(\theta) > \gamma.$$

We can extend  $p$  to  $p'$  also certified by  $\mathfrak{A}$  such that  $\text{dom}(p')$  contains all the relevant points. Then

$$p' \Vdash \check{\gamma} \notin D_{\dot{G}}.$$

Contradiction! The other direction is easy.

□(Claim 1)

Now if  $\beta \in \pi_{0,\gamma}(t_k)$  and  $\gamma \in D_G$  there is some  $p \in G$  with  $p \Vdash \check{\gamma} \in D_{\dot{G}}$  and  $\beta \in (\pi_\gamma^p)^{-1} \circ \pi_0^p(t_k)$  (Note the following subtlety:  $\pi_0^p$  is only defined on the ordinals, but using the well ordering  $<$  on  $H_\theta^\#$  we can assume that  $\text{dom}(\pi_0^p)$  contains  $t_k$ ). Let  $p' \leq p$  be arbitrary and let  $\mathfrak{A}$  certify  $p'$ . Then  $\aleph_{3+\beta+1}^{M_\gamma^\mathfrak{A}} \in S_k$  by the above claim and the fact that  $\mathfrak{A}$  certifies  $p'$ . So we may extend  $p'$  to  $p''$  making sure

$$p'' \Vdash \aleph_{3+\beta+1}^{M_\gamma} \in \tilde{\pi}_{0,\omega_1}(s_k).$$

Hence the set of  $p''$  forcing the desired result is dense below  $p$ . The other direction is similar. □

We can now literally adopt lemmata 16 and 17 of [CS09] and their proofs; i.e. it is clear that  $\mathbb{P}'(\theta, \mathbf{NS}_{\omega_1}, S, T)$  is stationary set preserving.

To finish the proof of 3.4 we have to show that in  $V[G]$  for all  $k \in \omega$

$$F \Vdash T_k \subset \tilde{S}_k.$$

For this fix  $k \in \omega$  and some  $\beta \in T_k$ . By 3.8 we have for all  $\gamma \in D_G \setminus (\beta + 1)$

$$\beta \in T_k \iff \aleph_{3+\beta+1}^{M_\gamma} \in S_k.$$

**Lemma 3.9** *The function  $f : D_G \setminus (\beta + 1) \rightarrow \omega_1$*

$$\gamma \mapsto \aleph_{3+\beta+1}^{M_\gamma}$$

*is a canonical function for  $\aleph_{3+\beta+1}^V < \omega_2^{V[G]}$  in  $V[G]$ .*

*Proof.* Let

$$\mathcal{J}_G := \langle \langle M_i, \tilde{\pi}_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1^V \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

denote the iteration that is added by  $G$ . Set  $\eta := \aleph_{3+\beta+1}^V$ . Fix some bijection  $g : \omega_1 \rightarrow \eta$  in  $V[G]$ . Let  $\langle X_i; i \in \omega_1 \rangle$  be a continuous elementary chain of countable submodels of  $H_{\omega_2}^{V[G]}$  such that  $g, H_\theta^V \in X_0$ . So clearly  $H_\theta^V \subset \cup \{X_i; i \in \omega\}$ . So for all  $i \in \omega_1$  we have

$$X_i \cap \eta = g \Vdash (X_i \cap \omega_1).$$

Clearly  $\langle X_i \cap H_\theta^V; i \in \omega_1 \rangle$  is club in  $[H_\theta^V]^\omega$ . Since the set  $\{\text{ran}(\tilde{\pi}_{i,\omega_1}) \cap H_\theta; i \in \omega_1\}$  is also a club in  $[H_\theta^V]^\omega$  there is a club  $C \subset \omega_1$  such that for all  $i \in C$

$$X_i \cap \eta = \text{ran}(\tilde{\pi}_{i,\omega_1}) \cap \eta.$$

So for all  $i \in C$  we have

$$i = \text{ran}(\tilde{\pi}_{i,\omega_1}) \cap \omega_1 = X_i \cap \omega_1$$

and thus

$$\text{otp}(g \Vdash i) = \text{otp}(\text{ran}(\tilde{\pi}_{i,\omega_1}) \cap \eta) = \aleph_{3+\beta+1}^{M_i} = f(i).$$

Hence  $f$  is a canonical function. □

So the club  $D_G \setminus (\beta + 1)$  and  $f$  from the previous lemma witness that in  $V[G]$

$$\mathbf{1}_\mathbb{B} \Vdash \aleph_{3+\beta+1}^V \in j(S_i),$$

where  $\mathbb{B}$  is  $(\mathcal{P}(\omega_1) \setminus \text{NS}_{\omega_1})^{V[G]}$  and  $j$  is a name for the generic embedding added by forcing with  $\mathbb{B}$ . Hence  $\aleph_{3+\beta+1}^V \in \tilde{S}_i$ . This finishes the proof of 3.4.

Observe that the single instance of  $\phi_{AC}$  that holds in  $V^{\mathbb{P}'(\theta, \text{NS}_{\omega_1}, \vec{S}, \vec{T})}$  is a  $\Sigma_1$  statement in  $H_{\omega_2}$  in the parameters  $\vec{S}$  and  $\vec{T}$ . Since  $\mathbb{P}'(\theta, \text{NS}_{\omega_1}, \vec{S}, \vec{T})$  preserves stationary subsets an application of BMM yields the following corollary.

**Corollary 3.10** *If  $\text{NS}_{\omega_1}$  is precipitous+BMM then  $\phi_{AC}$ .*

## 4 Obtaining $\psi_{AC}$

**Definition 4.1** (Woodin)  $\psi_{AC}$ : Let  $S \subset \omega_1$  and  $T \subset \omega_1$  be stationary, costationary sets. Then there exists a canonical function  $f$  for some  $\eta < \omega_2$  such that for some club  $C \subset \omega_1$

$$\{\alpha < \omega_1; f(\alpha) \in T\} \cap C = S \cap C.$$

Note the following reformulation of the above definition in terms of generic ultrapowers: let  $j$  be a name for the embedding induced by some generic  $G \subset \mathcal{P}(\omega_1) \setminus \text{NS}_{\omega_1}$ , with  $S, T$  as above we have

$$\mathbf{1}_{\mathcal{P}(\omega_1) \setminus \text{NS}_{\omega_1}} \Vdash \check{S} \in \dot{G} \iff \eta \in j(T).$$

Woodin has shown:

**Theorem 4.2** ([Woo99, 10.95]) *If BMM +  $\text{NS}_{\omega_1}$  is precipitous then  $\psi_{AC}$ .*

With the technology from the previous section on  $\phi_{AC}$  it is possible to give a different proof of 4.2. Since this is very similar to the section on  $\phi_{AC}$ , we shall only state the required results. The proofs are very similar to the  $\phi_{AC}$  case.

**Lemma 4.3** (Hitting regular cardinals lemma) *Let  $\langle M, I \rangle$  be a countable model of ZFC\* and let  $I$  be a precipitous ideal on  $\omega_1^M$ . Assume that  $\mathcal{P}(\mathcal{P}(\omega_1))$  exists in  $M$ . Let  $\theta \in M$  be such that*

$$M \models \text{Card}(\mathcal{P}(\mathcal{P}(\omega_1)))^+ = \theta,$$

and let  $\theta' \geq \theta$  such that  $\theta'$  is a regular cardinal in  $M$ . Then a genericity iteration

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \theta' \rangle, \langle G_i; i < \theta' \rangle \rangle$$

of  $M_0 = M$  exists in  $V$  such that  $\pi_{0,\theta'}(\omega_1^M) = \theta'$ .

We again modify the forcing  $\mathbb{P}'(\omega_2, \text{NS}_{\omega_1})$  to show a weak form of  $\psi_{AC}$  in the generic extension. An application of BMM will then give us the desired result.

**Theorem 4.4** *Let  $\text{NS}_{\omega_1}$  be precipitous and suppose  $H_\theta^\sharp$  exists, where  $\theta = 2^{2^{\aleph_1}}$ . For all  $S, T$  stationary and costationary there exists a forcing construction  $\mathbb{P} = \mathbb{P}'(\theta, \text{NS}_{\omega_1}, S, T)$  that preserves stationary subsets, such that if  $G$  is  $\mathbb{P}$ -generic over  $V$ , then in  $V[G]$  there is generic iteration*

$$\langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

such that if  $i < \omega_1$ , then  $M_i$  is countable and  $M_{\omega_1} = \langle H_\theta^\sharp; \in, \text{NS}_{\omega_1} \rangle$ . In particular,  $M_0$  is generically  $\omega_1$ -iterable. Additionally the following holds in  $V[G]$ : there is a club  $C \subset \omega_1$ , such that for all  $\alpha \in C$

$$\omega_1^{M_\alpha} \in S \iff \theta_\alpha \in T,$$

where  $\theta_\alpha = \pi_{\alpha, \omega_1}^{-1}(\theta)$ .

We will now define our modified forcing construction  $\mathbb{P} := \mathbb{P}'(\theta, \text{NS}_{\omega_1}, S, T)$ .

**Definition 4.5** Conditions  $p$  in  $\mathbb{P}'(\theta, \text{NS}_{\omega_1}, S, T)$  are triples

$$p = \langle \langle \kappa_i^p; i \in \text{dom}(p) \rangle, \langle \pi_i^p; i \in \text{dom}(p) \rangle, \langle \tau_i^p; i \in \text{dom}_-(p) \rangle \rangle$$

such that the following conditions hold:

Conditions i.,ii.,iii.,iv.,v. as in definition 3.5 hold. We replace condition vi. as follows:

vi. In  $V^{\text{Col}(\omega, \theta)}$ , there is a *model which certifies  $p$  with respect to  $\mathcal{M}$* , i.e. a model  $\mathfrak{A}$  such that  $H_\theta^\sharp \in \text{wfp}(\mathfrak{A})$ ,  $\mathfrak{A} \models \text{ZFC}^-$ , for all stationary  $S$ ,  $\mathfrak{A} \models$  “ $S$  is stationary”, and inside  $\mathfrak{A}$  there is a generic iteration

$$\mathcal{J}^\mathfrak{A} := \langle \langle M_i^\mathfrak{A}, \pi_{i,j}^\mathfrak{A}, I_i^\mathfrak{A}, \kappa_i^\mathfrak{A}; i \leq j \leq \omega_1 \rangle, \langle G_i^\mathfrak{A}; i < \omega_1 \rangle \rangle$$

such that conditions (a),(b),(c),(d) and (e) as in definition 3.5 hold. We replace (f).

(f) Let  $D^\mathfrak{A}$  be the club of *limits* of important ordinals of  $\mathcal{J}^\mathfrak{A}$  relative to  $\pi_{0, \omega_1}^{\mathfrak{A}-1}(\theta)$ . Let  $\alpha \in D^\mathfrak{A}$ . Let  $\beta$  be the next important ordinal above  $\alpha$ . Then

$$\omega_1^{M_\alpha^\mathfrak{A}} \in S \iff \pi_{\alpha, \omega_1}^{\mathfrak{A}-1}(\theta) = \omega_1^{M_\beta^\mathfrak{A}} \in T.$$

If  $p, q$  are conditions, then we write  $p \leq q$  iff  $p \leq_{\mathbb{P}'(\theta, \text{NS}_{\omega_1})} q$ .

Applying the Hitting regular cardinals lemma one can show that certifying structures exist. Hence one has:

**Lemma 4.6**  $\mathbb{P} \neq \emptyset$ .

We can now literally adopt lemmata 6 through 15 of [CS09]. So we have, using the definitions for  $\pi_i, M_i, I_i, \kappa_i, G_i, \tilde{\pi}_{i,j}$ , of [CS09]:

**Lemma 4.7** *Let  $G \subset \mathbb{P}$  is  $V$ -generic. Let  $\kappa_i = \kappa_i^p$  for some  $p \in G$ . Then*

$$H_\theta^\sharp \cap \text{OR} = \cup \{ \text{ran}(\pi_i); i < \omega_1 \}$$

and

$$\mathcal{J}_G := \langle \langle M_i, \tilde{\pi}_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1^V \rangle, \langle G_i; i < \omega_1 \rangle \rangle$$

is a generic iteration of  $M_0$  such that if  $i < \omega_1$ , then  $M_i$  is countable, and  $M_{\omega_1} = \langle H_\theta^\sharp; \in, I \rangle$ .

We set

$$\theta_i := \tilde{\pi}_{i, \omega_1}^{-1}(\theta),$$

and we let  $D_G$  denote the club of limits of important ordinal of  $\mathcal{J}$  relative to  $\theta_0$ . A density argument shows:

**Lemma 4.8**  $D_G$  is club and for all  $i \in D_G$

$$\omega_1^{M_i} \in S \iff \theta_i \in T.$$

Since the sequence  $\langle \theta_i; i \in D_G \rangle$  is a canonical function for  $\theta$  in the forcing extension, we have

$$\mathbf{1}_{\mathbb{P}(\omega_1) \setminus \text{NS}_{\omega_1}} \Vdash \check{S} \in \check{G} \iff \theta \in \check{j}(T).$$

We can now literally adopt lemmata 16 and 17 of [CS09] and their proofs; i.e. it is clear that  $\mathbb{P}'(\theta, \text{NS}_{\omega_1}, S, T)$  is stationary set preserving. Hence theorem 4.4 follows.

## 5 The semiproperness of $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$

In [CS09] it was shown that  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  preserves stationary subsets of  $\omega_1$  provided that  $\text{NS}_{\omega_1}$  is precipitous. Since it is consistent relative to large cardinals that all stationary set preserving forcings are semiproper, the forcing  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  can clearly be semiproper. We show that the semiproperness of the forcings  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  implies a generalization of Chang's Conjecture which in turn implies the semiproperness of all stationary set preserving forcings.

Recall the definition of semiproperness.

**Definition 5.1** A notion of forcing  $\mathbb{P}$  is semiproper if for every sufficiently large  $\lambda$ , every well-ordering  $<$  of  $H_\lambda$  and every countable elementary submodel  $X \prec \langle H_\lambda; \in, < \rangle$  the following holds:

$$\forall p \in X \cap \mathbb{P} \exists q \leq p : q \text{ is } (X, \mathbb{P})\text{-semigeneric,}$$

where  $q$  is  $(X, \mathbb{P})$ -semigeneric if for every name  $\dot{\alpha} \in X$  for a countable ordinal

$$\exists \beta \in X : q \Vdash \dot{\alpha} = \check{\beta}.$$

**Definition 5.2** ([She98, XIII. 1.5])

- Let  $x, y$  be countable. We write  $x \sqsubset y$  if  $x \cap \omega_1 = y \cap \omega_1$  and  $x \subset y$ .
- A set  $S \subset [W]^\omega$  is *semistationary* in  $[W]^\omega$  if  $\{y \in [W]^\omega; \exists x \in S : x \sqsubset y\}$  is stationary in  $[W]^\omega$ .
- Let  $\lambda \geq \omega_2$ . We denote by  $\text{SSR}([\lambda]^\omega)$  the following principle: For every  $S$  semistationary in  $[\lambda]^\omega$  there is  $W \subset \lambda$ ,  $\text{Card}(W) = \omega_1 \subset W$  and  $S \cap [W]^\omega$  is semistationary in  $[W]^\omega$ .
- If  $\text{SSR}([\lambda]^\omega)$  holds for all cardinals  $\lambda \geq \omega_2$  then we will say that *Semistationary Reflection* (SSR) holds.

Note that [She98] has a more general notation for the above reflection principles. In [She98] the principle  $\text{SSR}([\lambda]^\omega)$  is called  $\text{Rss}(\aleph_2, \lambda)$  and SSR is called  $\text{Rss}(\aleph_2)$ .

**Lemma 5.3** ([She98, XIII.1.7(3)]) *Semistationary Reflection implies that all stationary set preserving forcings are semiproper.*

**Definition 5.4** ([FMS88])  $(\dagger)$  is an abbreviation for: every stationary set preserving forcing is semiproper.

Foreman, Magidor and Shelah have shown:

**Lemma 5.5** ([FMS88, Theorem 26]) *If  $(\dagger)$  then  $\text{NS}_{\omega_1}$  is precipitous.*

We will consider a generalization of Chang's Conjecture that we call  $\text{CC}^{**}$ .

**Definition 5.6** Let  $\lambda \geq \omega_2$ .  $\text{CC}^*(\lambda)$  is the following axiom: There are arbitrarily large regular cardinals  $\theta > \lambda$  such that for all well-orderings  $<$  of  $H_\theta$  and for all  $a \in [\lambda]^{\omega_1}$  and for all countable  $X \prec \langle H_\theta; \in, < \rangle$  there is a countable  $Y \prec \langle H_\theta; \in, < \rangle$  such that  $X \sqsubset Y$  and there is some  $b \in Y \cap [\lambda]^{\omega_1}$  such that  $a \subset b$ .

$\text{CC}^{**}$  is  $\text{CC}^*(\lambda)$  for all cardinals  $\lambda \geq \omega_2$ .

Note that  $\text{CC}^*(\omega_2)$  implies Todorćević's  $\text{CC}^*$ ; in the case of  $\text{CC}^*$  one only requires for an  $X$  as above that  $X \sqsubset Y$  and  $X \cap \omega_2 \neq Y \cap \omega_2$ , see [Tod93]. Note that  $\text{CC}^*(\omega_2)$  (and also  $\text{CC}^*$ ) implies the usual Chang Conjecture by building a continuous chain of countable elementary submodels of length  $\omega_1$ ; at each successor stage apply  $\text{CC}^{**}$ . So the countable ordinals of the last model of the chain are the same as the first model's.

The next theorem answers a question of Todorćević who asked the second author under which circumstances  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  is semiproper.

The authors would like to thank Daisuke Ikegami for communicating valuable results about the relationship of  $\text{CC}^{**}$ , SSR and  $(\dagger)$ .

**Theorem 5.7** *The following are equivalent:*

1.  $\text{NS}_{\omega_1}$  is precipitous and for all regular  $\theta \geq \omega_2$  the partial ordering  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  is semiproper.
2. For arbitrarily large  $\theta \geq \omega_2$  there is a semiproper partial order  $\mathbb{P}$  that adds a generic iteration

$$\langle\langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle\rangle, \langle G_i; i < \omega_1 \rangle\rangle.$$

such that  $H_\theta \subset M_{\omega_1}$  and all  $M_i$  are countable.

3.  $\text{CC}^{**}$
4. SSR
5.  $(\dagger)$

Before we prove the above theorem note that the Namba-like forcing in [KLZ07] is stationary set preserving (cf. [Zap]) and hence  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  is not the only example witnessing the consistency of 2.

*Proof.* 1.  $\implies$  2. is trivial and 4.  $\implies$  5. is Lemma 5.3.

5.  $\implies$  1. is clear since by 5.5,  $\text{NS}_{\omega_1}$  is precipitous in this case and so by [CS09] the forcing  $\mathbb{P}(\text{NS}_{\omega_1}, \theta)$  exists for all regular  $\theta \leq \omega_2$  and preserves stationary subsets of  $\omega_1$ .

It remains to show 2.  $\implies$  3. and 3.  $\implies$  4. For the first implication we assume that  $\text{CC}^{**}$  does not hold and work toward a contradiction. So there is a least cardinal  $\lambda_0 \geq \aleph_2$  for which  $\text{CC}^{**}$  fails. Since 2. holds there is a least  $\theta_0 > \lambda_0$  such that a semiproper  $\mathbb{P}$  exists that adds an iteration

$$\langle\langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle\rangle, \langle G_i; i < \omega_1 \rangle\rangle.$$

such that  $H_{\theta_0} \subset M_{\omega_1}$  and all  $M_i$  are countable. Let  $\theta > \theta_0$  large enough so that a name for an iteration as above and  $\mathcal{P}(\mathbb{P})$  are both in  $H_\theta$ . Let  $<$  be some well-ordering of  $H_\theta$ . Now fix some arbitrary  $X \prec \langle H_\theta; \in, < \rangle$  and some  $a \in [\lambda_0]^\omega$ . Our aim is now to construct a  $Y \prec \langle H_\theta; \in, < \rangle$  like in  $\text{CC}^{**}$ . For this we first show that it suffices to do so in a generic extension:

**Claim 1.** If there is some generic extension of  $V$  that contains some  $Y \prec \langle H_\theta; \in, < \rangle$  such that  $X \sqsubset Y$  and there is some  $b \in Y \cap [\lambda_0]^{\omega_1} \cap V$  such that  $a \subset b$  then there is already some  $Z \in V$  with  $Z \prec \langle H_\theta; \in, < \rangle$ ,  $X \sqsubset Z$  and  $b \in Z$ .

*Proof of Claim 1.* If  $Y$  is in some generic extension  $W$  of  $V$ , then by  $b \in V$  there is a tree  $T \in V$  searching for a countable  $Z \prec \langle H_\theta; \in, < \rangle$  such that  $b \in Z$  and  $X \sqsubset Z$ .

So  $T$  has a branch in  $W$ , this is clearly witnessed by  $Y$ . By the absoluteness of well-foundedness we have a branch through  $T$  in  $V$  and hence there is some countable  $Z \prec \langle H_\theta; \in, < \rangle$  with  $X \sqsubset Z$  and  $b \in Z$  in  $V$ .  $\square$ (Claim 1)

By the minimality of  $\lambda_0$  and  $\theta_0$  some semiproper forcing and some name for an iteration as above exist in  $X$ . Let us call this forcing  $\mathbb{P}$  again. Let  $G \subset \mathbb{P}$  be generic over  $V$ .

**Claim 2.**  $X[G] \prec H_\lambda[G]$ .

This claim is part of the folklore. For the readers convenience we give a *Proof of Claim 2.* An induction along the first order formulae will yield the desired result: let  $\phi$  be a formula and let  $\sigma \in X$  denote some name such that

$$H_\lambda[G] \models \exists y \phi(y, \sigma^G).$$

Then by the fullness of the forcing names we have

$$H_\lambda \models \exists \tau \forall p \in \mathbb{P} (p \Vdash \exists y \phi(y, \sigma) \implies p \Vdash \phi(\tau, \sigma)).$$

So by elementarity such a  $\tau$  exists in  $X$ . By the inductive hypothesis we have

$$H_\lambda[G] \models \phi(\tau^G, \sigma^G) \iff X[G] \models \phi(\tau^G, \sigma^G).$$

$\square$ (Claim 2)

By our hypothesis we can force the existence of a generic iteration

$$\dot{J}^G = \langle \langle M_i, \pi_{i,j}, I_i, \kappa_i; i \leq j \leq \omega_1 \rangle, \langle G_i; i < \omega_1 \rangle \rangle.$$

with  $M_{\omega_1} \supset H_\theta$ . So by the regularity of  $\theta$  we have  $a \in M_{\omega_1}$ . Note that  $X[G]$  can calculate  $M_0$ .

**Claim 3.** Let  $\beta < \alpha \leq \omega_1$ . All elements of  $M_\alpha$  are of the form  $\pi_{\beta,\alpha}(f)(\vec{\xi})$  for some  $f : \kappa_\beta^n \rightarrow M_\beta$ ,  $f \in M_\beta$  and ordinals  $\xi_1, \dots, \xi_n < \omega_1^{M_\alpha}$ .

This claim is also part of the folklore. Nevertheless we give a proof for the readers convenience.

*Proof of Claim 3.* Fix  $\beta < \omega_1$ . We show this by induction on  $\alpha$ . Let  $\alpha = \gamma + 1$ . Then  $M_\alpha$  is isomorph to  $\text{Ult}(M_\gamma, G_\gamma)$ . Hence every element of  $M_\alpha$  has the form  $\pi_{\gamma,\alpha}(f)(\kappa_\gamma)$  for some  $f : \kappa_\gamma \rightarrow M_\gamma$ ,  $f \in M_\gamma$ . By the inductive hypothesis  $f$  is of the form  $\pi_{\beta,\gamma}(g)(\vec{\xi})$  for some  $g : \kappa_\beta^n \rightarrow M_\beta$ ,  $g \in M_\beta$  and  $\vec{\xi} \in \kappa_\gamma^n$ . Then

$$\pi_{\gamma,\alpha}(f)(\kappa_\gamma) = \pi_{\gamma,\alpha}(\pi_{\beta,\gamma}(g)(\vec{\xi}))(\kappa_\gamma) = \pi_{\beta,\alpha}(g)(\vec{\xi})(\kappa_\gamma),$$

since the critical point of  $\pi_{\gamma,\alpha}$  is  $\kappa_\gamma$ .

The case  $\text{Lim}(\alpha)$  simply uses the fact that  $M_\alpha$  is the direct limit of all  $M_\gamma$  for  $\gamma < \alpha$ : if  $x \in M_\alpha$ , then  $x = \pi_{\gamma,\alpha}(\bar{x})$  for some  $\gamma < \alpha$  and some  $\bar{x} \in M_\gamma$ . Without loss of generality we may assume  $\beta < \gamma$ . Then  $\bar{x}$  is of the form  $\pi_{\beta,\gamma}(g)(\vec{\xi})$  for some  $g : \kappa_\beta^n \rightarrow M_\beta$ ,  $g \in M_\beta$  and ordinals  $\vec{\xi} \in \kappa_\gamma^n$ . Then

$$x = \pi_{\gamma,\alpha}(\bar{x}) = \pi_{\gamma,\alpha}(\pi_{\beta,\gamma}(g)(\vec{\xi})) = \pi_{\beta,\alpha}(g)(\vec{\xi}).$$

$\square$ (Claim 3)

By setting  $\beta = 0$  and  $\alpha = \omega_1$  in the above claim, we have that there is some  $f \in M_0$ ,  $f : \kappa_0^n \rightarrow M_0$  and  $\vec{\xi} = \xi_1, \dots, \xi_n < \omega_1$  such that

$$a = \pi_{0,\omega_1}(f)(\vec{\xi}).$$

This  $f$  is in  $X[G]$ . We set

$$b := \bigcup \{ \pi_{0,\omega_1}(f)(\vec{\alpha}) ; \vec{\alpha} \in \omega_1^n \wedge \pi_{0,\omega_1}(f)(\vec{\alpha}) \in ([H_\theta]^{\omega_1})^V \}.$$

Clearly  $a \subset b$  and  $\text{Card}(b) = \omega_1$ . Since the parameters  $\pi_{0,\omega_1}(f), [H_\theta]^{\omega_1}$  used in the definition of  $b$  are in  $V$  we have that  $b \in V$ . Also  $b \in X[G]$ . By the semiproperness of  $\mathbb{P} X \sqsubset X[G]$ . So  $X[G]$  witnesses that in some generic extension of  $V$  there is some  $Y$  as desired. This suffices to show by claim 1.

We now show that 3.  $\implies$  4. This implication is a slight generalization of [Tod93, Lemma 6]. Fix an ordinal  $\lambda \geq \omega_2$  and a semistationary  $S \subset [\omega_2]^\omega$ . We set

$$\mathcal{W} := \{ W \subset \lambda ; \text{Card}(W) = \omega_1 \subset W \}$$

and

$$T := \{ y \in [\lambda]^\omega ; \exists x \in S : x \sqsubset y \}.$$

By the very definition of semistationarity  $T$  is stationary. Let us assume that SSR does not hold and work toward a contradiction. For all  $W \in \mathcal{W}$

$$S_W := \{ y \in [W]^\omega ; \exists x \in S \cap [W]^\omega : x \sqsubset y \}$$

is nonstationary. For each  $W \in \mathcal{W}$  we may hence pick a function

$$f_W : [W]^{<\omega} \rightarrow W$$

such that

$$S_W \cap \{ x \in [W]^\omega ; f_W \text{``}[x]^{<\omega} \subset x \} = \emptyset.$$

Let  $\mathcal{F}$  denote the collection of these  $f_W$ . Let  $\theta > \lambda$  be regular large enough such that  $\mathcal{F}, \mathcal{W}, S, T \in H_\theta$  and such that the implications of  $\text{CC}^{**}$  hold for this  $\theta$ . Let  $<$  be a well-ordering of  $H_\theta$ . Pick a countable  $M \prec \langle H_\theta; \in, < \rangle$  such that  $\mathcal{F}, \mathcal{W}, S, T, \lambda \in M$  and

$$M \cap \lambda \in T.$$

Let

$$a := (M \cap \lambda) \cup \omega_1.$$

Since  $\text{CC}^{**}$  holds, there is a countable  $M^* \prec H_\theta$  and some  $b \in [\theta]^{\omega_1}$  such that  $M \sqsubset M^*$ ,  $a \subset b$  and  $b \in M^*$ . Clearly  $W := b \cap \lambda \in \mathcal{W} \cap M^*$ . So  $f_W \in M^*$ . Then by elementarity of  $M^*$

$$f_W \text{``}[W \cap M^*]^{<\omega} \subset W \cap M^*.$$

By the choice of  $a$  and the properties of  $M^*$  we have

$$M \cap \lambda \sqsubset W \cap M^*.$$

Since we have  $M \cap \lambda \in T$  there is some  $x \in S$  such that  $x \sqsubset M \cap \lambda$ . Note that  $x \in [W]^\omega$ . By the transitivity of  $\sqsubset$ ,

$$x \sqsubset W \cap M^*.$$

This implies  $W \cap M^* \in S_W$ . We thus have a contradiction to the choice of  $f_W$ . This finishes the proof.  $\square$



## References

- [AW02] David Asperó and Philip D. Welch. Bounded Martin's maximum, weak Erdős cardinals, and  $\psi_{AC}$ . *J. Symbolic Logic*, 67(3):1141–1152, 2002.
- [CS09] Benjamin Claverie and Ralf Schindler. Increasing  $u_2$  by a stationary set preserving forcing. *J. Symbolic Logic*, 74(1):187–200, 2009.
- [FMS88] M. Foreman, M. Magidor, and S. Shelah. Martin's maximum, saturated ideals, and nonregular ultrafilters. I. *Ann. of Math. (2)*, 127(1):1–47, 1988.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.
- [KLZ07] Richard Ketchersid, Paul Larson, and Jindřich Zapletal. Increasing  $\delta_2^1$  and Namba-style forcing. *J. Symbolic Logic*, 72(4):1372–1378, 2007.
- [She98] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, second edition, 1998.
- [Tod93] Stevo Todorčević. Conjectures of Rado and Chang and cardinal arithmetic. In *Finite and infinite combinatorics in sets and logic (Banff, AB, 1991)*, volume 411 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 385–398. Kluwer Acad. Publ., Dordrecht, 1993.
- [Woo99] W. Hugh Woodin. *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, volume 1 of *de Gruyter Series in Logic and its Applications*. Walter de Gruyter & Co., Berlin, 1999.
- [Zap] Jindřich Zapletal. private communication.