

Pure Anderson Motives over Finite Fields

Matthias Bornhofen, Urs Hartl *

October 30, 2008

Abstract

In the arithmetic of function fields Drinfeld modules play the role that elliptic curves take on in the arithmetic of number fields. As higher dimensional generalizations of Drinfeld modules, and as the appropriate analogues of abelian varieties, G. Anderson introduced pure t -motives. In this article we study the arithmetic of the latter. We investigate which pure t -motives are semisimple, that is, isogenous to direct sums of simple ones. We give examples for pure t -motives which are not semisimple. Over finite fields the semisimplicity is equivalent to the semisimplicity of the endomorphism algebra, but also this fails over infinite fields. Still over finite fields we study the Zeta function and the endomorphism rings of pure t -motives and criteria for the existence of isogenies. We obtain answers which are similar to Tate's famous results for abelian varieties.

Mathematics Subject Classification (2000): 11G09, (13A35, 16K20)

Introduction

In the last decades the Arithmetic of Function Fields has acquired great impetus caused by Drinfeld's [Dr1, Dr2] invention of the concepts of *elliptic modules* (today called *Drinfeld modules*) and *elliptic sheaves* in the 1970s. Both are analogues of elliptic curves. The latter live in the Arithmetic of Number Fields, like their higher dimensional generalizations abelian varieties. In [BH1, Ha1] we claimed that pure Anderson motives (a slight generalization of the pure t -motives introduced by Anderson [An1]) and abelian τ -sheaves should be viewed as the appropriate analogues for abelian varieties and higher dimensional generalizations of elliptic sheaves or modules. We want to further support this claim in the present article by developing the theory of pure Anderson motives over finite fields.

To give the definition of pure Anderson motives let C be a connected smooth projective curve over \mathbb{F}_q , let $\infty \in C(\mathbb{F}_q)$ be a fixed point, and let $A = \Gamma(C \setminus \{\infty\}, \mathcal{O}_C)$. For a field $L \supset \mathbb{F}_q$ let σ^* be the endomorphism of $A_L := A \otimes_{\mathbb{F}_q} L$ sending $a \otimes b$ to $a \otimes b^q$ for $a \in A$ and $b \in L$. Let $c^* : A \rightarrow L$ be an \mathbb{F}_q -homomorphism and let $J = (a \otimes 1 - 1 \otimes c^*(a) : a \in A) \subset A_L$. A *pure Anderson motive* $\underline{M} = (M, \tau)$ of rank r , dimension d and characteristic c^* consists of a locally free A_L -module M of rank r and an A_L -homomorphism $\tau : \sigma^* M := M \otimes_{A_L, \sigma^*} A_L \rightarrow M$ with $\dim_L \operatorname{coker} \tau = d$ and $J^d \cdot \operatorname{coker} \tau = 0$, such that M possesses an extension to a locally free sheaf \mathcal{M} on $C \times_{\mathbb{F}_q} L$ on which $\tau^l : (\sigma^*)^l \mathcal{M} \rightarrow \mathcal{M}(k \cdot \infty)$ is an isomorphism near ∞ for some positive integers k and l . The last condition is the purity condition. The ratio $\frac{k}{l}$ equals $\frac{d}{r}$ and is called the *weight of \underline{M}* . Anderson's definition of pure t -motives [An1] is recovered by setting $C = \mathbb{P}_{\mathbb{F}_q}^1$ and $A = \mathbb{F}_q[t]$. In the first two sections we recall the definition of morphisms and isogenies between pure Anderson motives as well as some facts from [BH1]. Also for an isogeny f between pure Anderson motives we define the degree of f as an ideal of A (2.8) which annihilates $\operatorname{coker} f$ (2.10). If \underline{M} is a semisimple (see below) pure Anderson motive over a finite field, the degree of any isogeny $f : \underline{M} \rightarrow \underline{M}$ is a principal ideal and has a canonical generator (7.3). In particular f has a canonical dual.

*The second author acknowledges support of the Deutsche Forschungsgemeinschaft in form of DFG-grant HA3006/2-1 and SFB 478

Next we address the question whether every pure Anderson motive is *semisimple*, that is, isogenous to a direct sum of simple pure Anderson motives. A pure Anderson motive is called *simple* if it has no non-trivial factor motives. This question is the analogue of the classical theorem of Poincaré-Weil on the semisimplicity of abelian varieties. By giving a counterexample (Example 6.1) we demonstrate that the answer to this question is negative in general. On the positive side we show that every pure Anderson motive over a finite base field becomes semisimple after a field extension whose degree is a power of q (6.15), and then stays semisimple after any further field extension (6.16). Let Q be the function field of C . Then the endomorphism Q -algebra $\mathrm{QEnd}(\underline{M}) := \mathrm{End}(\underline{M}) \otimes_A Q$ of a semisimple pure Anderson motive is semisimple (2.7) and over a finite field also the converse is true (6.11). This is false however over an infinite field (6.13).

Like for abelian varieties the behavior of a pure Anderson motive \underline{M} over a finite field is controlled by its Frobenius endomorphism π (defined in 5.2). If \underline{M} is semisimple we determine the dimension and the local Hasse invariants of its endomorphism Q -algebra $\mathrm{QEnd}(\underline{M})$ in terms of π (6.5, 9.1). We define a Zeta function $Z_{\underline{M}}$ for a pure Anderson motive \underline{M} (Definition 7.6) and we show that it satisfies the Riemann hypothesis (7.8), and has an expression in terms of the degrees $\deg(1 - \pi^i)$ for all i if \underline{M} is semisimple (7.7). We prove the following isogeny criterion.

Theorem 8.1. *Let \underline{M} and \underline{M}' be semisimple pure Anderson motives over a finite field and let π , respectively π' , be their Frobenius endomorphisms. Then the following are equivalent:*

1. \underline{M} and \underline{M}' are isogenous.
2. The characteristic polynomials of π and π' acting on the v -adic Tate modules of \underline{M} , respectively \underline{M}' , coincide for some (any) place $v \in \mathrm{Spec} A$.
3. There exists an isomorphism of Q -algebras $\mathrm{QEnd}(\underline{M}) \cong \mathrm{QEnd}(\underline{M}')$ mapping π to π' .
4. $Z_{\underline{M}} = Z_{\underline{M}'}$.

In the last section we sketch a few results for the question, which orders of $\mathrm{QEnd}(\underline{M})$ occur as the endomorphism rings of pure Anderson motives (10.7, 10.11). There is a relation between the breaking up of the isogeny class of a semisimple pure Anderson motive into isomorphism classes, and the arithmetic of $\mathrm{QEnd}(\underline{M})$. We indicate this by treating the case of pure Anderson motives defined over the minimal field \mathbb{F}_q . In this case $\mathrm{QEnd}(\underline{M})$ is commutative (10.11). Many of our results parallel Tate's celebrated article [Tat] on abelian varieties over finite fields. To prove them, a major tool are the Tate modules and local shtuka attached to pure Anderson motives, which we recall in Sections 4 and 3, and the analogue [Tag, Tam] of Tate's conjecture on endomorphisms. These local structures behave like in the classical case of abelian varieties, local shtuka playing the role of the p -divisible groups of the abelian varieties. The only difference is that p -divisible groups are only useful for abelian varieties in characteristic p , whereas the local shtuka at any place of Q are important for the investigation of abelian τ -sheaves and pure Anderson motives. One of the aims of this article is to demonstrate the utility of local shtuka. For instance we apply them in the computation of the Hasse invariants of $\mathrm{QEnd}(\underline{M})$ in Theorem 9.1. We also used them in [BH1] to reprove the standard fact that the set of morphisms between two pure Anderson motives is a projective A -module (1.3). Scattered in the text are several interesting examples displaying various phenomena (6.1, 6.13, 9.4, 9.5). Note that there is a two in one version [BH2] of the present article and [BH1] on the arXiv.

Contents

Introduction	1
1 Pure Anderson Motives and Abelian τ-Sheaves	4

2	Isogenies and Quasi-Isogenies	6
3	Local Shtuka	8
4	Tate Modules	12
5	The Frobenius Endomorphism	13
6	The Poincaré-Weil Theorem	14
7	Zeta Functions and Reduced Norms	21
8	A Quasi-Isogeny Criterion	23
9	The Endomorphism Q-Algebra	25
10	Kernel Ideals for Pure Anderson Motives	31
	References	34

Notation

In this article we denote by

\mathbb{F}_q	the finite field with q elements and characteristic p ,
C	a smooth projective geometrically irreducible curve over \mathbb{F}_q ,
$\infty \in C(\mathbb{F}_q)$	a fixed \mathbb{F}_q -rational point on C ,
$A = \Gamma(C \setminus \{\infty\}, \mathcal{O}_C)$	the ring of regular functions on C outside ∞ ,
$Q = \mathbb{F}_q(C) = \text{Quot}(A)$	the function field of C ,
Q_v	the completion of Q at the place $v \in C$,
A_v	the ring of integers in Q_v . For $v \neq \infty$ it is the completion of A at v .
\mathbb{F}_v	the residue field of A_v . In particular $\mathbb{F}_\infty \cong \mathbb{F}_q$.

For a field L containing \mathbb{F}_q we write

$C_L = C \times_{\text{Spec } \mathbb{F}_q} \text{Spec } L$,	
$A_L = A \otimes_{\mathbb{F}_q} L$,	
$Q_L = Q \otimes_{\mathbb{F}_q} L$,	
$A_{v,L} = A_v \widehat{\otimes}_{\mathbb{F}_q} L$	for the completion of \mathcal{O}_{C_L} at the closed subscheme $v \times \text{Spec } L$,
$Q_{v,L} = A_{v,L}[\frac{1}{v}]$.	Note that this is not a field if $\mathbb{F}_v \cap L \supsetneq \mathbb{F}_q$,
$\text{Frob}_q : L \rightarrow L$	for the q -Frobenius endomorphism mapping x to x^q ,
$\sigma = \text{id}_C \times \text{Spec}(\text{Frob}_q)$	for the endomorphism of C_L which acts as the identity on the points and on \mathcal{O}_C and as the q -Frobenius on L ,
σ^*	for the endomorphisms induced by σ on all the above rings. For instance $\sigma^*(a \otimes b) = a \otimes b^q$ for $a \in A$ and $b \in L$.
$\sigma^*M = M \otimes_{A_L, \sigma^*} A_L$	for an A_L -module M and similarly for the other rings.

For a divisor D on C we denote by $\mathcal{O}_{C_L}(D)$ the invertible sheaf on C_L whose sections φ have divisor $(\varphi) \geq -D$. For a coherent sheaf \mathcal{F} on C_L we set $\mathcal{F}(D) := \mathcal{F} \otimes_{\mathcal{O}_{C_L}} \mathcal{O}_{C_L}(D)$. This notation applies in particular to the divisor $D = n \cdot \infty$ for $n \in \mathbb{Z}$.

We will fix the further notation $\pi, F, E, \mu_\pi, \pi_v, F_v, E_v$, and χ_v in formula (6.1) on page 15.

1 Pure Anderson Motives and Abelian τ -Sheaves

Pure Anderson motives were introduced by G. Anderson [An1] under the name *pure t -motives* in the case where $A = \mathbb{F}_p[t]$. They were further studied in [BH1]. To give their definition let L be a field extension of \mathbb{F}_q and fix an \mathbb{F}_q -homomorphism $c^* : A \rightarrow L$. Let $J \subset A_L$ be the ideal generated by $a \otimes 1 - 1 \otimes c^*(a)$ for all $a \in A$.

Definition 1.1 (pure Anderson motives). *A pure Anderson motive $\underline{M} = (M, \tau)$ of rank r , dimension d , and characteristic c^* over L consists of a locally free A_L -module M of rank r and an injective A_L -module homomorphism $\tau : \sigma^* M \rightarrow M$ such that*

1. *the cokernel of τ is an L -vector space of dimension d and annihilated by J^d , and*
2. *M extends to a locally free sheaf \mathcal{M} of rank r on C_L such that for some positive integers k, l the map $\tau^l := \tau \circ \sigma^*(\tau) \circ \dots \circ (\sigma^*)^{l-1}(\tau) : (\sigma^*)^l M \rightarrow M$ induces an isomorphism $(\sigma^*)^l \mathcal{M}_\infty \rightarrow \mathcal{M}(k \cdot \infty)_\infty$ of the stalks at ∞ .*

We call $\varepsilon := \ker c^* \in \text{Spec } A$ the characteristic point of \underline{M} and we say that \underline{M} has finite characteristic (respectively generic characteristic) if ε is a closed (respectively the generic) point. The ratio $\text{wt}(M, \tau) := \frac{k}{l}$ equals $\frac{d}{r}$ and is called the weight of (M, τ) ; see [BH1, Proposition 1.2].

Definition 1.2. (Compare [PT, 4.5])

1. *A morphism $f : (M, \tau) \rightarrow (M', \tau')$ between Anderson motives of the same characteristic c^* is a morphism $f : M \rightarrow M'$ of A_L -modules which satisfies $f \circ \tau = \tau' \circ \sigma^*(f)$.*
2. *If $f : \underline{M} \rightarrow \underline{M}'$ is surjective, \underline{M}' is called a factor motive of \underline{M} .*
3. *A morphism $f : \underline{M} \rightarrow \underline{M}'$ is called an isogeny if f is injective with torsion cokernel.*
4. *An isogeny is called separable (respectively purely inseparable) if the induced morphism $\tau : \sigma^* \text{coker } f \rightarrow \text{coker } f$ is an isomorphism (respectively is nilpotent, that is, if for some n the morphism $\tau \circ \sigma^* \tau \circ \dots \circ (\sigma^*)^n \tau$ is zero).*

We denote the set of morphisms between \underline{M} and \underline{M}' by $\text{Hom}(\underline{M}, \underline{M}')$. It is an A -module.

If \underline{M} and \underline{M}' are pure Anderson motives of different weights then $\text{Hom}(\underline{M}, \underline{M}') = \{0\}$ by [BH1, Corollary 3.5]. This justifies the terminology *pure*. The following fact is well known. A proof can be found for instance in [BH1, Theorem 9.5].

Theorem 1.3. *Let \underline{M} and \underline{M}' be pure Anderson motives over an arbitrary field L . Then $\text{Hom}(\underline{M}, \underline{M}')$ is a projective A -module of rank $\leq rr'$. The minimal polynomial of every endomorphism of a pure Anderson motive \underline{M} lies in $A[x]$. \square*

Corollary 1.4. ([BH1, Corollary 5.4]) *Let $f : \underline{M} \rightarrow \underline{M}'$ be an isogeny between pure Anderson motives. Then*

1. *there exists an element $a \in A$ which annihilates $\text{coker } f$,*
2. *there exists a dual isogeny $f^\vee : \underline{M}' \rightarrow \underline{M}$ such that $f \circ f^\vee = a \cdot \text{id}_{\underline{M}'}$ and $f^\vee \circ f = a \cdot \text{id}_{\underline{M}}$.*

Next we come to the notion of abelian (τ -)sheaves. It was introduced in [Ha1] in order to construct moduli spaces for pure Anderson motives. We briefly recall the results from [BH1] on the relation between pure Anderson motives and abelian τ -sheaves. Although our primary interest is on pure Anderson motives we present abelian τ -sheaves here because they can have characteristic point $\varepsilon = \infty \in C$ in contrast to pure Anderson motives, and many results for the later extend to this more general situation. Moreover, some results are proved most naturally via the use of abelian τ -sheaves (e.g. 9.1 and 7.3 below). The fact that $\varepsilon = \infty$ is allowed for abelian τ -sheaves was crucial for the uniformization of the moduli spaces of pure Anderson motives in [Ha1] and the derived consequences on analytic uniformization of pure Anderson motives. Let $L \supset \mathbb{F}_q$ be a field and fix a morphism $c : \text{Spec } L \rightarrow C$. Let \mathcal{J} be the ideal sheaf on C_L of the graph of c . Let r and d be non-negative integers.

Definition 1.5 (Abelian τ -sheaf). An abelian τ -sheaf $\underline{\mathcal{F}} = (\mathcal{F}_i, \Pi_i, \tau_i)$ of rank r , dimension d and characteristic c over L is a collection of locally free sheaves \mathcal{F}_i on C_L of rank r together with injective morphisms Π_i, τ_i of \mathcal{O}_{C_L} -modules ($i \in \mathbb{Z}$) of the form

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \mathcal{F}_{i-1} & \xrightarrow{\Pi_{i-1}} & \mathcal{F}_i & \xrightarrow{\Pi_i} & \mathcal{F}_{i+1} \xrightarrow{\Pi_{i+1}} \cdots \\ & & \uparrow \tau_{i-2} & & \uparrow \tau_{i-1} & & \uparrow \tau_i \\ \cdots & \longrightarrow & \sigma^* \mathcal{F}_{i-2} & \xrightarrow{\sigma^* \Pi_{i-2}} & \sigma^* \mathcal{F}_{i-1} & \xrightarrow{\sigma^* \Pi_{i-1}} & \sigma^* \mathcal{F}_i \xrightarrow{\sigma^* \Pi_i} \cdots \end{array}$$

subject to the following conditions:

1. the above diagram is commutative,
2. there exist integers $k, l > 0$ with $ld = kr$ such that the morphism $\Pi_{i+l-1} \circ \cdots \circ \Pi_i$ identifies \mathcal{F}_i with the subsheaf $\mathcal{F}_{i+l}(-k \cdot \infty)$ of \mathcal{F}_{i+l} for all $i \in \mathbb{Z}$,
3. $\text{coker } \Pi_i$ is an L -vector space of dimension d for all $i \in \mathbb{Z}$,
4. $\text{coker } \tau_i$ is an L -vector space of dimension d and annihilated by \mathcal{J}^d for all $i \in \mathbb{Z}$.

We call $\varepsilon := c(\text{Spec } L) \in C$ the characteristic point and say that $\underline{\mathcal{F}}$ has finite (respectively generic) characteristic if ε is a closed (respectively the generic) point. If $r \neq 0$ we call $\text{wt}(\underline{\mathcal{F}}) := \frac{d}{r}$ the weight of $\underline{\mathcal{F}}$.

Remark. 1. If $\underline{\mathcal{F}}$ is an abelian τ -sheaf and D a divisor on C , then $\underline{\mathcal{F}}(D) := (\mathcal{F}_i(D), \Pi_i \otimes 1, \tau_i \otimes 1)$ is an abelian τ -sheaf of the same rank and dimension as $\underline{\mathcal{F}}$.

2. Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf and let $n \in \mathbb{Z}$. We denote by $\underline{\mathcal{F}}[n] := (\mathcal{F}_{i+n}, \Pi_{i+n}, \tau_{i+n})$ the n -shifted abelian τ -sheaf of $\underline{\mathcal{F}}$ whose collection of \mathcal{F} 's, Π 's and τ 's is just shifted by n .

Definition 1.6. A morphism f between two abelian τ -sheaves $\underline{\mathcal{F}} = (\mathcal{F}_i, \Pi_i, \tau_i)$ and $\underline{\mathcal{F}}' = (\mathcal{F}'_i, \Pi'_i, \tau'_i)$ of the same characteristic $c : \text{Spec } L \rightarrow C$ is a collection of morphisms $f_i : \mathcal{F}_i \rightarrow \mathcal{F}'_i$ ($i \in \mathbb{Z}$) which commute with the Π 's and the τ 's, that is, $f_{i+1} \circ \Pi_i = \Pi'_i \circ f_i$ and $f_{i+1} \circ \tau_i = \tau'_i \circ \sigma^* f_i$. We denote the set of morphisms between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ by $\text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$. It is an \mathbb{F}_q -vector space.

For example, the collection of morphisms $(\Pi_i) : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}[1]$ defines a morphism between the abelian τ -sheaves $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}[1]$.

Definition 1.7. Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ be abelian τ -sheaves and let $f \in \text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ be a morphism. Then f is called injective (respectively surjective, respectively an isomorphism), if f_i is injective (respectively surjective, respectively bijective) for all $i \in \mathbb{Z}$. We call $\underline{\mathcal{F}}$ an abelian factor τ -sheaf of $\underline{\mathcal{F}}'$, if there is a surjective morphism from $\underline{\mathcal{F}}'$ onto $\underline{\mathcal{F}}$.

If $\underline{\mathcal{F}} = (\mathcal{F}_i, \Pi_i, \tau_i)$ is an abelian τ -sheaf of rank r , dimension d , and characteristic $c : \text{Spec } L \rightarrow C$ with $\varepsilon = \text{im } c \neq \infty$ then

$$\underline{M}(\underline{\mathcal{F}}) := (M, \tau) := \left(\Gamma(C_L \setminus \{\infty\}, \mathcal{F}_0), \Pi_0^{-1} \circ \tau_0 \right) \quad (1.1)$$

is a pure Anderson motive of the same rank and dimension and of characteristic $c^* : A \rightarrow L$. Conversely we have the following result.

Proposition 1.8. ([BH1, Theorem 3.1]) Let (M, τ) be a pure Anderson motive of rank r , dimension d , and characteristic $c^* : A \rightarrow L$ over L . Then $(M, \tau) = \underline{M}(\underline{\mathcal{F}})$ for an abelian τ -sheaf $\underline{\mathcal{F}}$ over L of same rank and dimension with characteristic $c := \text{Spec } c^* : \text{Spec } L \rightarrow \text{Spec } A \subset C$. One can even find the abelian τ -sheaf $\underline{\mathcal{F}}$ with k, l relatively prime.

2 Isogenies and Quasi-Isogenies

We recall the basic facts about isogenies from [BH1].

Proposition 2.1. ([BH1, Proposition 5.1]) *Let $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'$ be a morphism between two abelian τ -sheaves $\underline{\mathcal{F}} = (\mathcal{F}_i, \Pi_i, \tau_i)$ and $\underline{\mathcal{F}}' = (\mathcal{F}'_i, \Pi'_i, \tau'_i)$. Then the following assertions are equivalent:*

1. f is injective and the support of all $\text{coker } f_i$ is contained in $D \times \text{Spec } L$ for a finite closed subscheme $D \subset C$,
2. f is injective and $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ have the same rank and dimension,
3. $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ have the same weight and the fiber $f_{i,\eta}$ at the generic point η of C_L is an isomorphism for some (any) $i \in \mathbb{Z}$.

Definition 2.2. 1. A morphism $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'$ satisfying the equivalent conditions of Proposition 2.1 is called an *isogeny*. We denote the set of isogenies between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ by $\text{Isog}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$.

2. An isogeny $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'$ is called *separable* (respectively *purely inseparable*) if for all i the induced morphism $\tau_i : \sigma^* \text{coker } f_i \rightarrow \text{coker } f_{i+1}$ is an isomorphism (respectively is nilpotent, that is, $\tau_i \circ \sigma^* \tau_{i-1} \circ \dots \circ (\sigma^*)^n \tau_{i-n} = 0$ for some n).

The endomorphism rings of abelian τ -sheaves are finite rings. But if we allow the (endo-)morphisms to have ‘‘poles’’ we get rings which are related to the endomorphism rings of the associated pure Anderson motives. We make the following:

Definition 2.3 (Quasi-morphism and quasi-isogeny). *Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ be abelian τ -sheaves.*

1. A quasi-morphism f between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ is a morphism $f \in \text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}'(D))$ for some effective divisor D on C .
2. A quasi-isogeny f between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ is an isogeny $f \in \text{Isog}(\underline{\mathcal{F}}, \underline{\mathcal{F}}'(D))$ for some effective divisor D on C .

If $D_1 \leq D_2$ the composition with the inclusion isogeny $\underline{\mathcal{F}}'(D_1) \subset \underline{\mathcal{F}}'(D_2)$ defines an injection $\text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}'(D_1)) \hookrightarrow \text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}'(D_2))$. This yields an equivalence relation for quasi-morphisms and quasi-isogenies. We let $\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ and $\text{QIsog}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ be the set of quasi-morphisms, respectively quasi-isogenies, between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ modulo this equivalence relation. We write $\text{QEnd}(\underline{\mathcal{F}}) := \text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}})$ and $\text{QIsog}(\underline{\mathcal{F}}) := \text{QIsog}(\underline{\mathcal{F}}, \underline{\mathcal{F}})$.

The Q -vector spaces $\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ and $\text{QEnd}(\underline{\mathcal{F}})$ are finite dimensional, and $\text{QIsog}(\underline{\mathcal{F}})$ is the group of units in the Q -algebra $\text{QEnd}(\underline{\mathcal{F}})$, see [BH1, Propositions 6.5 and 9.4].

Two abelian τ -sheaves $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ are called *quasi-isogenous* (notation: $\underline{\mathcal{F}} \approx \underline{\mathcal{F}}'$), if there exists a quasi-isogeny between $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$. The relation \approx is an equivalence relation. If $\underline{\mathcal{F}} \approx \underline{\mathcal{F}}'$, then the Q -algebras $\text{QEnd}(\underline{\mathcal{F}})$ and $\text{QEnd}(\underline{\mathcal{F}}')$ are isomorphic, and $\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ is free of rank 1 both as a left module over $\text{QEnd}(\underline{\mathcal{F}}')$ and as a right module over $\text{QEnd}(\underline{\mathcal{F}})$.

Proposition 2.4. ([BH1, Proposition 6.10]) *Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ be two abelian τ -sheaves of characteristic $\varepsilon \neq \infty$ and let $\underline{M}(\underline{\mathcal{F}})$ and $\underline{M}(\underline{\mathcal{F}}')$ be their associated pure Anderson motives. Then there is a canonical isomorphism of Q -vector spaces*

$$\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}') = \text{Hom}(\underline{M}(\underline{\mathcal{F}}), \underline{M}(\underline{\mathcal{F}}')) \otimes_A Q.$$

If \underline{M} and \underline{M}' are pure Anderson motives, then the elements of $\text{Hom}(\underline{M}, \underline{M}') \otimes_A Q$ which admit an inverse in $\text{Hom}(\underline{M}', \underline{M}) \otimes_A Q$ are called *quasi-isogenies*. With this definition we can reformulate Propositions 1.8 and 2.4 as follows.

Corollary 2.5. *Let $\varepsilon \neq \infty$. Then the functor $\underline{\mathcal{F}} \mapsto \underline{M}(\underline{\mathcal{F}})$ defines an equivalence of categories between*

1. the category with abelian τ -sheaves as objects and with $\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ as the set of morphisms,

2. and the category with pure Anderson motives as objects and with $\text{Hom}(\underline{M}, \underline{M}') \otimes_A Q$ as the set of morphisms.

We call these the quasi-isogeny categories of abelian τ -sheaves of characteristic different from ∞ and of pure Anderson motives, respectively. \square

Definition 2.6. Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf.

1. $\underline{\mathcal{F}}$ is called simple, if $\underline{\mathcal{F}} \neq \underline{0}$ and $\underline{\mathcal{F}}$ has no abelian factor τ -sheaves other than $\underline{0}$ and $\underline{\mathcal{F}}$.
2. $\underline{\mathcal{F}}$ is called semisimple, if $\underline{\mathcal{F}}$ admits, up to quasi-isogeny, a decomposition into a direct sum $\underline{\mathcal{F}} \approx \underline{\mathcal{F}}_1 \oplus \cdots \oplus \underline{\mathcal{F}}_n$ of simple abelian τ -sheaves $\underline{\mathcal{F}}_j$ ($1 \leq j \leq n$).

We make the same definition for a pure Anderson motive.

Remark. 1. Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf with characteristic different from ∞ . Then $\underline{\mathcal{F}}$ is (semi-)simple if and only if the pure Anderson motive $\underline{M}(\underline{\mathcal{F}})$ is (semi-)simple by [BH1, Proposition 7.3].

2. It is not sensible to try defining *simple* pure Anderson motives via sub-motives, since for example $a\underline{M} \subset \underline{M}$ is a proper sub-motive for any $a \in A \setminus A^\times$. This shows that pure Anderson motives behave dually to abelian varieties. Namely an abelian variety is called simple if it has no non-trivial abelian subvarieties.

Theorem 2.7. ([BH1, Theorem 7.8]) Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf of characteristic different from ∞ .

1. If $\underline{\mathcal{F}}$ is simple, then $\text{QEnd}(\underline{\mathcal{F}})$ is a division algebra over Q .
2. If $\underline{\mathcal{F}}$ is semisimple with decomposition $\underline{\mathcal{F}} \approx \underline{\mathcal{F}}_1 \oplus \cdots \oplus \underline{\mathcal{F}}_n$ up to quasi-isogeny into simple abelian τ -sheaves $\underline{\mathcal{F}}_j$, then $\text{QEnd}(\underline{\mathcal{F}})$ decomposes into a finite direct sum of full matrix algebras over the division algebras $\text{QEnd}(\underline{\mathcal{F}}_j)$ over Q .

In the following we want to define the *degree* of an isogeny which should be an ideal of A since in the function field case we have substituted A for \mathbb{Z} . Let $f : \underline{M} \rightarrow \underline{M}'$ be an isogeny between pure Anderson motives. Then the A_L -module $\text{coker } f$ is a finite L -vector space equipped with a morphism of A_L -modules $\tau' : \sigma^* \text{coker } f \rightarrow \text{coker } f$. Since $\text{coker } f$ is annihilated by an element of A it decomposes by the Chinese remainder theorem

$$(\text{coker } f, \tau') = \bigoplus_{v \in \text{supp}(\text{coker } f)} (\text{coker } f, \tau') \otimes_A A_v =: \bigoplus_{v \in \text{supp}(\text{coker } f)} \underline{K}_v.$$

If $v \neq \varepsilon$ the morphism τ' on \underline{K}_v is an isomorphism and so Lang's theorem implies that

$$(\underline{K}_v \otimes_L L^{\text{sep}})^\tau \otimes_{\mathbb{F}_q} L^{\text{sep}} \xrightarrow{\sim} \underline{K}_v \otimes_L L^{\text{sep}}$$

is an isomorphism; see for instance [An1, Lemma 1.8.2]. In particular

$$[\mathbb{F}_v : \mathbb{F}_q] \cdot \dim_{\mathbb{F}_v} (\underline{K}_v \otimes_L L^{\text{sep}})^\tau = \dim_{\mathbb{F}_q} (\underline{K}_v \otimes_L L^{\text{sep}})^\tau = \dim_{L^{\text{sep}}} (\underline{K}_v \otimes_L L^{\text{sep}}) = \dim_L \underline{K}_v.$$

On the other hand if the characteristic is finite and $v = \varepsilon$, the characteristic morphism $c^* : A \rightarrow L$ yields $\mathbb{F}_\varepsilon \subset L$ and determines the distinguished prime ideal

$$\mathfrak{a}_0 := (b \otimes 1 - 1 \otimes c^*(b) : b \in \mathbb{F}_\varepsilon) \subset A_{\varepsilon, L}.$$

If we set $n := [\mathbb{F}_\varepsilon : \mathbb{F}_q]$ and $\mathfrak{a}_i := (\sigma^*)^i \mathfrak{a}_0 = (b \otimes 1 - 1 \otimes c^*(b)^{q^i} : b \in \mathbb{F}_\varepsilon)$, then we can decompose $A_{\varepsilon, L} = \bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} A_{\varepsilon, L} / \mathfrak{a}_i$ and τ is an isomorphism

$$\sigma^* (\underline{K}_\varepsilon / \mathfrak{a}_{i-1} \underline{K}_\varepsilon) \xrightarrow{\sim} \underline{K}_\varepsilon / \mathfrak{a}_i$$

for $i \neq 0$ since τ is an isomorphism on \underline{M} and \underline{M}' outside the graph of c^* . (This argument will be used again in Proposition 3.8.) In particular

$$[\mathbb{F}_\varepsilon : \mathbb{F}_q] \cdot \dim_L (\underline{K}_\varepsilon / \mathfrak{a}_0 \underline{K}_\varepsilon) = \dim_L \underline{K}_\varepsilon.$$

Definition 2.8. We assign to the isogeny f the ideal

$$\deg(f) := \prod_{v \in \text{supp}(\text{coker } f)} v^{(\dim_L \underline{K}_v)/[\mathbb{F}_v : \mathbb{F}_q]} = \varepsilon^{\dim_L(\underline{K}_\varepsilon/\mathfrak{a}_0 \underline{K}_\varepsilon)} \cdot \prod_{v \neq \varepsilon} v^{\dim_{\mathbb{F}_v}(\underline{K}_v \otimes_L L^{\text{sep}})^\tau}$$

of A and call it the degree of f . We call $\varepsilon^{\dim_L(\underline{K}_\varepsilon/\mathfrak{a}_0 \underline{K}_\varepsilon)}$ the inseparability degree of f and $\prod_{v \neq \varepsilon} v^{\dim_{\mathbb{F}_v}(\underline{K}_v \otimes_L L^{\text{sep}})^\tau}$ the separability degree of f .

Remark. The separability degree of f is the Euler-Poincaré characteristic $EP(\bigoplus_{v \neq \varepsilon} \underline{K}_v \otimes_L L^{\text{sep}})^\tau$; see Gekeler [Gek, 3.9] or Pink and Traulsen [PT, 4.6]. Recall that the Euler-Poincaré characteristic of a finite torsion A -module is the ideal of A defined by requiring that EP is multiplicative in short exact sequences, and that $EP(A/v) := v$ for any maximal ideal v of A .

Lemma 2.9. 1. If $f : \underline{M} \rightarrow \underline{M}'$ and $g : \underline{M}' \rightarrow \underline{M}''$ are isogenies then $\deg(gf) = \deg(f) \cdot \deg(g)$.
2. $\dim_{\mathbb{F}_q} A/\deg(f) = \dim_L \text{coker } f$.

Proof. 1 is immediate from the short exact sequence

$$0 \longrightarrow \text{coker } f \xrightarrow{g} \text{coker}(gf) \longrightarrow \text{coker } g \longrightarrow 0$$

and 2 is obvious. □

Proposition 2.10. The ideal $\deg(f)$ annihilates $\text{coker } f$.

Proof. If $v = \varepsilon$ and a is a uniformizer at ε , then multiplication with a is nilpotent on the L -vector space $\underline{K}_\varepsilon/\mathfrak{a}_0 \underline{K}_\varepsilon$. In particular $a^{\dim_L(\underline{K}_\varepsilon/\mathfrak{a}_0 \underline{K}_\varepsilon)}$ annihilates $\underline{K}_\varepsilon/\mathfrak{a}_0 \underline{K}_\varepsilon$, and hence also $\underline{K}_\varepsilon$.

If $v \neq \varepsilon$ and a is a uniformizer at v , we obtain analogously that $a^{\dim_{\mathbb{F}_v}(\underline{K}_v \otimes_L L^{\text{sep}})^\tau}$ annihilates the \mathbb{F}_v -vector space $(\underline{K}_v \otimes_L L^{\text{sep}})^\tau$ and therefore also the L -vector space \underline{K}_v . □

Proposition 2.11. Let $f : \underline{M} \rightarrow \underline{M}'$ be an isogeny such that $\deg(f) = aA$ is principal (for example this is the case if $C = \mathbb{P}^1$ and $A = \mathbb{F}_q[t]$). Then there is a uniquely determined dual isogeny $f^\vee : \underline{M}' \rightarrow \underline{M}$ (depending on a), which satisfies $f \circ f^\vee = a \cdot \text{id}_{\underline{M}'}$ and $f^\vee \circ f = a \cdot \text{id}_{\underline{M}}$.

Proof. Since $\deg(f)$ annihilates $\text{coker } f$ the proposition is immediate. □

In Theorem 7.3 we will see that any isogeny $f \in \text{End}(\underline{M})$ of a semisimple pure Anderson motive over a finite field satisfies the assumption that $\deg(f)$ is principal.

3 Local Shtuka

There are mainly two local structures which one can attach to pure Anderson motives and abelian τ -sheaves, namely the *local (iso-)shtuka* and the *Tate module*. We treat the Tate module in the next section. The local (iso-)shtuka is the analogue of the Dieudonné module of the p -divisible group attached to an abelian variety. Note however one fundamental difference. While the Dieudonné module exists only if p equals the characteristic of the base field, there is no such restriction in our theory here. And in fact this would even allow to dispense with Tate modules at all and only work with local (iso-)shtuka. Local (iso-)shtuka were introduced in [Ha1] under the name *Dieudonné $\mathbb{F}_q[[z]]$ -modules* (respectively *Dieudonné $\mathbb{F}_q((z))$ -modules*). They are studied in [An2, Ha2, Lau]. Over a field their definition takes the following form.

Definition 3.1. Let $v \in C$ be a place of Q and let $L \supset \mathbb{F}_q$ be a field. An (effective) local σ -shtuka at v of rank r over L is a pair $\hat{\underline{M}} = (\hat{M}, \phi)$ consisting of a free $A_{v,L}$ -module \hat{M} of rank r and an injective $A_{v,L}$ -module homomorphism $\phi : \sigma^* \hat{M} \rightarrow \hat{M}$.

A local σ -isoshtuka at v of rank r over L is a pair $\hat{\underline{N}} = (\hat{N}, \phi)$ consisting of a free $Q_{v,L}$ -module \hat{N} of rank r and an isomorphism $\phi : \sigma^* \hat{N} \xrightarrow{\sim} \hat{N}$ of $Q_{v,L}$ -modules.

Remark 3.2. Note that so far in the literature [An2, Ha1, Ha2, Lau] it is always assumed that A_v has residue field \mathbb{F}_q , the fixed field of σ on L . So in particular $A_{v,L}$ is an integral domain and $Q_{v,L}$ is a field. For applications to pure Anderson motives this is not a problem since we may reduce to this case by Propositions 3.5 and 3.8 below.

Definition 3.3. A local shtuka $\underline{\hat{M}} = (\hat{M}, \phi)$ is called étale if ϕ is an isomorphism. The Tate module of an étale local σ -shtuka $\underline{\hat{M}}$ at v is the $G := \text{Gal}(L^{\text{sep}}/L)$ -module of ϕ -invariants

$$T_v \underline{\hat{M}} := (\underline{\hat{M}} \otimes_{A_{v,L}} A_{v,L^{\text{sep}}})^\phi.$$

The rational Tate module of $\underline{\hat{M}}$ is the G -module

$$V_v \underline{\hat{M}} := T_v \underline{\hat{M}} \otimes_{A_v} Q_v.$$

It follows from [TW, Proposition 6.1] that $T_v \underline{\hat{M}}$ is a free A_v -module of the same rank than $\underline{\hat{M}}$ and that the natural morphism

$$T_v \underline{\hat{M}} \otimes_{A_v} A_{v,L^{\text{sep}}} \xrightarrow{\sim} \underline{\hat{M}} \otimes_{A_{v,L}} A_{v,L^{\text{sep}}}$$

is a G - and ϕ -equivariant isomorphism of $A_{v,L^{\text{sep}}}$ -modules, where on the left module G acts on both factors and ϕ is $\text{id} \otimes \sigma^*$. Since $(L^{\text{sep}})^G = L$ we obtain:

Proposition 3.4. Let $\underline{\hat{M}}$ and $\underline{\hat{M}'}$ be étale local σ -shtuka at v over L . Then

1. $\underline{\hat{M}} = (T_v \underline{\hat{M}} \otimes_{A_v} A_{v,L^{\text{sep}}})^G$, the Galois invariants,
2. $\text{Hom}_{A_{v,L}[\phi]}(\underline{\hat{M}}, \underline{\hat{M}'}) \xrightarrow{\sim} \text{Hom}_{A_v[G]}(T_v \underline{\hat{M}}, T_v \underline{\hat{M}'})$, $f \mapsto T_v f$ is an isomorphism.

In particular the Tate module functor yields an equivalence of the category of étale local shtuka at v over L with the category of $A_v[G]$ -modules, which are finite free over A_v . \square

Proof. 1 and 2 are immediate. Hence clearly the Tate module functor is fully faithful. That it is an equivalence follows analogously to [Kat, Proposition 4.1.1]. \square

If the residue field \mathbb{F}_v of v is larger than \mathbb{F}_q one has to be a bit careful with local (iso-)shtuka since $Q_{v,L}$ is then in general not a field. Namely let $\#\mathbb{F}_v = q^n$ and let $\mathbb{F}_{q^f} := \{\alpha \in L : \alpha^{q^n} = \alpha\}$ be the “intersection” of \mathbb{F}_v with L . Then

$$\mathbb{F}_v \otimes_{\mathbb{F}_q} L = \prod_{\text{Gal}(\mathbb{F}_{q^f}/\mathbb{F}_q)} \mathbb{F}_v \otimes_{\mathbb{F}_{q^f}} L = \prod_{i \in \mathbb{Z}/f\mathbb{Z}} \mathbb{F}_v \otimes_{\mathbb{F}_q} L / (b \otimes 1 - 1 \otimes b^{q^i} : b \in \mathbb{F}_{q^f})$$

and σ^* transports the i -th factor to the $(i+1)$ -th factor. (Of course, the indexing of the factors depends on a choice of embeddings $\mathbb{F}_{q^f} \subset \mathbb{F}_v$ and $\mathbb{F}_{q^f} \subset L$.) Denote by \mathfrak{a}_i the ideal of $A_{v,L}$ (or $Q_{v,L}$) generated by $\{b \otimes 1 - 1 \otimes b^{q^i} : b \in \mathbb{F}_{q^f}\}$. Then

$$A_{v,L} = \prod_{\text{Gal}(\mathbb{F}_{q^f}/\mathbb{F}_q)} A_v \widehat{\otimes}_{\mathbb{F}_{q^f}} L = \prod_{i \in \mathbb{Z}/f\mathbb{Z}} A_{v,L} / \mathfrak{a}_i$$

and similarly for Q_v . Note that the factors in this decomposition and the ideals \mathfrak{a}_i correspond precisely to the places v_i of $C_{\mathbb{F}_{q^f}}$ lying above v .

Proposition 3.5. Fix an i . The reduction modulo \mathfrak{a}_i induces equivalences of categories

1. $(\hat{N}, \phi) \mapsto (\hat{N}/\mathfrak{a}_i \hat{N}, \phi^f : (\sigma^*)^f \hat{N}/\mathfrak{a}_i \hat{N} \rightarrow \hat{N}/\mathfrak{a}_i \hat{N})$

between local σ -isoshtuka at v over L and local σ^f -isoshtuka at v_i over L of the same rank.

$$2. (\hat{M}, \phi) \longmapsto (\hat{M}/\mathfrak{a}_i \hat{M}, \phi^f : (\sigma^*)^f \hat{M}/\mathfrak{a}_i \hat{M} \rightarrow \hat{M}/\mathfrak{a}_i \hat{M})$$

between étale local σ -shtuka at v over L and étale local σ^f -shtuka at v_i over L preserving Tate modules

$$T_v(\hat{M}, \phi) \xrightarrow{\sim} T_{v_i}(\hat{M}/\mathfrak{a}_i \hat{M}, \phi^f).$$

Proof. Since $\sigma^* \mathfrak{a}_i = \mathfrak{a}_{i+1}$ the isomorphism ϕ yields isomorphisms $\sigma^*(\hat{N}/\mathfrak{a}_i \hat{N}) \rightarrow \hat{N}/\mathfrak{a}_{i+1} \hat{N}$ and similarly for \hat{M} . These allow to reconstruct the other factors from $(\hat{N}/\mathfrak{a}_i \hat{N}, \phi^f)$, and likewise for \hat{M} . The isomorphism between the Tate modules follows from the observation that an element $(x_j)_{j \in \mathbb{Z}/f\mathbb{Z}}$ is ϕ -invariant if and only if $x_{j+1} = \phi(\sigma^* x_j)$ for all j and $x_i = \phi^f((\sigma^*)^f x_i)$. \square

Remark. The advantage of the étale local σ^f -shtuka at v_i is that it is a free module over the integral domain $A_{v,L}/\mathfrak{a}_i = A_v \widehat{\otimes}_{\mathbb{F}_{q^f}} L$, and similarly for local σ^f -isoshtuka. So the results from [An2, Ha1, Ha2, Lau] apply.

Now let $\underline{\mathcal{F}}$ be an abelian τ -sheaf and $v \in C$ an arbitrary place of Q . We define the *local σ -isoshtuka of $\underline{\mathcal{F}}$ at v* as

$$\underline{N}_v(\underline{\mathcal{F}}) := \left(\mathcal{F}_0 \otimes_{\mathcal{O}_{C_L}} Q_{v,L}, \Pi_0^{-1} \circ \tau_0 \right).$$

If $v \neq \infty$ we define the *local σ -shtuka of $\underline{\mathcal{F}}$ at v* as

$$\underline{M}_v(\underline{\mathcal{F}}) := \left(\mathcal{F}_0 \otimes_{\mathcal{O}_{C_L}} A_{v,L}, \Pi_0^{-1} \circ \tau_0 \right).$$

Likewise if \underline{M} is a pure Anderson motive over L and $v \in \text{Spec } A$ we define the *local σ -(iso)-shtuka of \underline{M} at v* as

$$\underline{M}_v(\underline{M}) := \underline{M} \otimes_{A_L} A_{v,L} \quad \text{respectively} \quad \underline{N}_v(\underline{M}) := \underline{M} \otimes_{A_L} Q_{v,L}.$$

These local (iso)-shtuka all have rank r . The local shtuka are étale if $v \neq \varepsilon$. Note that $\underline{N}_\infty(\underline{\mathcal{F}})$ does not contain a local σ -shtuka if $\varepsilon \neq \infty$, since then it is isoclinic of slope $-\text{wt}(\underline{\mathcal{F}}) < 0$.

However, if $v = \infty$ the periodicity condition allows to define a different local (iso)-shtuka at ∞ which is of slope ≥ 0 . Namely, choose a uniformizer z on C at ∞ and set $\hat{M}_i := \mathcal{F}_i \otimes_{\mathcal{O}_{C_L}} A_{\infty,L}$. Recall the integers k, l from Definition 1.5/2 and set $\tilde{\Pi} := \Pi_{l-1} \circ \cdots \circ \Pi_0$. We define the *big local σ -shtuka of $\underline{\mathcal{F}}$ at ∞* as

$$\tilde{\underline{M}}_\infty(\underline{\mathcal{F}}) := \hat{M}_0 \oplus \cdots \oplus \hat{M}_{l-1} \quad \text{with} \quad \phi := \begin{pmatrix} 0 & \cdots & 0 & \tilde{\Pi}^{-1} \circ z^k \tau_{l-1} \\ \tau_0 & \cdots & & 0 \\ 0 & \cdots & & \vdots \\ \vdots & \cdots & & 0 \\ 0 & \cdots & 0 & \tau_{l-2} & 0 \end{pmatrix} \quad (3.1)$$

We also define the *big local σ -isoshtuka of $\underline{\mathcal{F}}$ at ∞* as

$$\tilde{\underline{N}}_\infty(\underline{\mathcal{F}}) := \tilde{\underline{M}}_\infty(\underline{\mathcal{F}}) \otimes_{A_{\infty,L}} Q_{\infty,L}.$$

Both have rank rl and depend on the choice of k, l and z . If $\varepsilon \neq \infty$ then $\tilde{\underline{M}}_\infty(\underline{\mathcal{F}})$ is étale. Note that $\tilde{\underline{M}}_\infty(\underline{\mathcal{F}})$ and $\tilde{\underline{N}}_\infty(\underline{\mathcal{F}})$ were used in [Ha1] to construct the uniformization at ∞ of the moduli spaces of abelian τ -sheaves.

The big local (iso)-shtuka at ∞ , $\tilde{\underline{M}}_\infty(\underline{\mathcal{F}})$ and $\tilde{\underline{N}}_\infty(\underline{\mathcal{F}})$ are always equipped with the endomorphisms

$$\Pi := \begin{pmatrix} 0 & \cdots & 0 & \tilde{\Pi}^{-1} \circ z^k \Pi_{l-1} \\ \Pi_0 & \cdots & & 0 \\ 0 & \cdots & & \vdots \\ \vdots & \cdots & & 0 \\ 0 & \cdots & 0 & \Pi_{l-2} & 0 \end{pmatrix}, \quad \Lambda(\lambda) := \begin{pmatrix} \lambda \cdot \text{id}_{M_0} & & & \\ & \lambda^q \cdot \text{id}_{M_1} & & \\ & & \ddots & \\ & & & \lambda^{q^{l-1}} \cdot \text{id}_{M_{l-1}} \end{pmatrix} \quad (3.2)$$

for all $\lambda \in \mathbb{F}_{q^l} \cap L$. They satisfy the relations $\Pi^l = z^k$ and $\Pi \circ \Lambda(\lambda^q) = \Lambda(\lambda) \circ \Pi$. We let Δ_∞ be “the” central division algebra over Q_∞ of rank l with Hasse invariant $-\frac{k}{l}$, or explicitly

$$\Delta_\infty := \mathbb{F}_{q^l}((z))[\Pi] / (\Pi^l - z^k, \lambda z - z\lambda, \Pi\lambda^q - \lambda\Pi \text{ for all } \lambda \in \mathbb{F}_{q^l}). \quad (3.3)$$

If $\mathbb{F}_{q^l} \subset L$ we identify Δ_∞ with a subalgebra of $\text{End}_{Q_\infty, L[\phi]}(\tilde{N}_\infty(\mathcal{F}))$ by mapping $\lambda \in \mathbb{F}_{q^l} \subset \Delta_\infty$ to $\Lambda(\lambda)$.

The following two results were proved in [BH1, Theorems 8.6 and 8.7].

Theorem 3.6. *Let \mathcal{F} and \mathcal{F}' be abelian τ -sheaves of the same weight over a finite field L and let v be an arbitrary place of Q .*

1. *Then there is a canonical isomorphism of Q_v -vector spaces*

$$\text{QHom}(\mathcal{F}, \mathcal{F}') \otimes_Q Q_v \xrightarrow{\sim} \text{Hom}_{Q_{v, L[\phi]}}(\underline{N}_v(\mathcal{F}), \underline{N}_v(\mathcal{F}')).$$

2. *If $v = \infty$ choose an l which satisfies 1.5/2 for both \mathcal{F} and \mathcal{F}' and assume $\mathbb{F}_{q^l} \subset L$. Then there is a canonical isomorphism of Q_∞ -vector spaces*

$$\text{QHom}(\mathcal{F}, \mathcal{F}') \otimes_Q Q_\infty \xrightarrow{\sim} \text{Hom}_{\Delta_\infty \hat{\otimes}_{\mathbb{F}_q} L[\phi]}(\tilde{N}_\infty(\mathcal{F}), \tilde{N}_\infty(\mathcal{F}')).$$

Theorem 3.7. *Let \underline{M} and \underline{M}' be pure Anderson motives over a finite field L and let $v \in \text{Spec } A$ be an arbitrary maximal ideal. Then*

$$\text{Hom}(\underline{M}, \underline{M}') \otimes_A A_v \xrightarrow{\sim} \text{Hom}_{A_{v, L[\phi]}}(\underline{M}_v(\underline{M}), \underline{M}_v(\underline{M}')).$$

Let now the characteristic be finite and $v = \varepsilon$ be the characteristic point. Consider a pure Anderson motive \underline{M} of characteristic c , its local σ -shtuka $\underline{M}_\varepsilon(\underline{M}) = (\hat{M}, \phi)$ at ε and the decomposition of the later described before Proposition 3.5

$$\underline{M}_\varepsilon(\underline{M}) = \prod_{i \in \mathbb{Z}/f\mathbb{Z}} \underline{M}_\varepsilon(\underline{M}) / \mathfrak{a}_i \underline{M}_\varepsilon(\underline{M}).$$

From the morphism $c : \text{Spec } L \rightarrow \text{Spec } \mathbb{F}_\varepsilon \subset C$ we see that $\mathbb{F}_\varepsilon \subset L$, $f = [\mathbb{F}_\varepsilon : \mathbb{F}_q]$ and that there is a distinguished place v_0 of $C_{\mathbb{F}_\varepsilon}$ above $v = \varepsilon \neq \infty$, namely the image of $c \times c : \text{Spec } L \rightarrow C \times \text{Spec } \mathbb{F}_\varepsilon$. Then ϕ has no cokernel on $\underline{M}_\varepsilon(\underline{M}) / \mathfrak{a}_i \underline{M}_\varepsilon(\underline{M})$ for $i \neq 0$ and the reasoning of Proposition 3.5 yields

Proposition 3.8. *The reduction modulo \mathfrak{a}_0 induces an equivalence of categories*

$$\underline{M}_\varepsilon(\underline{M}) \longmapsto (\underline{M}_\varepsilon(\underline{M}) / \mathfrak{a}_0 \underline{M}_\varepsilon(\underline{M}), \phi^f)$$

between the local σ -shtuka at ε associated with pure Anderson motives of characteristic c and the local σ^f -shtuka at v_0 associated with pure Anderson motives of characteristic c . The same is true for abelian τ -sheaves. \square

Remark. Now the fixed field of σ^f on L equals \mathbb{F}_ε , the residue field of A_ε . Also $\underline{M}_\varepsilon(\mathcal{F}) / \mathfrak{a}_0 \underline{M}_\varepsilon(\mathcal{F})$ is a module over the integral domain $A_\varepsilon \hat{\otimes}_{\mathbb{F}_\varepsilon} L$. So again [An2, Ha1, Ha2, Lau] apply to $(\underline{M}_\varepsilon(\mathcal{F}) / \mathfrak{a}_0 \underline{M}_\varepsilon(\mathcal{F}), \phi^f)$.

Proposition 3.9. *Let \underline{M} be a pure Anderson motive over L and let $\hat{\underline{M}}'_\varepsilon$ be a local σ^f -subshtuka of $\underline{M}_\varepsilon(\underline{M}) / \mathfrak{a}_0 \underline{M}_\varepsilon(\underline{M})$ of the same rank. Then there is a pure Anderson motive \underline{M}' and an isogeny $f : \underline{M}' \rightarrow \underline{M}$ with $\underline{M}_\varepsilon(f)(\underline{M}_\varepsilon(\underline{M}') / \mathfrak{a}_0 \underline{M}_\varepsilon(\underline{M}')) = \hat{\underline{M}}'_\varepsilon$. The same is true for abelian τ -sheaves.*

Proof. Extend \hat{M}'_ε to the local σ -subshutuka $\bigoplus_{i \in \mathbb{Z}/f\mathbb{Z}} \phi^i((\sigma^*)^i \hat{M}'_\varepsilon)$ of $\underline{M}_\varepsilon(\underline{M})$ and consider

$$\underline{K} := \underline{M}_\varepsilon(\underline{M}) / \bigoplus_{i \in \mathbb{Z}/f\mathbb{Z}} \phi^i((\sigma^*)^i \hat{M}'_\varepsilon).$$

The induced morphism $\phi_K : \sigma^*K \rightarrow K$ has its kernel and cokernel supported on the graph of c . Set $\underline{M}' = (M', \tau') := (\ker(\underline{M} \rightarrow \underline{K}), \tau|_{M'})$. Then \underline{M}' is a pure Anderson motive with the required properties by [BH1, Proposition 1.6]. \square

There is a corresponding result at the places $v \neq \varepsilon$ which is stated in Proposition 4.4.

4 Tate Modules

Definition 4.1. *If $\underline{\mathcal{F}}$ is an abelian τ -sheaf over L , respectively \underline{M} a pure Anderson motive over L and $v \in C$ (respectively $v \in \text{Spec } A$) is a place of Q different from the characteristic point ε , we define*

$$\begin{aligned} T_v \underline{\mathcal{F}} &:= T_v(\underline{M}_v(\underline{\mathcal{F}})) & \text{and} & & V_v \underline{\mathcal{F}} &:= V_v(\underline{M}_v(\underline{\mathcal{F}})) & \text{for } v \neq \infty, \\ T_\infty \underline{\mathcal{F}} &:= T_\infty(\tilde{\underline{M}}_\infty(\underline{\mathcal{F}})) & \text{and} & & V_\infty \underline{\mathcal{F}} &:= V_\infty(\tilde{\underline{M}}_\infty(\underline{\mathcal{F}})) & \text{for } v = \infty \neq \varepsilon, \\ \text{respectively } T_v \underline{M} &:= T_v(\underline{M}_v(\underline{M})) & \text{and} & & V_v \underline{M} &:= V_v(\underline{M}_v(\underline{M})). \end{aligned}$$

We call $T_v \underline{\mathcal{F}}$ (respectively $V_v \underline{\mathcal{F}}$) the (rational) v -adic Tate module of $\underline{\mathcal{F}}$. If $v = \infty$ they both depend on the choice of k, l , and z ; see page 10.

By [TW, Proposition 6.1], $T_v \underline{\mathcal{F}}$ (and $V_v \underline{\mathcal{F}}$) are free A_v -modules (respectively Q_v -vector spaces) of rank r for $v \neq \infty$ and rl for $v = \infty$, which carry a continuous $G = \text{Gal}(L^{\text{sep}}/L)$ -action.

Also the Tate modules $T_\infty \underline{\mathcal{F}}$ and $V_\infty \underline{\mathcal{F}}$ are always equipped with the endomorphisms Π and $\Lambda(\lambda)$ for $\lambda \in \mathbb{F}_{q^l} \cap L$ from (3.2). And if $\mathbb{F}_{q^l} \subset L$ we identify the algebra Δ_∞ from (3.3) with a subalgebra of $\text{End}_{Q_\infty}(V_\infty \underline{\mathcal{F}})$ by mapping $\lambda \in \mathbb{F}_{q^l}$ to $\Lambda(\lambda)$.

Remark. Our functor T_v is covariant. In the literature usually the A_v -dual of our $T_v \underline{M}$ is called the v -adic Tate module of \underline{M} . With that convention the Tate module functor is contravariant on Anderson motives but covariant on Drinfeld modules and Anderson's abelian t -modules [An1] (which both give rise to Anderson motives). Similarly the classical Tate module functor on abelian varieties is covariant. We chose our non-standard convention here solely to avoid perpetual dualizations. This agrees also with the remark after Definition 2.6 that abelian τ -sheaves behave dually to abelian varieties.

The following analogues of the Tate conjecture for abelian varieties are due to Taguchi [Tag] and Tamagawa [Tam, §2].

Theorem 4.2. *Let \underline{M} and \underline{M}' be pure Anderson motives over a finitely generated field L and let $G := \text{Gal}(L^{\text{sep}}/L)$. Let $\varepsilon \neq v \in \text{Spec } A$ be a maximal ideal. Then the Tate conjecture holds:*

$$\text{Hom}(\underline{M}, \underline{M}') \otimes_A A_v \cong \text{Hom}_{A_v[G]}(T_v \underline{M}, T_v \underline{M}'). \quad \square$$

Theorem 4.3. *([BH1, Theorem 9.9]) Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}'}$ be abelian τ -sheaves over a finitely generated field L and let $G := \text{Gal}(L^{\text{sep}}/L)$. Let $v \in C$ be a place different from the characteristic point ε .*

1. *If $v \neq \infty$ assume $\varepsilon \neq \infty$ or $\text{wt}(\underline{\mathcal{F}}) = \text{wt}(\underline{\mathcal{F}'})$. Then*

$$\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}'}) \otimes_Q Q_v \cong \text{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}, V_v \underline{\mathcal{F}'}).$$

2. *If $v = \infty$ choose an integer l which satisfies 1.5/2 for both $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}'}$ and assume $\mathbb{F}_{q^l} \subset L$. Then*

$$\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}'}) \otimes_Q Q_\infty \cong \text{Hom}_{\Delta_\infty[G]}(V_\infty \underline{\mathcal{F}}, V_\infty \underline{\mathcal{F}'}).$$

As expected, there is the following relation between Tate modules and isogenies.

Proposition 4.4. (*[BH1, Proposition 9.11]*)

1. Let $f : \underline{M}' \rightarrow \underline{M}$ be an isogeny between pure Anderson motives then $T_v f(T_v \underline{M}')$ is a G -stable lattice in $V_v \underline{M}$ contained in $T_v \underline{M}$.
2. Conversely if \underline{M} is a pure Anderson motive and Λ_v is a G -stable lattice in $V_v \underline{M}$ contained in $T_v \underline{M}$, then there exists a pure Anderson motive \underline{M}' and a separable isogeny $f : \underline{M}' \rightarrow \underline{M}$ with $T_v f(T_v \underline{M}') = \Lambda_v$.

Proposition 4.5. Let $\underline{\mathcal{F}}'$ be an abelian factor τ -sheaf of $\underline{\mathcal{F}}$. Then $V_v \underline{\mathcal{F}}'$ is a G -factor space of $V_v \underline{\mathcal{F}}$. The same holds if \underline{M}' is a factor motive of a pure Anderson motive \underline{M} .

Proof. Let $f \in \text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ be surjective and let $\hat{\underline{M}}$ and $\hat{\underline{M}}'$ be the (big, if $v = \infty$) local σ -shtuka of $\underline{\mathcal{F}}$, respectively $\underline{\mathcal{F}}'$, at v . Then the induced morphism $\underline{M}_v(f) \in \text{Hom}(\hat{\underline{M}}, \hat{\underline{M}}')$ is surjective and $\hat{\underline{M}}'' := \ker \underline{M}_v(f)$ is also a local σ -shtuka at v . We get an exact sequence of local σ -shtuka which we tensor with $A_{v, L^{\text{sep}}}$ yielding

$$0 \longrightarrow \hat{\underline{M}}'' \otimes_{A_{v, L}} A_{v, L^{\text{sep}}} \longrightarrow \hat{\underline{M}} \otimes_{A_{v, L}} A_{v, L^{\text{sep}}} \xrightarrow{\underline{M}_v(f)} \hat{\underline{M}}' \otimes_{A_{v, L}} A_{v, L^{\text{sep}}} \longrightarrow 0.$$

The Tate module functor is left exact, because considering the morphism of $A_{v, L^{\text{sep}}}$ -modules

$$1 - \tau : \hat{\underline{M}} \otimes_{A_{v, L}} A_{v, L^{\text{sep}}} \longrightarrow \hat{\underline{M}} \otimes_{A_{v, L}} A_{v, L^{\text{sep}}}$$

we have by definition $T_v \hat{\underline{M}} = \ker(1 - \tau)$, and the desired left exactness follows from the snake lemma. After tensoring with $\otimes_{A_v} Q_v$ we get

$$0 \longrightarrow V_v \hat{\underline{M}}'' \longrightarrow V_v \hat{\underline{M}} \xrightarrow{V_v f} V_v \hat{\underline{M}}'.$$

Counting the dimensions of these Q_v -vector spaces, we finally also get right exactness, as desired. \square

5 The Frobenius Endomorphism

Suppose that the characteristic is finite, that is, the characteristic point ε is a closed point of C with finite residue field \mathbb{F}_ε , and the map $c : \text{Spec } L \rightarrow C$ factors through the finite field $\varepsilon = \text{Spec } \mathbb{F}_\varepsilon$.

Definition 5.1 (*s -Frobenius on abelian τ -sheaves*). Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf with finite characteristic point $\varepsilon = \text{Spec } \mathbb{F}_\varepsilon$ and let $s = q^e$ be a power of the cardinality of \mathbb{F}_ε . We define the s -Frobenius on $\underline{\mathcal{F}}$ by

$$\pi := (\pi_i) : (\sigma^*)^e \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}[e], \quad \pi_i := \tau_{i+e-1} \circ \dots \circ (\sigma^*)^{e-1} \tau_i : (\sigma^*)^e \mathcal{F}_i \rightarrow \mathcal{F}_{i+e}.$$

Clearly π is an isogeny. Observe that $\mathbb{F}_\varepsilon \subset \mathbb{F}_s$ implies that $(\sigma^*)^e \underline{\mathcal{F}}$ has the same characteristic as $\underline{\mathcal{F}}$.

Similarly if $\varepsilon \in \text{Spec } A$ is a closed point we define

Definition 5.2 (*s -Frobenius on pure Anderson motives*). Let \underline{M} be a pure Anderson motive with finite characteristic point $\varepsilon = \text{Spec } \mathbb{F}_\varepsilon$ and let $s = q^e$ be a power of the cardinality of \mathbb{F}_ε . We define the s -Frobenius isogeny on \underline{M} by

$$\pi := \tau \circ \dots \circ (\sigma^*)^{e-1} \tau : (\sigma^*)^e \underline{M} \rightarrow \underline{M}.$$

Remark 5.3. Classically for (abelian) varieties X over a field K of characteristic p one defines the Frobenius morphism $X \rightarrow \phi^* X$ where ϕ is the p -Frobenius on K . There p equals the cardinality of the “characteristic field” $\text{im}(\mathbb{Z} \rightarrow K) = \mathbb{F}_p$. In view of the dual behavior of abelian τ -sheaves and pure Anderson motives our definition is a perfect analogue since here we consider the s -Frobenius for s being the cardinality of (a power of) the “characteristic field” $\text{im}(c^* : A \rightarrow L) = \mathbb{F}_\varepsilon$.

Now we suppose $L = \mathbb{F}_s$ to be a finite field with $s = q^e$ ($e \in \mathbb{N}$). Let $\overline{\mathbb{F}}_s$ denote a fixed algebraic closure of \mathbb{F}_s and set $G = \text{Gal}(\overline{\mathbb{F}}_s/\mathbb{F}_s)$. It is topologically generated by $\text{Frob}_s : x \mapsto x^s$. The following results for the Frobenius endomorphism of τ -modules can be found in Taguchi and Wan [TW, §6].

Proposition 5.4. *Let \underline{M} be a pure Anderson motive over \mathbb{F}_s of rank r and let $\varepsilon \neq v \in \text{Spec } A$ be a maximal ideal.*

1. *The generator Frob_s of G acts on $T_v \underline{M}$ like $(T_v \pi)^{-1}$.*
2. *Let $\Psi : A_v[G] \rightarrow \text{End}_{A_v}(T_v \underline{M})$ denote the continuous morphism of A_v -modules which is induced by the action of G on $T_v \underline{M}$. Then $\text{im } \Psi = A_v[T_v \pi]$.*

Proof. 1 was proved in [TW, Ch. 6] and 2 follows from the continuity of Ψ . □

Remark. The inversion of $T_v \pi$ in the first statement results from the dual definition of our Tate module.

Proposition 5.5. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over $L = \mathbb{F}_s$ with $s = q^e$ and let π be its s -Frobenius. Then $(\sigma^*)^e \underline{\mathcal{F}} = \underline{\mathcal{F}}$. Let $v \in C$ be a place different from ∞ and from the characteristic point ε .*

1. *The s -Frobenius π can be considered as a quasi-isogeny of $\underline{\mathcal{F}}$.*
2. *The generator Frob_s of G acts on $T_v \underline{\mathcal{F}}$ like $(T_v \pi)^{-1}$.*
3. *The image of the continuous morphism of Q_v -vector spaces $Q_v[G] \rightarrow \text{End}_{Q_v}(V_v \underline{\mathcal{F}})$ is $Q_v[V_v \pi]$.*
4. *$\underline{M}(\pi)$ coincides with the s -Frobenius on the pure Anderson motive $\underline{M}(\underline{\mathcal{F}})$ from definition 5.2.*

Proof. 1. Due to the periodicity condition, we have $\underline{\mathcal{F}}[e] \subset \underline{\mathcal{F}}(nk \cdot \infty)$ for a sufficiently large $n \in \mathbb{N}$, since $\mathcal{F}_{i+e} \subset \mathcal{F}_{i+nl} = \mathcal{F}_i(nk \cdot \infty)$ for $e \leq nl$. Thus $\pi \in \text{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}(nk \cdot \infty))$, and therefore $\pi \in \text{QEnd}(\underline{\mathcal{F}})$. By 2.1, we have $\pi \in \text{QIsog}(\underline{\mathcal{F}})$.

2 and 3 again follow from [TW, Ch. 6] and the continuity of Ψ ; see [BH2, Proposition 2.29] for more details.

4 follows from the definition of π and the commutation of the H 's and the τ 's. □

6 The Poincaré-Weil Theorem

In this section we study the analogue for pure Anderson motives and abelian τ -sheaves of the Poincaré-Weil theorem. Originally, this theorem states that every abelian variety is semisimple, that is, isogenous to a product of simple abelian varieties, see [Lan, Corollary of Theorem II.1/6]. Unfortunately, we cannot expect a full analogue of this statement for abelian τ -sheaves or pure Anderson motives as our next example illustrates. On the positive side we show that every abelian τ -sheaf or pure Anderson motive over a finite field becomes semisimple after a finite base field extension.

Example 6.1. Let $C = \mathbb{P}_{\mathbb{F}_q}^1$, $C \setminus \{\infty\} = \text{Spec } \mathbb{F}_q[t]$ and $\zeta := c^*(1/t) \in \mathbb{F}_q^\times$. We construct an abelian τ -sheaf $\underline{\mathcal{F}}$ over $L = \mathbb{F}_q$ with $r = d = 2$ which is not semisimple. Let

$$\Delta = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \cdot t$$

with $\alpha, \beta, \gamma, \delta \in \mathbb{F}_q$. To obtain characteristic c we need $\det \Delta = (1 - \zeta t)^2$, and thus we require the conditions $\alpha + \delta = -2\zeta$ and $\alpha\delta - \beta\gamma = \zeta^2$. We set $\mathcal{F}_i := \mathcal{O}_{C_L}(i \cdot \infty)^{\oplus 2}$, we let H_i be the natural inclusion, and we let $\tau_i := \Delta$. Then $\underline{\mathcal{F}}$ is an abelian τ -sheaf with $r = d = 2$ and $k = l = 1$. The associated pure Anderson motive is $\underline{M} = (L[t]^{\oplus 2}, \Delta)$.

We see that $\underline{\mathcal{F}}$ is not simple. If $\Delta = \begin{pmatrix} 1-\zeta t & 0 \\ 0 & 1-\zeta t \end{pmatrix}$ then $\underline{\mathcal{F}}$ is semisimple as a direct sum of two simple abelian τ -sheaves. Otherwise, if $\Delta \neq \begin{pmatrix} 1-\zeta t & 0 \\ 0 & 1-\zeta t \end{pmatrix}$ which is the case for example if $\beta \neq 0$, consider

$$\tilde{\Delta} := \begin{pmatrix} \beta & 0 \\ \delta + \zeta & 1 \end{pmatrix}^{-1} \cdot \Delta \cdot \sigma^* \begin{pmatrix} \beta & 0 \\ \delta + \zeta & 1 \end{pmatrix} = \begin{pmatrix} 1-\zeta t & t \\ 0 & 1-\zeta t \end{pmatrix}$$

and the abelian τ -sheaf with $\tilde{\mathcal{F}}_i = \mathcal{O}_{C_L}(i \cdot \infty)^{\oplus 2}$ and $\tilde{\tau}_i = \tilde{\Delta}$ which is isomorphic to $\underline{\mathcal{F}}$. There is an exact sequence

$$0 \longrightarrow \underline{\mathcal{F}}' \xrightarrow[\tau' = 1 - \zeta t]{\varphi} \tilde{\mathcal{F}} \xrightarrow[\tilde{\tau}]{\psi} \underline{\mathcal{F}}'' \longrightarrow 0$$

with $\varphi : 1 \mapsto \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\psi : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto y$ where $\underline{\mathcal{F}}' = \underline{\mathcal{F}}''$ is the abelian τ -sheaf with $\mathcal{F}'_i = \mathcal{O}_{C_L}(i \cdot \infty)$ and $\tau'_i = 1 - \zeta t$. If $\tilde{\mathcal{F}}$ were semisimple, then there would be a quasi-morphism $\omega : \underline{\mathcal{F}}'' \rightarrow \tilde{\mathcal{F}}$ with $\psi \circ \omega = \text{id}_{\underline{\mathcal{F}}''}$, hence $\omega : y \mapsto \begin{pmatrix} e \\ 1 \end{pmatrix} \cdot y$ for some $e \in \mathbb{F}_q(t)$. Thus, a necessary condition for the semisimplicity of $\underline{\mathcal{F}}$ is

$$(1 - \zeta t) \cdot \sigma^*(y) \cdot \begin{pmatrix} e \\ 1 \end{pmatrix} = \begin{pmatrix} 1 - \zeta t & t \\ 0 & 1 - \zeta t \end{pmatrix} \cdot \begin{pmatrix} \sigma^*(e) \\ 1 \end{pmatrix} \cdot \sigma^*(y)$$

which is equivalent to the condition

$$e - \sigma^*(e) = \frac{t}{1 - \zeta t}.$$

But this cannot be true since $e - \sigma^*(e) = 0$, thus $\underline{\mathcal{F}}$ is *not* semisimple. However, this last formula is satisfied if $e = \lambda \cdot \frac{t}{1 - \zeta t}$ for $\lambda \in \mathbb{F}_{q^q}$ with $\lambda^q - \lambda = -1$. That means that after field extension $\mathbb{F}_q(\lambda) / \mathbb{F}_q$ we get $\underline{\mathcal{F}} \cong \underline{\mathcal{F}}'^{\oplus 2}$ and we have $\text{QEnd}(\underline{\mathcal{F}}) = M_2(\text{QEnd}(\underline{\mathcal{F}}')) = M_2(Q)$. Note that this phenomenon generally appears, and we will state and prove it in Theorem 6.15.

From now on we fix a place $v \in \text{Spec } A$ which is different from the characteristic point ε of c . For a morphism $f \in \text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ between two abelian τ -sheaves $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ we denote its image $V_v f \in \text{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}, V_v \underline{\mathcal{F}}')$ just by f_v . If $\underline{\mathcal{F}}$ is defined over \mathbb{F}_s this applies in particular to the s -Frobenius endomorphism π of $\underline{\mathcal{F}}$ (Definition 5.1).

Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over the finite field $L = \mathbb{F}_s$. We set

$$\begin{aligned} E &:= \text{QEnd}(\underline{\mathcal{F}}) \ni \pi & E_v &:= \text{End}_{Q_v[G]}(V_v \underline{\mathcal{F}}) \ni \pi_v \\ F &:= Q[\pi] \subset E & F_v &:= \text{im}(Q_v[G] \rightarrow \text{End}_{Q_v}(V_v \underline{\mathcal{F}})) \end{aligned} \quad (6.1)$$

with $Q_v[G] \rightarrow \text{End}_{Q_v}(V_v \underline{\mathcal{F}})$ induced by the action of G on $V_v \underline{\mathcal{F}}$. Clearly, we have $F \subset E$ and $F_v \subset E_v$ by Proposition 5.5/3. By [BH1, Proposition 9.4], we know that $\dim_Q E < \infty$. Thus π is algebraic over Q . We denote its minimal polynomial by $\mu_\pi \in Q[x]$, and the characteristic polynomial of the endomorphism π_v of $V_v \underline{\mathcal{F}}$ by $\chi_v \in Q_v[x]$. If $\varepsilon \neq \infty$, Theorem 1.3 shows that π is integral over A , $\mu_\pi \in A[x]$. The zeroes of π in $\text{Spec } A[\pi]$ all lie above ε because π is an isomorphism locally at all $v \neq \varepsilon$; compare with [BH1, Remark 5.5].

Due to the Tate conjecture, our situation can be represented by the following diagram where we want to fit the missing bottom right arrow with an isomorphism.

$$\begin{array}{ccccc} E & \longrightarrow & E \otimes_Q Q_v & \xrightarrow{\sim} & E_v \\ \uparrow & & \uparrow & & \uparrow \\ F & \longrightarrow & F \otimes_Q Q_v & \xrightarrow{\sim} & F_v \end{array}$$

Lemma 6.2. *The natural morphism between $F \otimes_Q Q_v$ and F_v is an isomorphism.*

Proof. Consider the isomorphism $\psi : E \otimes_Q Q_v \cong E_v \subset \text{End}_{Q_v}(V_v \underline{\mathcal{F}})$ and set $\varphi := \psi|_{F \otimes_Q Q_v}$. Then φ is injective and maps into F_v . Since $\text{im } \varphi = Q_v[\pi_v]$, the surjectivity follows from Proposition 5.5. \square

To evaluate the dimension of E we need the following notation.

Definition 6.3. *Let K be a field. Let $f, g \in K[x]$ be two polynomials and let*

$$f = \prod_{\substack{\mu \in K[x] \\ \text{irred.}}} \mu^{m(\mu)}, \quad g = \prod_{\substack{\mu \in K[x] \\ \text{irred.}}} \mu^{n(\mu)}$$

be their respective factorizations in powers of irreducible polynomials. Then we define the integer

$$r_K(f, g) := \prod_{\substack{\mu \in K[x] \\ \text{irred.}}} m(\mu) \cdot n(\mu) \cdot \deg \mu.$$

Remark. In contrast to characteristic zero, we have for $\text{char}(K) \neq 0$ in general different values of the integer r_K for different ground fields K . Namely, if $K \subset L$ then $r_K(f, g) \leq r_L(f, g)$ with equality if and only if all irreducible $\mu \in K[x]$ which are contained both in f and in g have no multiple factors in $L[x]$. This is satisfied for example if the greatest common divisor of f and g has only separable irreducible factors, or if L is separable over K . See 9.4 below for an example where $r_K(f, g) < r_L(f, g)$.

Before we discuss semisimplicity criteria in 6.8 – 6.16, let us compute the dimension of $\text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$.

Lemma 6.4. *Let v be a place of Q different from ε and ∞ . Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ be abelian τ -sheaves over \mathbb{F}_s and assume that π_v and π'_v are semisimple. Factor their characteristic polynomials $\chi_v = \mu_1^{m_1} \cdots \mu_n^{m_n}$ and $\chi'_v = \mu_1^{m'_1} \cdots \mu_n^{m'_n}$ with distinct monic irreducible polynomials $\mu_1, \dots, \mu_n \in Q_v[x]$ and $m_i, m'_i \in \mathbb{N}_0$. Then*

1. $\text{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}, V_v \underline{\mathcal{F}}') \cong \bigoplus_{i=1}^n M_{m'_i \times m_i}(Q_v[x]/(\mu_i))$ as Q_v -vector spaces,
2. $\text{End}_{Q_v[G]}(V_v \underline{\mathcal{F}}) \cong \bigoplus_{i=1}^n M_{m_i}(Q_v[x]/(\mu_i))$ as Q_v -algebras, and
3. $\dim_{Q_v} \text{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}, V_v \underline{\mathcal{F}}') = r_{Q_v}(\chi_v, \chi'_v)$.

Proof. Clearly 2 and 3 are consequences of 1 which we now prove. Since π_v and π'_v are semisimple, we have the following decomposition of $Q_v[G]$ -modules

$$V_v \underline{\mathcal{F}} \cong \bigoplus_{i=1}^n (Q_v[x]/(\mu_i))^{\oplus m_i}, \quad V_v \underline{\mathcal{F}}' \cong \bigoplus_{i=1}^n (Q_v[x]/(\mu_i))^{\oplus m'_i}$$

where $Q_v[x]/(\mu_i) =: K_i$ are fields. Obviously, we only have non-zero $Q_v[G]$ -morphisms $K_i \rightarrow K_j$ if $i = j$, since otherwise $\mu_i(\pi) \neq 0$ in K_j . Since π_v operates on $K_i^{\oplus m_i}$ as multiplication by the scalar x , the lemma follows. \square

Theorem 6.5. *Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ be abelian τ -sheaves of the same weight over \mathbb{F}_s and assume that the endomorphisms π_v and π'_v of $V_v \underline{\mathcal{F}}$ and $V_v \underline{\mathcal{F}}'$ are semisimple at a place $v \neq \varepsilon, \infty$ of Q . Let χ_v and χ'_v be their characteristic polynomials. Then*

$$\dim_Q \text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}') = r_{Q_v}(\chi_v, \chi'_v).$$

Proof. This follows from the lemma and the Tate conjecture, Theorem 4.3. \square

Corollary 6.6. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf of rank r over \mathbb{F}_s with Frobenius endomorphism π and let μ_π be the minimal polynomial of π . Assume that $F = Q[x]/(\mu_\pi)$ is a field and set $h := [F : Q] = \deg \mu_\pi$. Then*

1. $h|r$ and $\dim_Q \text{QEnd}(\underline{\mathcal{F}}) = \frac{r^2}{h}$ and $\dim_F \text{QEnd}(\underline{\mathcal{F}}) = \frac{r^2}{h^2}$.
2. For any place v of Q different from ε and ∞ we have $\text{QEnd}(\underline{\mathcal{F}}) \otimes_Q Q_v \cong M_{r/h}(F \otimes_Q Q_v)$ and $\chi_v = (\mu_\pi)^{r/h}$ independent of v .

Proof. Since F is a field, π_v is semisimple by 6.8 below. So general facts of linear algebra imply that $\mu_\pi = \mu_1 \cdots \mu_n$ with pairwise different irreducible monic polynomials $\mu_i \in Q_v[x]$ and $\chi_v = \mu_1^{m_1} \cdots \mu_n^{m_n}$ with $m_i \geq 1$. We set $K_i = Q_v[x]/(\mu_i)$ and use the notation from (6.1). By Lemma 6.4 the semisimple Q_v -algebra E_v decomposes $E_v \cong \bigoplus_{i=1}^n E_i$ into the simple constituents $E_i = M_{m_i}(K_i)$. By [Bou,

Théorème 5.3/1 and Proposition 5.4/12], $E_i = E_v \cdot e_i$ where e_i are the central idempotents with $K_i = F_v \cdot e_i$. Thus there are epimorphisms of K_i -vector spaces

$$\mathrm{QEnd}(\underline{\mathcal{F}}) \otimes_F K_i = E_v \otimes_{F_v} K_i \longrightarrow E_i.$$

This shows that $m_i^2 \leq \dim_F E$. So by Lemma 6.4

$$\begin{aligned} [F : Q] \cdot \dim_F E &= \dim_{Q_v} E_v = \sum_{i=1}^n m_i^2 \deg \mu_i \leq \\ &\leq \dim_F E \cdot \sum_{i=1}^n \deg \mu_i = \dim_F E \cdot \deg \mu_\pi = [F : Q] \cdot \dim_F E. \end{aligned}$$

Therefore $m_i^2 = \dim_F E$ for all i . Since $r = \deg \chi_v = \sum_i m_i \deg \mu_i = \sqrt{\dim_F E} \cdot [F : Q]$. We find $r = m_i h$ and $\dim_F E = \frac{r^2}{h^2}$, proving 1. For 2 we use that

$$E_v \cong \bigoplus_i M_{r/h}(Q_v[x]/(\mu_i)) = M_{r/h}\left(\bigoplus_i Q_v[x]/(\mu_i)\right) = M_{r/h}(Q_v[x]/(\mu_\pi)).$$

□

Next we investigate when π_v is semisimple.

Remark 6.7. Notice that the completion Q_v is separable over Q . Namely, in terms of [EGA, IV.7.8.1–3], we can state that $\mathcal{O}_{C,v}$ is an excellent ring. Thus the formal fibers of $\widehat{\mathcal{O}}_{C,v} \longrightarrow \mathcal{O}_{C,v}$ and therefore $Q_v = \widehat{\mathcal{O}}_{C,v} \otimes_{\mathcal{O}_{C,v}} Q \longrightarrow Q$ are geometrically regular. This means that $Q_v \otimes_Q K$ is regular for every finite field extension K over Q . Since "regular" implies "reduced", we conclude that Q_v is separable over Q .

Proposition 6.8. *In the notation of (6.1) the following statements are equivalent:*

1. π is semisimple.
2. F is semisimple.
3. $F \otimes_Q Q_v \cong F_v$ is semisimple.
4. π_v is semisimple.
5. $E \otimes_Q Q_v \cong E_v$ is semisimple.
6. E is semisimple.

Proof. 1. and 2. are equivalent by definition. So we show the equivalences from 2. to 6.

Let F be semisimple. Since Q_v is separable over Q , we conclude that $F \otimes_Q Q_v \cong Q_v[\pi_v]$ is semisimple by [Bou, Corollaire 7.6/4]. Hence π_v is semisimple by definition, and we showed in Lemma 6.4/2 that then $E_v \cong E \otimes_Q Q_v$ is semisimple. Again by [Bou, Corollaire 7.6/4] this implies that E is semisimple. Since $F \subset Z(E)$ is a finite dimensional Q -subalgebra of the center of E , we conclude by [Bou, Corollaire de Proposition 6.4/9] that F is semisimple, and our proof is complete. □

Remark. If more generally $\underline{\mathcal{F}}$ is defined over a finitely generated field, then one cannot consider π , π_v , nor F . Nevertheless 5 and 6 remain equivalent and are still implied by 3 due to the following well-known lemma. Namely E_v is the commutant of F_v in $\mathrm{End}_{Q_v}(V_v \underline{\mathcal{F}})$. We thank O. Gabber for mentioning this fact to us and we include its proof for lack of reference.

Lemma 6.9. *Let B be a central simple algebra of finite dimension over a field K and let F be a semisimple K -subalgebra of B . Then the commutant of F in B is semisimple.*

Proof. Let $F = \bigoplus_i F_i$ be the decomposition into simple constituents and let e_i be the corresponding central idempotents, that is, $F_i = F e_i$. Consider $B_i = e_i B e_i$ which is again central simple over K by [Bou, Corollaire 6.4/4], since if $I \subset B_i$ is a non-zero two sided ideal then BIB contains 1 and so I contains the unit e_i of B_i . By [Bou, Théorème 10.2/2] the commutant E_i of F_i in B_i is simple. Clearly the commutant E of F in B satisfies $E_i = e_i E e_i = E e_i$ and $E = \bigoplus_i E_i$ proving the lemma. □

Corollary 6.10. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over \mathbb{F}_s of rank r with semisimple Frobenius endomorphism π . Then the algebra $F = Q(\pi)$ is the center of the semisimple algebra $E = \mathrm{QEnd}(\underline{\mathcal{F}})$.*

Proof. Since F_v is semisimple, we know by [Bou, Proposition 5.1/1] that the F_v -module $V_v \underline{\mathcal{F}}$ is semisimple. The commutant of F_v in $\mathrm{End}_{Q_v}(V_v \underline{\mathcal{F}})$ is E_v by definition. Trivially $V_v \underline{\mathcal{F}}$ is of finite type over E_v . Thus, by the theorem of bicommutation [Bou, Corollaire 4.2/1], the commutant of E_v in $\mathrm{End}(V_v \underline{\mathcal{F}})$ is again F_v . We conclude $Z(E_v) = E_v \cap F_v = F_v$ and we have $F \otimes_Q Q_v = F_v = Z(E_v) = Z(E) \otimes_Q Q_v$ by [Bou, Corollaire de Proposition 1.2/3]. Considering the dimensions, we obtain $\dim_Q F = \dim_Q Z(E)$. Since $F \subset Z(E)$ and the dimensions are finite, we finish by $F = Z(E)$. \square

Theorem 6.11. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over a finite field L .*

1. *If $\mathrm{QEnd}(\underline{\mathcal{F}})$ is a division algebra over Q then $\underline{\mathcal{F}}$ is simple. If in addition $\varepsilon \neq \infty$ then both statements are equivalent.*
2. *If the characteristic point ε is different from ∞ then $\underline{\mathcal{F}}$ is semisimple if and only if $\mathrm{QEnd}(\underline{\mathcal{F}})$ is semisimple.*

Proof. 1. Let $\mathrm{QEnd}(\underline{\mathcal{F}}) = E$ be a division algebra and let $f \in \mathrm{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$ be the morphism onto a non-zero factor sheaf $\underline{\mathcal{F}}'$ of $\underline{\mathcal{F}}$. We show that f is an isomorphism. We know by 4.5 that $f_v \in \mathrm{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}, V_v \underline{\mathcal{F}}')$ is surjective. By the semisimplicity of E and Proposition 6.8, F_v is semisimple, and therefore $V_v \underline{\mathcal{F}}$ is a finitely generated semisimple F_v -module. Thus we get a morphism $g_v \in \mathrm{Hom}_{Q_v[G]}(V_v \underline{\mathcal{F}}', V_v \underline{\mathcal{F}})$ with $f_v \circ g_v = \mathrm{id}_{V_v \underline{\mathcal{F}}'}$. Consider the integral Tate modules $T_v \underline{\mathcal{F}}$ and $T_v \underline{\mathcal{F}}'$. We can find some $n \in \mathbb{N}$ such that

$$v^n g_v \in \mathrm{Hom}_{A_v[G]}(T_v \underline{\mathcal{F}}', T_v \underline{\mathcal{F}}) \cong \mathrm{Hom}(\underline{M}(\underline{\mathcal{F}}'), \underline{M}(\underline{\mathcal{F}})) \otimes_A A_v$$

and we choose $g \in \mathrm{Hom}(\underline{M}(\underline{\mathcal{F}}'), \underline{M}(\underline{\mathcal{F}})) \subset \mathrm{QHom}(\underline{\mathcal{F}}', \underline{\mathcal{F}})$ with $g \equiv v^n g_v$ modulo v^m for a sufficiently large $m > n$. If $g \circ f = 0$ in E , then $f \circ g \circ f = 0$, and therefore $f \circ g = 0$ in $\mathrm{QEnd}(\underline{\mathcal{F}}')$ due to the surjectivity of f . This would imply

$$v^n \cdot \mathrm{id}_{V_v \underline{\mathcal{F}}'} = v^n (f_v \circ g_v) = f_v \circ (v^n g_v) \equiv f \circ g = 0 \quad (\text{modulo } v^m)$$

which is a contradiction. Thus $g \circ f \neq 0$ is invertible in E , and therefore f is injective. By that, f gives the desired isomorphism between $\underline{\mathcal{F}}'$ and $\underline{\mathcal{F}}$. The second assertion follows from Theorem 2.7.

2. We already saw one direction in Theorem 2.7/2. So let now $\mathrm{QEnd}(\underline{\mathcal{F}})$ be semisimple and let

$$\mathrm{QEnd}(\underline{\mathcal{F}}) = \bigoplus_{j=1}^m M_{\lambda_j}(E_j)$$

be the decomposition into full matrix algebras $M_{\lambda_j}(E_j)$ over division algebras E_j over Q ($1 \leq j \leq m$). For each j we find λ_j distinct idempotents $e_{j,1}, \dots, e_{j,\lambda_j} \in M_{\lambda_j}(E_j)$ such that $e_{j,\alpha} \cdot \mathrm{QEnd}(\underline{\mathcal{F}}) \cdot e_{j,\alpha} = E_j$ for all $1 \leq \alpha \leq \lambda_j$ with $\sum_{\alpha=1}^{\lambda_j} e_{j,\alpha} = 1$ in $M_{\lambda_j}(E_j)$. Let e_1, \dots, e_n denote all these idempotents, $n = \sum_{j=1}^m \lambda_j$, and choose a divisor D on C such that $e_i \in \mathrm{Hom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}(D))$ for all $1 \leq i \leq n$. Then $\sum_{i=1}^n e_i = \mathrm{id}_{\underline{\mathcal{F}}}$ in $\mathrm{QEnd}(\underline{\mathcal{F}})$ and therefore

$$\underline{\mathcal{F}} \xrightarrow{\sum_i e_i} \bigoplus_{i=1}^n \mathrm{im} e_i \subset \underline{\mathcal{F}}(D).$$

The image $\underline{\mathcal{F}}_i := \mathrm{im} e_i$ is an abelian τ -sheaf by [BH1, Proposition 4.2] because $\varepsilon \neq \infty$. Since $\sum_i e_i$ is injective it is an isogeny by 2.1. Since $\mathrm{QEnd}(\underline{\mathcal{F}}_i) = e_i \cdot \mathrm{QEnd}(\underline{\mathcal{F}}) \cdot e_i$ is a division algebra, $\underline{\mathcal{F}}_i$ is a simple abelian τ -sheaf by 1. Thus $\underline{\mathcal{F}} \approx \underline{\mathcal{F}}_1 \oplus \dots \oplus \underline{\mathcal{F}}_n$ gives the decomposition into a direct sum of simple abelian τ -sheaves $\underline{\mathcal{F}}_i$ as desired. \square

Remark 6.12. Unfortunately the theorem fails if L is not finite, as Example 6.13 below shows. The reason is, that then E_v may still be semisimple while the image F_v of $Q_v[G]$ in $\text{End}_{Q_v}(V_v\mathcal{F})$ is not. Nevertheless, if one adds the assumption that F_v is semisimple, the assertions of Theorem 6.11 remain valid over an arbitrary field L . (See also the remark after Proposition 6.8.)

Example 6.13. We construct a pure Anderson motive \underline{M} over a non-finite field L which is not semisimple, but has $\text{End}(\underline{M}) = A$. Any associated abelian τ -sheaf \mathcal{F} has $\text{QEnd}(\mathcal{F}) = Q$. Let $C = \mathbb{P}_{\mathbb{F}_q}^1$, $A = \mathbb{F}_q[t]$ with $q > 2$, and $L = \mathbb{F}_q(\alpha)$ where α is transcendental over \mathbb{F}_q . Let $M = A_L^{\oplus 2}$ and $\tau = \begin{pmatrix} \alpha t & t \\ 0 & t \end{pmatrix}$. Then $\underline{M} = (M, \tau)$ is a pure Anderson motive of rank and dimension 2. Clearly \underline{M} is not simple, since $\underline{M}' = (A_L, \tau' = t)$ is a factor motive by projecting onto the second coordinate. We will see below that \underline{M} is not even semisimple.

Let $\begin{pmatrix} e & f \\ g & h \end{pmatrix} \in M_2(A_L)$ be an endomorphism of \underline{M} , that is,

$$\begin{pmatrix} \alpha\sigma^*e + \sigma^*g & \alpha\sigma^*f + \sigma^*h \\ \sigma^*g & \sigma^*h \end{pmatrix} = \begin{pmatrix} \alpha e & e + f \\ \alpha g & g + h \end{pmatrix}.$$

Choose $\beta \in \mathbb{F}_q(\alpha)^{\text{alg}} \setminus \mathbb{F}_q(\alpha)$ satisfying $\beta^{q-1} = \alpha$ (for $\beta \notin \mathbb{F}_q(\alpha)$ we use $q > 2$). Then $\sigma^*g = \alpha g$ implies $g \in \beta \cdot \mathbb{F}_q[t]$. Since also $g \in \mathbb{F}_q(\alpha)[t]$ we must have $g = 0$. Now $\sigma^*e = e$ and $\sigma^*h = h$ yielding $e, h \in \mathbb{F}_q[t]$.

Let $\gamma \in \mathbb{F}_q(\alpha)^{\text{alg}} \setminus \mathbb{F}_q(\beta)$ with $\gamma^q - \gamma = \beta$ and set $\tilde{f} := \beta f - \gamma \cdot (e - h)$. Then $\alpha\sigma^*f - f = e - \sigma^*h = e - h$ implies $\sigma^*\tilde{f} - \tilde{f} = \beta^q\sigma^*f - \beta f - (\gamma^q - \gamma)(e - h) = \beta(\alpha\sigma^*f - f - (e - h)) = 0$. Thus $\tilde{f} \in \mathbb{F}_q[t]$ and $\gamma \cdot (e - h) \in \mathbb{F}_q(\beta)[t]$. So we must have $e = h$ and then $\beta f = \tilde{f} \in \mathbb{F}_q[t]$ implies $f = 0$. This shows that $\text{End}(\underline{M}) = \mathbb{F}_q[t] = A$.

The same argument shows that \underline{M} is not even semisimple. Namely, the projection $\underline{M} \rightarrow \underline{M}'$ has no section $\underline{M}' \rightarrow \underline{M}, 1 \mapsto \begin{pmatrix} f \\ 1 \end{pmatrix}$, since there is no solution f for the equation $\alpha t \sigma^* f + t = t f$.

It is also not hard to compute F_v for instance at the place $v = (t - 1)$. Let $z = t - 1$ and $\beta \in L^{\text{sep}}$ with $\beta^{q-1} = \alpha$, and consider the basis $\begin{pmatrix} y/\beta \\ 0 \end{pmatrix}, \begin{pmatrix} x \\ y \end{pmatrix}$ of the Tate module $T_v(\underline{M})$ with

$$\begin{pmatrix} x \\ y \end{pmatrix} = \sum_{i=0}^{\infty} \begin{pmatrix} x_i \\ y_i \end{pmatrix} z^i \quad \text{and} \quad x_i, y_i \in L^{\text{sep}}, y_0 \neq 0.$$

They are subject to the equations $y = t\sigma^*y = (1+z)\sigma^*y$ and $x = \alpha t\sigma^*x + t\sigma^*y = \alpha(1+z)\sigma^*x + y$, that is,

$$\begin{aligned} y_i - y_i^q &= y_{i-1}^q, \text{ and} \\ x_i - \alpha x_i^q &= \alpha x_{i-1}^q + y_i. \end{aligned}$$

There are elements γ and δ of $G = \text{Gal}(L^{\text{sep}}/L)$ operating as $\gamma(y_i) = y_i$, $\gamma(x_i) = x_i$, $\gamma(\beta) = \beta/\eta$ for an $\eta \in \mathbb{F}_q^\times \setminus \{1\}$, respectively as $\delta(y_i) = y_i$, $\delta(\beta) = \beta$, $\delta(x_i) = x_i + y_i/\beta$. With respect to our basis of $T_v(\underline{M})$ they correspond to matrices $\gamma_v = \begin{pmatrix} \eta & 0 \\ 0 & 1 \end{pmatrix}$ and $\delta_v = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. We conclude that F_v is the Q_v -algebra of upper triangular matrices. Its commutant in $M_2(Q_v)$ equals $Q_v \cdot \text{Id}_2 \cong \text{End}(\underline{M}) \otimes_A Q_v$.

Remark. If $q = 2$ any pure Anderson motive of rank $\text{rk } \underline{M} = 2$ on $A = \mathbb{F}_q[t]$, which is not semisimple has $\text{End}(\underline{M}) \supsetneq A$. One easily sees this by choosing a basis of \underline{M} for which τ has the form $\begin{pmatrix} \alpha(t-\theta)^d & * \\ 0 & \beta(t-\theta)^d \end{pmatrix}$ with $\alpha, \beta, \theta \in L$. Then $\begin{pmatrix} 0 & \beta/\alpha \\ 0 & 0 \end{pmatrix}$ is an endomorphism.

However, we expect that also for $q = 2$ there are examples similar to 6.13 (of $\text{rk } \underline{M} \geq 3$), although we have not tried to find one.

Let \mathcal{F} be an abelian τ -sheaf over \mathbb{F}_s and let $\mathbb{F}_{s'}/\mathbb{F}_s$ be a finite field extension. The base extension

$$\mathcal{F} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'} := (\mathcal{F}_i \otimes_{\mathcal{O}_{C_{\mathbb{F}_s}}} \mathcal{O}_{C_{\mathbb{F}_{s'}}}, \Pi_i \otimes 1, \tau_i \otimes 1)$$

is an abelian τ -sheaf over $\mathbb{F}_{s'}$ with $\pi' = (\pi \otimes 1)^t$ for $s' = s^t$, and we have a canonical isomorphism between $V_v \underline{\mathcal{F}}$ and $V_v \underline{\mathcal{F}'}$.

For the next result recall that an endomorphism φ of a finite dimensional vector space V over a field K is called *absolutely semisimple* if for every field extension K'/K the endomorphism $\varphi \otimes 1 \in \text{End}_{K'}(V \otimes_K K')$ is semisimple. The following characterization is taken from [Bou, Proposition 9.2/4 and Proposition 9.2/5].

Lemma 6.14. *Let K be a field and let V be a finite dimensional K -vector space. Let $\varphi \in \text{End}_K(V)$ be an endomorphism.*

1. φ is absolutely semisimple, if and only if there exists a perfect field extension K'/K such that $\varphi \otimes 1 \in \text{End}_{K'}(V \otimes_K K')$ is semisimple.
2. φ is absolutely semisimple, if and only if its minimal polynomial is separable.

Theorem 6.15. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over the finite field \mathbb{F}_s . Then there exists a finite field extension $\mathbb{F}_{s'}/\mathbb{F}_s$ whose degree is a power of $\text{char } \mathbb{F}_s$ such that $\underline{\mathcal{F}} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'}$ has an absolutely semisimple Frobenius endomorphism. Thus if moreover $\varepsilon \neq \infty$ then $\underline{\mathcal{F}} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'}$ is semisimple.*

Remark. It suffices to take $[\mathbb{F}_{s'} : \mathbb{F}_s]$ as the smallest power of $\text{char } \mathbb{F}_s$ which is $\geq \text{rk } \underline{\mathcal{F}}$.

Proof. Let $s' = s^t$ for some arbitrary $t \in \mathbb{N}$. Let $\underline{\mathcal{F}'} := \underline{\mathcal{F}} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'}$ be the abelian τ -sheaf over $\mathbb{F}_{s'}$ induced by $\underline{\mathcal{F}}$. Let $v \in \text{Spec } A$ be a place different from ε . Over Q_v^{alg} we can write $\pi_v \in \text{End}_{Q_v}(V_v \underline{\mathcal{F}})$ in Jordan normal form

$$B^{-1}(\pi_v \otimes 1)B = \begin{pmatrix} \lambda_1 & * & & 0 \\ & \lambda_2 & \ddots & \\ & & \ddots & * \\ 0 & & & \lambda_r \end{pmatrix}$$

for $B \in GL_r(Q_v^{\text{alg}})$ and for some $\lambda_j \in Q_v^{\text{alg}}$, $1 \leq j \leq r$. Thus, by a suitable choice of $t \in \mathbb{N}$ as a power of $\text{char } \mathbb{F}_q$ (as in the remark), we can achieve that $\pi'_v = (\pi_v \otimes 1)^t$ is of the form

$$B^{-1}\pi'_v B = \begin{pmatrix} \lambda_1^t & & & 0 \\ & \lambda_2^t & & \\ & & \ddots & \\ 0 & & & \lambda_r^t \end{pmatrix}.$$

Since Q_v^{alg} is perfect, we conclude by 6.14/1 that π'_v and thus π' is absolutely semisimple. \square

The following corollary illustrates that, in contrast to endomorphisms of vector spaces, there is no need of the term "absolutely semisimple" for abelian τ -sheaves or pure Anderson motives over finite fields.

Corollary 6.16. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over \mathbb{F}_s of characteristic different from ∞ . If $\underline{\mathcal{F}}$ is semisimple, then $\underline{\mathcal{F}} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'}$ is semisimple for every finite field extension $\mathbb{F}_{s'}/\mathbb{F}_s$. The same is true for pure Anderson motives.*

Proof. Let $\underline{\mathcal{F}}$ be semisimple and let $\mathbb{F}_{s'}/\mathbb{F}_s$ be a finite field extension with $s' = s^t$. We set $\underline{\mathcal{F}'} := \underline{\mathcal{F}} \otimes_{\mathbb{F}_s} \mathbb{F}_{s'}$. By 6.11 and 6.8, we know that $\text{QEnd}(\underline{\mathcal{F}}) \otimes_Q Q_v \cong \text{End}_{Q_v[\pi_v]}(V_v \underline{\mathcal{F}})$ is semisimple. Since $Q_v[\pi_v^t] \subset Q_v[\pi_v]$ we conclude by [Bou, Corollaire de Proposition 6.4/9] that $Q_v[\pi_v^t]$ is semisimple, as well. As $V_v \underline{\mathcal{F}'} = V_v \underline{\mathcal{F}}$, we have $\pi'_v = \pi_v^t$, and therefore π'_v is semisimple. Thus, by 6.8, $\text{QEnd}(\underline{\mathcal{F}'})$ is semisimple and $\underline{\mathcal{F}'}$ is semisimple by 6.11/2. \square

7 Zeta Functions and Reduced Norms

In this section we generalize Gekeler's results [Gek] on Zeta functions for Drinfeld modules to pure Anderson motives. But let us begin by recalling a few facts about reduced norms; see for instance [Rei, §9]. Let \underline{M} be a semisimple pure Anderson motive over a finite field and let π be its Frobenius endomorphism. Then $F = Q(\pi)$ is the center of the semisimple algebra E by Corollary 6.10. Write $F = \bigoplus_i F_i$ and $E = \bigoplus_i E_i$ where the F_i are fields and E_i is central simple over F_i . Note that by 6.11 the pure Anderson motive \underline{M} decomposes correspondingly up to isogeny $\underline{M} \approx \bigoplus_i \underline{M}_i$ with $E_i = \text{End}(\underline{M}_i) \otimes_A Q$. We apply 6.6 to \underline{M}_i and obtain $\sum_i [E_i : F_i]^{1/2} \cdot [F_i : Q] = r$. Let $f \in E$ and write it as $f = \sum_i f_i$ with $f_i \in E_i$. Choose for each i a splitting field K_i of E_i with $\alpha_i : E_i \otimes_{F_i} K_i \xrightarrow{\sim} M_{n_i}(K_i)$ where $n_i^2 = [E_i : F_i]$. The *reduced norm of f* is then defined by

$$N(f) := nr_{E/Q}(f) := \prod_i N_{F_i/Q}(\det \alpha_i(f_i \otimes 1)),$$

where $N_{F_i/Q}$ is the usual field norm. The reduced norm is an element of Q which is independent of the choices of K_i and α_i . It satisfies $N(a) = a^r$ for all $a \in Q$, and $N(f) \neq 0$ if and only if $f \in E^\times$, that is, f is a quasi-isogeny. If $f \in \text{End}(\underline{M})$ or more generally f is contained in a finite A -algebra then $N(f) \in A$ since A is normal.

Theorem 7.1. *Let $\underline{\mathcal{F}}$ be a semisimple abelian τ -sheaf over a finite field L and let $f \in \text{QEnd}(\underline{\mathcal{F}})$ be a quasi-isogeny. Then for any place $v \neq \varepsilon, \infty$ of Q we have $N(f) = \det V_v f$, the determinant of the endomorphism $V_v f \in \text{End}_{Q_v}(V_v \underline{\mathcal{F}})$. For $v = \infty \neq \varepsilon$ we have $N(f)^l = \det V_\infty f$, where l comes from Definition 4.1 and satisfies $\dim_{Q_\infty} V_\infty \underline{\mathcal{F}} = l \cdot \text{rk } \underline{\mathcal{F}}$.*

Proof. Clearly, if t is a power of q then $N(f^t) = \det V_v f^t$ implies $N(f) = \det V_v f$ since 1 is the only t -th root of unity in Q_v for $v \neq \infty$, and likewise for $v = \infty$. Writing $V_v f$ in Jordan canonical form over Q_v^{alg} we find as in the proof of Theorem 6.15 a power t of q such that $V_v f^t$ is absolutely semisimple over Q_v and hence its minimal polynomial is separable by 6.14. Then $F_v(f^t)$ and $F(f^t)$ are semisimple by [Bou, Proposition 9.1/1 and Corollaire 7.7/4]. We now replace f by f^t and thus assume that $F(f)$ is semisimple.

As is well known there is a semisimple commutative subalgebra $H = \bigoplus_i H_i$ of E containing $F(f)$ with $\dim_{F_i} H_i = n_i$ and hence $\dim_Q H = r$. Then $nr_{E/Q}(f)$ equals the determinant of the Q -endomorphism $\tilde{f} : x \mapsto fx$ of H . The reason for this is that $H_i \otimes_{F_i} K_i$ is still semisimple and commutative if we choose a splitting field K_i which is separable over F_i . By Lemma 7.2 below $H_i \otimes_{F_i} K_i$ is isomorphic to $K_i^{n_i}$ as left $H_i \otimes_{F_i} K_i$ -modules, and this implies that $nr_{E_i/F_i}(f_i) = \det \alpha_i(f_i) = \det \tilde{f}_i$, the determinant of the F_i -endomorphism $\tilde{f}_i : x \mapsto f_i x$ of H_i , and $N(f) = \det \tilde{f}$ the determinant of the Q -endomorphism \tilde{f} of H .

If $v \neq \infty$ then again by Lemma 7.2, H_v is H_v -isomorphic to $V_v \underline{\mathcal{F}}$ and $N(f) = \det \tilde{f} = \det V_v f$.

If $v = \infty$ we embed $E_\infty^{\oplus l}$ into $\text{End}_{Q_{\infty,L}[\phi]}(\tilde{N}_\infty(\underline{\mathcal{F}}))$. Namely, if $(f^{(0)}, \dots, f^{(l-1)}) \in E_\infty^{\oplus l}$, where $f^{(m)} = (f_i^{(m)} : \mathcal{F}_i \otimes_{\mathcal{O}_{C_L}} Q_{\infty,L} \rightarrow \mathcal{F}_i \otimes_{\mathcal{O}_{C_L}} Q_{\infty,L})$, we set

$$g_{ij} := \begin{cases} \Pi_{i-1} \circ \dots \circ \Pi_j \circ f_j^{(i-j)} & \text{if } 0 \leq j \leq i \leq l-1 \\ z^k \Pi_i^{-1} \circ \dots \circ \Pi_{j-1}^{-1} \circ f_j^{(l+i-j)} & \text{if } 0 \leq i < j \leq l-1. \end{cases}$$

Then $g_{ij} : \mathcal{F}_j \otimes_{\mathcal{O}_{C_L}} Q_{\infty,L} \rightarrow \mathcal{F}_i \otimes_{\mathcal{O}_{C_L}} Q_{\infty,L}$ and a straightforward computation shows that the homomorphism $g = (g_{ij})_{i,j=0..l-1}$ commutes with ϕ from (3.1) on page 10, that is, g is an element of $\text{End}_{Q_{\infty,L}[\phi]}(\tilde{N}_\infty(\underline{\mathcal{F}})) = \text{End}_{Q_\infty[G]}(V_\infty \underline{\mathcal{F}})$; use Proposition 3.4. Now we apply Lemma 7.2 to $H_\infty^{\oplus l} \subset E_\infty^{\oplus l} \subset \text{End}_{Q_\infty}(V_\infty \underline{\mathcal{F}})$, and we compute $N(f)^l = (\det \tilde{f})^l = \det_{Q_\infty}(H_\infty^{\oplus l} \rightarrow H_\infty^{\oplus l}, h \mapsto fh) = \det V_\infty f$ as desired. \square

Lemma 7.2. *Let K be a field and let $H \subset M_n(K)$ be a semisimple commutative K -algebra with $\dim_K H = n$. Then as a (left) module over itself H is isomorphic to K^n .*

Proof. Decomposing H into a direct sum of fields $\bigoplus_{\kappa} L_{\kappa}$ and K^n into a direct sum $\bigoplus_{\lambda} V_{\lambda}$ of simple H -modules, each V_{λ} is isomorphic to an $L_{\kappa(\lambda)}$. The injectivity of $H \rightarrow M_n(K)$ and $\dim_K H = n$ imply that H is isomorphic to $\bigoplus_{\lambda} \text{End}_{L_{\kappa(\lambda)}}(V_{\lambda})$ and a fortiori isomorphic as left module over itself to K^n . \square

Theorem 7.3. *Let \underline{M} be a semisimple pure Anderson motive of rank r over a finite field L and let $f \in \text{End}(\underline{M})$ be an isogeny. Then*

1. $\dim_L \text{coker } N(f) = r \cdot \dim_L \text{coker } f$.
2. The ideal $\deg(f) = N(f) \cdot A$ is principal and has a canonical generator.
3. There exists a canonical dual isogeny $f^{\vee} \in \text{End}(\underline{M})$ satisfying $f \circ f^{\vee} = N(f) = f^{\vee} \circ f$.

Remark. 1. This shows that $N(1 - \pi^n) \in A$ is the analogue for pure Anderson motives of the number of rational points $X(\mathbb{F}_{q^n}) = \deg(1 - \text{Frob}_q^n) \in \mathbb{Z}$ on an abelian variety X over the finite field \mathbb{F}_q ; see also Theorem 7.7 below.

2. The dual isogeny satisfies $(fg)^{\vee} = g^{\vee} f^{\vee}$, because $N(fg) = N(f)N(g)$. Note however, that we cannot expect that $(f + g)^{\vee} = f^{\vee} + g^{\vee}$ unless $r = 2$ because for $f = a \in A$ we have $N(a) = a^r$ and $a^{\vee} = a^{r-1}$.

Proof. 1. Clearly for any $a \in A$ we have $\dim_L \underline{M}/a\underline{M} = r \cdot \dim_{\mathbb{F}_q} A/(a) = -r \cdot \infty(a)$ where $\infty(a)$ denotes the ∞ -adic valuation of a . Now let $\underline{\mathcal{F}}$ be an abelian τ -sheaf with $\underline{M} = \underline{M}(\underline{\mathcal{F}})$, and let $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}(n \cdot \infty)$ for some n be the isogeny induced by f . Using Theorem 7.1 we compute the dimension

$$\begin{aligned} l \cdot \dim_L \text{coker } f &= nrl - \dim_L \bigoplus_{j=0}^{l-1} (\mathcal{F}_j(n \cdot \infty)/f_j(\mathcal{F}_j))_{\infty} \\ &= nrl - \dim_L \tilde{M}_{\infty}(\underline{\mathcal{F}}(n \cdot \infty))/\tilde{M}_{\infty}(f)(\tilde{M}_{\infty}(\underline{\mathcal{F}})) \\ &= nrl - \dim_{\mathbb{F}_q} (T_{\infty} \underline{\mathcal{F}}(n \cdot \infty)/T_{\infty} f(T_{\infty} \underline{\mathcal{F}})) \\ &= -\infty(\det V_{\infty} f) = -l \cdot \infty(N(f)). \end{aligned}$$

Here the first equality follows from the identities $\mathcal{F}_j(n \cdot \infty)/f_j(\mathcal{F}_j) = (\mathcal{F}_j(n \cdot \infty)/f_j(\mathcal{F}_j))_{\infty} \oplus \text{coker } f$ and $\dim_L(\mathcal{F}_j(n \cdot \infty)/f_j(\mathcal{F}_j)) = \deg \mathcal{F}_j(n \cdot \infty) - \deg f_j(\mathcal{F}_j) = nr$. The second equality is the definition of \tilde{M}_{∞} , and the third follows from the isomorphism $\tilde{M}_{\infty}(\underline{\mathcal{F}}) \otimes_{A_{\infty, L}} A_{\infty, L^{\text{sep}}} \cong T_{\infty} \underline{\mathcal{F}} \otimes_{A_{\infty}} A_{\infty, L^{\text{sep}}}$. The fourth equality follows from the elementary divisor theorem. From this we obtain 1.

2. Let $v \neq \varepsilon$ be a maximal ideal of A . Using Theorem 7.1 we compute the v -adic valuation of $N(f)$

$$v(N(f)) = v(\det T_v f) = \dim_{\mathbb{F}_v} (T_v \underline{M}/T_v f(T_v \underline{M})) = \dim_{\mathbb{F}_v} ((\text{coker } f)_v \otimes_L L^{\text{sep}})^{\tau} = v(\deg f).$$

Again the second equality follows from the elementary divisor theorem, the third equality comes from the fact that the τ -invariants of the v -primary part $(\text{coker } f)_v \otimes_L L^{\text{sep}}$ are isomorphic to $T_v \underline{M}/T_v f(T_v \underline{M})$, and the last equality is the definition of $\deg f$. From 1 and Lemma 2.9 we obtain

$$\begin{aligned} r \cdot \dim_{\mathbb{F}_q} A/\deg(f) &= r \cdot \dim_L \text{coker } f = \dim_L \text{coker } N(f) \\ &= \dim_L ((A/N(f))^r \otimes_{\mathbb{F}_q} L) = r \cdot \dim_{\mathbb{F}_q} A/N(f). \end{aligned}$$

From the identity $\dim_{\mathbb{F}_q} A/\mathfrak{a} = \sum_v [\mathbb{F}_v : \mathbb{F}_q] \cdot v(\mathfrak{a})$ for any ideal $\mathfrak{a} \subset A$ we conclude $\varepsilon(\deg f) = \varepsilon(N(f))$ and therefore $\deg(f) = N(f) \cdot A$.

Finally 3 is immediate since $N(f)$ annihilates $\text{coker } f$ by Proposition 2.10. \square

Remark 7.4. We do not know of a proof of 1 and 2 for arbitrary pure Anderson motives which does not make use of the associated abelian τ -sheaf $\underline{\mathcal{F}}$. In the special case when \underline{M} comes from a Drinfeld module, Gekeler [Gek, Lemma 3.1] argued that both sides of the equation in 2 are extensions to E of the ∞ -adic valuation on \mathcal{Q} . But this argument fails in general, since there may be more than one such extension as one sees from Example 9.5 below.

Corollary 7.5. *Let \underline{M} be a semisimple pure Anderson motive of dimension d over a finite field L and let π be its Frobenius endomorphism. Let $v \neq \varepsilon$ be a maximal ideal of A and let χ_v be the characteristic polynomial of π_v . Then*

1. $\chi_v \in A[x]$ is independent of v and $\chi_v(a) \cdot A = \det V_v(a - \pi) \cdot A = \deg(a - \pi)$ for every $a \in A$,
2. $\varepsilon^{d \cdot [L : \mathbb{F}_\varepsilon]} = \deg(\pi) = \chi_v(0) \cdot A = N(\pi) \cdot A$ is principal.

Proof. 1 is a direct consequence of Theorems 7.1 and 7.3 and the Lagrange interpolation theorem applied to the fact that $\chi_v(a) = N(a - \pi) = \chi_w(a) \in A$ for all $a \in A$.

2 follows from the fact that $\text{coker } \pi$ is supported on ε and from the equation $\dim_L \text{coker } \pi = [L : \mathbb{F}_q] \cdot \dim_L \text{coker } \tau = d \cdot [L : \mathbb{F}_q]$. \square

Definition 7.6. *We define the Zeta function of a pure Anderson motive \underline{M} over a finite field \mathbb{F}_s as*

$$Z_{\underline{M}}(t) := \prod_{0 \leq i \leq r} \det(1 - t \wedge^i \pi_v)^{(-1)^{i+1}}$$

where $\varepsilon \neq v \in \text{Spec } A$ is a maximal ideal and $\wedge^i \pi_v \in \text{End}_{Q_v}(\wedge^i V_v \underline{M})$.

By 7.5/1 the Zeta function $Z_{\underline{M}}(t)$ is independent of the place v and lies in $Q(t)$. This also follows from work of Böckle [Boe] and Gardeyn [Gar, §7]. The name ‘‘Zeta function’’ is justified by the following theorem (see also the remark after Theorem 7.3).

Theorem 7.7. *If \underline{M} is semisimple and $\sum_i a_i t^i$ is the power series expansion of $t \frac{d}{dt} \log Z_{\underline{M}}(t)$, then $a_i = N(1 - \pi^i) \in A$.*

Proof. By standard arguments $a_i = \det(1 - \pi_v^i)$; see [Gek, Lemma 5.6]. Now our assertion follows from Theorem 7.1 \square

This Zeta function satisfies the Riemann hypothesis:

Theorem 7.8. *In an algebraic closure of Q_∞ all eigenvalues of $\wedge^i \pi_v \in \text{End}_{Q_v}(\wedge^i V_v \underline{M})$ have the same absolute value $(\#\mathbb{F}_s)^{i \cdot \text{wt}(\underline{M})}$.*

Proof. This was proved by Goss [Gos, Theorem 5.6.10] for $i = 1$ and follows for the remaining i by general arguments of linear algebra. \square

8 A Quasi-Isogeny Criterion

Similarly to the theory for abelian varieties, the characteristic polynomials of the Frobenius endomorphisms on the associated Tate modules play an important role for the study of abelian τ -sheaves. For example, we can decide on quasi-isogeny of two abelian τ -sheaves $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}'}$ just by considering these characteristic polynomials.

Theorem 8.1. *Let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}'}$ be abelian τ -sheaves over \mathbb{F}_s with respective Frobenius endomorphisms π and π' , and let μ_π and $\mu_{\pi'}$ be their minimal polynomials over Q . Let $v \in C$ be a place different from ∞ and ε . Let χ_v and χ'_v be the characteristic polynomials of π_v and π'_v , respectively, and let $G := \text{Gal}(L^{\text{sep}}/L)$. Assume in addition that $\varepsilon \neq \infty$, or that $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}'}$ have the same weight.*

1. Consider the following statements:

- 1.1. $\underline{\mathcal{F}'}$ is quasi-isogenous to an abelian factor τ -sheaf of $\underline{\mathcal{F}}$.
- 1.2. $V_v \underline{\mathcal{F}'}$ is G -isomorphic to a G -factor space of $V_v \underline{\mathcal{F}}$.
- 1.3. χ'_v divides χ_v in $Q_v[x]$.

1.4. $\mu_{\pi'}$ divides μ_{π} in $Q[x]$ and $\text{rk } \underline{\mathcal{F}}' \leq \text{rk } \underline{\mathcal{F}}$

We have $1.1 \Rightarrow 1.2 \Rightarrow 1.3$ and 1.4 always,
 $1.2 \Leftarrow 1.3$ if π_v and π'_v are semisimple,
 $1.2 \Leftarrow 1.3 \Leftarrow 1.4$ if μ_{π} is irreducible in $Q[x]$,
 $1.1 \Leftarrow 1.2$ if the characteristic is different from ∞ .

2. Consider the following statements:

- 2.1. $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ are quasi-isogenous.
- 2.2. $V_v \underline{\mathcal{F}}$ and $V_v \underline{\mathcal{F}}'$ are G -isomorphic.
- 2.3. $\chi_v = \chi'_v$.
- 2.4. $\mu_{\pi} = \mu_{\pi'}$ and $\text{rk } \underline{\mathcal{F}} = \text{rk } \underline{\mathcal{F}}'$.
- 2.5. There is an isomorphism of Q -algebras $\text{QEnd}(\underline{\mathcal{F}}) \cong \text{QEnd}(\underline{\mathcal{F}}')$ mapping π to π' .
- 2.6. There is a Q_v -isomorphism $\text{QEnd}(\underline{\mathcal{F}}) \otimes_Q Q_v \cong \text{QEnd}(\underline{\mathcal{F}}') \otimes_Q Q_v$ mapping π_v to π'_v .
- 2.7. If $\varepsilon \neq \infty$ also consider the statement $Z_{\underline{M}(\underline{\mathcal{F}})} = Z_{\underline{M}(\underline{\mathcal{F}}')}$.

We have $2.1 \Leftrightarrow 2.2 \Rightarrow 2.3, 2.4, 2.5$ always,
 $2.5 \Rightarrow 2.6$ always,
 $2.3 \Leftrightarrow 2.7$ if the characteristic is different from ∞ ,
 $2.2 \Leftarrow 2.3 \Leftarrow 2.6$ if π_v and π'_v are semisimple,
 $2.2 \Leftarrow 2.3 \Leftarrow 2.4 \Leftarrow 2.6$ if μ_{π} and $\mu_{\pi'}$ are irreducible in $Q[x]$.

Proof. 1. For the implication $1.1 \Rightarrow 1.2$ without loss of generality, $\underline{\mathcal{F}}'$ can itself be considered as abelian factor τ -sheaf of $\underline{\mathcal{F}}$ and the implication follows from Proposition 4.5. The implication $1.2 \Rightarrow 1.3$ is obvious.

For $1.2 \Rightarrow 1.4$ note that μ_{π} is also the minimal polynomial of π_v over Q_v by Lemma 6.2. By Proposition 5.5 statement 1.2 implies $\mu_{\pi}(\pi'_v) = 0$, whence 1.4.

For $1.3 \Rightarrow 1.2$ let π_v and π'_v be semisimple. Let $\chi_v = \mu_1 \cdots \mu_n$ and $\chi'_v = \mu'_1 \cdots \mu'_{n'}$ be the factorization in $Q_v[x]$ into irreducible factors and set $V_i := Q_v[x]/(\mu_i)$ and $V'_i := Q_v[x]/(\mu'_i)$. Then we can decompose $V_v \underline{\mathcal{F}} = V_1 \oplus \cdots \oplus V_n$ and $V_v \underline{\mathcal{F}}' = V'_1 \oplus \cdots \oplus V'_{n'}$. Since χ'_v divides χ_v , we can now easily construct a surjective G -morphism from $V_v \underline{\mathcal{F}}$ onto $V_v \underline{\mathcal{F}}'$ which gives the desired result.

Next if μ_{π} is irreducible, 1.4 implies $\mu_{\pi'} = \mu_{\pi}$ and 1.3 follows from Corollary 6.6. It further follows from Proposition 6.8 that π_v and π'_v are semisimple and this implies 1.2 by the above.

For $1.2 \Rightarrow 1.1$ we first do not assume that $\varepsilon \neq \infty$. Let $f_v : V_v \underline{\mathcal{F}} \rightarrow V_v \underline{\mathcal{F}}'$ be a surjective morphism of $Q_v[G]$ -modules. We may multiply f_v by a suitable power of v to get a morphism $f_v : T_v \underline{\mathcal{F}} \rightarrow T_v \underline{\mathcal{F}}'$ of the integral Tate modules which is not necessarily surjective, but satisfies $v^n T_v \underline{\mathcal{F}}' \subset f_v(T_v \underline{\mathcal{F}})$ for a sufficiently large n . Let $\underline{M} := (\Gamma(C_L \setminus \{\infty\}, \mathcal{F}_0), \Pi_0^{-1} \circ \tau)$. This is a “ τ -module on A ” in the sense of [BH1, Definition 3.2]. If $\varepsilon \neq \infty$ then \underline{M} is the pure Anderson motive $\underline{M}(\underline{\mathcal{F}})$ associated with $\underline{\mathcal{F}}$ in (1.1). Also let $\underline{M}' := (\Gamma(C_L \setminus \{\infty\}, \mathcal{F}'_0), \Pi_0'^{-1} \circ \tau')$. By [BH1, Theorem 9.8] (or Theorem 4.2 if $\varepsilon \neq \infty$), f_v lies inside $\text{Hom}(\underline{M}, \underline{M}') \otimes_A A_v$, so we can approximate f_v by some $f \in \text{Hom}(\underline{M}, \underline{M}')$ with $T_v(f) \equiv f_v$ modulo $v^{n+1} T_v \underline{M}'$. Since $v^n T_v \underline{M}' \subset f_v(T_v \underline{M})$ we find inside $\text{im } T_v(f)$ generators of $v^n T_v \underline{M}' / v^{n+1} T_v \underline{M}'$. They generate an A_v -submodule of $v^n T_v \underline{M}'$ whose rank must at least be r' since $v^n T_v \underline{M}' / v^{n+1} T_v \underline{M}' \cong (A_v / v A_v)^{r'}$. Thus $\text{im } T_v(f)$ has rank r' . Either by assumption or by [BH1, Corollary 3.5] if $\varepsilon \neq \infty$, both $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}'$ have the same weight. So by [BH1, Proposition 6.10/1], f comes from a quasi-morphism $f \in \text{QHom}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$, that is, a morphism $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'(D)$ for a suitable divisor D . Now we finally assume that the characteristic is different from ∞ . By [BH1, Proposition 4.2], the image $\text{im}(f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'(D))$ is an abelian factor τ -sheaf of $\underline{\mathcal{F}}$ and $\text{im } f \rightarrow \underline{\mathcal{F}}'(D)$ is an injective morphism between abelian τ -sheaves of the same rank and weight, hence an isogeny by Proposition 2.1.

2. A large part of 2 follows from 1. We prove the rest. To show $2.2 \Rightarrow 2.1$ without the hypothesis on the characteristic, we just replace the last argument of the proof of $1.2 \Rightarrow 1.1$ by the following: Since $r = \dim_{Q_v} V_v \underline{\mathcal{F}} = \dim_{Q_v} V_v \underline{\mathcal{F}}' = r'$, the morphism $f : \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}'(D)$ is an injective morphism between abelian τ -sheaves of the same rank and weight, hence an isogeny by Proposition 2.1.

For the implication $2.1 \Rightarrow 2.5$ let $g \in \text{QIsog}(\underline{\mathcal{F}}, \underline{\mathcal{F}}')$. Then the map $\text{QEnd}(\underline{\mathcal{F}}) \rightarrow \text{QEnd}(\underline{\mathcal{F}}')$ sending $f \mapsto gfg^{-1}$ is an isomorphism with $\pi' = g\pi g^{-1}$. The implication $2.5 \Rightarrow 2.6$ is obvious.

For the implication $2.3 \Rightarrow 2.7$ note that knowledge of χ_v yields the knowledge of $\det(1 - t \wedge^i \pi_v)$ and thus of $Z_{\underline{M}}(\underline{\mathcal{F}})$ by linear algebra. Conversely we know from Theorem 7.8 that all zeroes of $\det(1 - t \wedge^i \pi_v)$ have absolute value $s^{-i \text{wt}(\underline{\mathcal{F}})}$ in an algebraic closure of Q_∞ . So we can recover χ_v from $Z_{\underline{M}}(\underline{\mathcal{F}})$ by simply looking at this absolute value. This proves $2.3 \Leftarrow 2.7$.

Next if π_v and π'_v are semisimple $2.6 \Rightarrow 2.3$ follows from Lemma 6.4/2, and $2.3 \Rightarrow 2.2$ was already established in 1.

Finally if μ_π and $\mu_{\pi'}$ are irreducible, 2.4 follows from 2.6 by Corollary 6.6 since μ_π is also the minimal polynomial of π_v over Q_v by Lemma 6.2. Also 2.3 follows from 2.4 by Corollary 6.6 and π_v and π'_v are semisimple, so $2.3 \Rightarrow 2.2$ by the above. \square

9 The Endomorphism Q -Algebra

In this section we study the structure of $\text{QEnd}(\underline{\mathcal{F}})$ for a semisimple abelian τ -sheaf $\underline{\mathcal{F}}$ over a finite field and calculate the local Hasse invariants of $\text{QEnd}(\underline{\mathcal{F}})$ as a central simple algebra over $Q(\pi)$. For a detailed introduction to central simple algebras, Hasse invariants and the Brauer group, we refer to [Rei, Ch. 7, §§28–31].

Theorem 9.1. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over the finite field \mathbb{F}_s of rank r with semisimple Frobenius endomorphism π , that is, $Q(\pi)$ is semisimple. Let $v \in C$ be a place different from ∞ and from the characteristic point ε . Let χ_v be the characteristic polynomial of π_v .*

1. *The algebra $F = Q(\pi)$ is the center of the semisimple algebra $E = \text{QEnd}(\underline{\mathcal{F}})$.*
2. *We have $r \leq [E : Q] = r_{Q_v}(\chi_v, \chi_v) \leq r^2$.*
3. *Consider the following statements:*

- 3.1. $E = F$.
- 3.2. E is commutative.
- 3.3. $[F : Q] = r$.
- 3.4. $[E : Q] = r$.
- 3.5. χ_v has no multiple factor in $Q_v[x]$.
- 3.6. χ_v is separable.

*We have $3.1 \Leftrightarrow 3.2 \Leftrightarrow 3.3 \Leftrightarrow 3.4 \Leftrightarrow 3.5 \Leftarrow 3.6$ always,
 $3.5 \Rightarrow 3.6$ if π_v is absolutely semisimple.*

4. *Consider the following statements:*

- 4.1. $F = Q$.
- 4.2. E is a central simple algebra over Q .
- 4.3. $[E : Q] = r^2$.
- 4.4. χ_v is the r -th power of a linear polynomial in $Q_v[x]$.
- 4.5. χ_v is purely inseparable.

*We have $4.1 \Leftrightarrow 4.2 \Leftrightarrow 4.3 \Leftrightarrow 4.4 \Rightarrow 4.5$ always,
 $4.4 \Leftarrow 4.5$ if π_v is absolutely semisimple.*

If 4.2 holds and moreover the characteristic point $\varepsilon := c(\text{Spec } \mathbb{F}_s) \in C_{\mathbb{F}_s}$ is different from ∞ , E is characterized by $\text{inv}_\infty E = \text{wt}(\underline{\mathcal{F}})$, $\text{inv}_\varepsilon E = -\text{wt}(\underline{\mathcal{F}})$ and $\text{inv}_v E = 0$ for any other place $v \in C$.

5. *In general the local Hasse invariants of E at the places v of F equal $\text{inv}_v E = -\frac{[\mathbb{F}_v : \mathbb{F}_q]}{[\mathbb{F}_s : \mathbb{F}_q]} \cdot v(\pi)$. In particular*

$$\text{inv}_v E = \begin{cases} 0 & \text{if } v \nmid \varepsilon \infty, \\ \text{wt}(\underline{\mathcal{F}}) \cdot [F_v : Q_\infty] & \text{if } v \mid \infty \text{ and } \varepsilon \neq \infty. \end{cases}$$

(Here F_v denotes the completion of F at the place v and \mathbb{F}_v is the residue field of the place v .)

Remark 9.2. If $\varepsilon \neq \infty$ and $\underline{\mathcal{F}}$ is an elliptic sheaf, that is, $d = 1$ and $\underline{M}(\underline{\mathcal{F}})$ is the Anderson motive of a Drinfeld module, Gekeler [Gek, Theorem 2.9] has shown that there is exactly one place v of F above ε , and exactly one place w of F above ∞ , and that $\text{inv}_w E = [F : Q] \cdot \text{wt}(\underline{\mathcal{F}})$ and $\text{inv}_v E = -[F : Q] \cdot \text{wt}(\underline{\mathcal{F}})$. Note that Gekeler actually computes the Hasse invariants of the endomorphism algebra of the Drinfeld module. So his invariants differ from ours by a minus sign, since passing from Drinfeld modules to abelian τ -sheaves is a contravariant functor, see [BS, Theorem 3.2.1].

Corollary 9.3. *Let $\underline{\mathcal{F}}$ be an abelian τ -sheaf over the smallest possible field $L = \mathbb{F}_q$ such that $\text{QEnd}(\underline{\mathcal{F}})$ is a division algebra. Then $\text{QEnd}(\underline{\mathcal{F}})$ is commutative and equals $Q(\pi)$.*

Proof. $\text{QEnd}(\underline{\mathcal{F}})$ is a central division algebra over F by Theorem 9.1, which splits at all places of F by 9.1/5, hence equals F . \square

Proof (of Theorem 9.1). 1 was already proved in Corollary 6.10.

2. Let

$$\chi_v = \prod_{i=1}^n \mu_i^{m_i} \in Q_v[x]$$

with distinct irreducible $\mu_i \in Q_v[x]$ and $m_i > 0$ for $1 \leq i \leq n$. Then $\sum_{i=1}^n m_i \cdot \deg \mu_i = \deg \chi_v = r$, and by Theorem 6.5 we have $[E : Q] = r_{Q_v}(\chi_v, \chi_v) = \sum_{i=1}^n m_i^2 \cdot \deg \mu_i$. The result now follows from the obvious inequalities

$$r = \sum_{i=1}^n m_i \cdot \deg \mu_i \stackrel{(1)}{\leq} \sum_{i=1}^n m_i^2 \cdot \deg \mu_i \stackrel{(2)}{\leq} \left(\sum_{i=1}^n m_i \cdot \deg \mu_i \right)^2 = r^2. \quad (9.1)$$

3. Since $F = Z(E)$, the equivalence 3.1 \Leftrightarrow 3.2 is evident. We have equality in (1) of Equation (9.1) if and only if $m_i = 1$ for all $1 \leq i \leq s$ which establishes the equivalence 3.4 \Leftrightarrow 3.5. In order to prove 3.5 \Rightarrow 3.3 we consider the minimal polynomial μ_v of π_v over Q_v . If χ_v has no multiple factor, then $\mu_v = \chi_v$ and therefore $[F : Q] = [Q_v(\pi_v) : Q_v] = r$. Next 3.3 \Rightarrow 3.1 because $F \subset E$ and $(\dim_{Q_v} F_v)(\dim_{Q_v} E_v) = \dim_{Q_v} \text{End}_{Q_v}(V_v \underline{\mathcal{F}}) = r^2$ by [Bou, Théorème 10.2/2], since E_v is the commutant of F_v in $\text{End}_{Q_v}(V_v \underline{\mathcal{F}})$. Note that 3.3 \Rightarrow 3.1 also follows from Lemma 7.2. Conversely 3.1 \Rightarrow 3.4 because $E = F$ implies $r \geq [Q_v(\pi_v) : Q_v] = [F : Q] = [E : Q] \geq r$. For 3.5 \Rightarrow 3.6 we use Lemma 6.14/2 as we know that $\chi_v = \mu_v$. 3.6 \Rightarrow 3.5 is clear.

4. If $F = Q$, then E is simple with center Q , so E is a central simple algebra over Q . Since $F = Z(E)$, the converse is obvious. This shows 4.1 \Leftrightarrow 4.2. We have equality in (2) of (9.1) if and only if $n = 1$, $\deg \mu_1 = 1$ and $m_1 = r$ which establishes 4.3 \Leftrightarrow 4.4. In order to connect 4.1 \Leftrightarrow 4.2 with 4.3 \Leftrightarrow 4.4 let χ_v be a power of a linear polynomial. By [Bou, Proposition 9.1/1] the minimal polynomial of π_v over Q_v is linear and thus $F = Q$. The converse is trivial. For 4.5 \Rightarrow 4.4 we use again 6.14/2 to see that μ_v is linear. 4.4 \Rightarrow 4.5 is clear.

The statement about the Hasse invariants follows from 5. Nevertheless, we give a separate proof in case $(k, l) = 1$ using Tate modules, since this is much shorter here and exhibits a different technique than 5. By the Tate conjecture 4.3, $E \otimes_Q Q_v$ is isomorphic to $\text{End}_{Q_v}(V_v \underline{\mathcal{F}}) \cong M_r(Q_v)$ for all places $v \in C$ which are different from ε and ∞ , so the Hasse invariants of E at these places are 0. Since the sum of all Hasse invariants is 0 (modulo 1), we only need to calculate $\text{inv}_\infty E$.

As a first step, we show that \mathbb{F}_{q^l} is contained in \mathbb{F}_s . In our situation, π lies inside Q . Thus, by 7.8 we get $s^{k/l} = |\pi|_\infty = q^m$ for some $m \in \mathbb{Z}$ as $|Q_\infty^\times|_\infty = q^{\mathbb{Z}}$. Since $q^e = s$, we conclude that $e \cdot k/l = m \in \mathbb{Z}$ and hence $l \mid e$, since k and l are assumed to be relatively prime. Therefore $\mathbb{F}_{q^l} \subset \mathbb{F}_{q^e} = \mathbb{F}_s$.

Consider the rational Tate module $V_\infty(\underline{\mathcal{F}})$ at ∞ and the isomorphism of Q_∞ -algebras

$$E \otimes_Q Q_\infty \cong \text{End}_{\Delta_\infty[G]}(V_\infty \underline{\mathcal{F}}) = \text{End}_{\Delta_\infty}(V_\infty \underline{\mathcal{F}})$$

from Theorem 4.3. Since $\dim_{Q_\infty} \Delta_\infty = l^2$ and $\dim_{Q_\infty} V_\infty \underline{\mathcal{F}} = rl$, we conclude that $V_\infty \underline{\mathcal{F}}$ is a left r/l -dimensional Δ_∞ -vector space and hence isomorphic to $\Delta_\infty^{r/l}$. Thus we have

$$E \otimes_Q Q_\infty \cong \text{End}_{\Delta_\infty}(\Delta_\infty^{r/l}) = M_{r/l}(\text{End}_{\Delta_\infty}(\Delta_\infty)) = M_{r/l}(\Delta_\infty^{\text{op}}).$$

Our proof now completes by $\text{inv}_\infty E = \text{inv} \Delta_\infty^{\text{op}} = -\text{inv} \Delta_\infty = \frac{k}{l} = \text{wt}(\underline{\mathcal{F}})$.

5. We prove the general case using local (iso-)shtuka rather than Tate modules which were used in 4. Our method is inspired by Milne's and Waterhouse' computation for abelian varieties [WM, Theorem 8]. However in the function field case this method can be used to calculate the Hasse invariant at *all* places, whereas in the number field case it applies only to the place which equals the characteristic of the ground field. Let w be a place of Q and let $\underline{N}_w := \underline{N}_w(\underline{\mathcal{F}})$ be the local σ -isoshtuka of $\underline{\mathcal{F}}$ at w . Let \mathbb{F}_w be the residue field of w and $\mathbb{F}_{q^f} = \mathbb{F}_w \cap \mathbb{F}_s$ the intersection inside an algebraic closure of \mathbb{F}_q . Let \mathfrak{a}_0 be the ideal $(b \otimes 1 - 1 \otimes b : b \in \mathbb{F}_{q^f})$ of $Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s$ and let $R := (Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s / \mathfrak{a}_0)[T] = Q_w \otimes_{\mathbb{F}_{q^f}} \mathbb{F}_s[T]$ be the non-commutative polynomial ring with $T \cdot (a \otimes b) = (a \otimes b^{q^f}) \cdot T$ for $a \in Q_w$ and $b \in \mathbb{F}_s$. Since $Q_w \otimes_{\mathbb{F}_{q^f}} \mathbb{F}_s$ is a field, R is a non-commutative principal ideal domain as studied by Jacobson [Jac, Chapter 3]. Its center is the commutative polynomial ring $Q_w[T^g]$ where $g = [\mathbb{F}_s : \mathbb{F}_{q^f}] = \frac{e}{f}$. From Theorem 3.6 and Proposition 3.5 we get isomorphisms

$$\text{QEnd}(\underline{\mathcal{F}}) \otimes_Q Q_w \cong \text{End}_{Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s / [\phi]}(\underline{N}_w) \cong \text{End}_R(\underline{N}_w / \mathfrak{a}_0 \underline{N}_w)$$

where T operates on $\underline{N}_w / \mathfrak{a}_0 \underline{N}_w$ as ϕ^f .

By [Jac, Theorem 3.19] the R -module $\underline{N}_w / \mathfrak{a}_0 \underline{N}_w$ decomposes into a finite direct sum indexed by some set I

$$\underline{N}_w / \mathfrak{a}_0 \underline{N}_w \cong \bigoplus_{v \in I} \underline{N}_v^{\oplus n_v} \quad (9.2)$$

of indecomposable R -modules \underline{N}_v with $\underline{N}_v \not\cong \underline{N}_{v'}$ for $v \neq v'$. The annihilator of \underline{N}_v is a two sided ideal of R generated by a central element $\mu_v \in Q_w[T^g]$ by [Jac, §3.6], which can be chosen to be monic. In particular (9.2) is an isomorphism of $Q_w[T^g]$ -modules and μ_v is the minimal polynomial of T^g on \underline{N}_v by [Jac, Lemma 3.1]. Therefore the least common multiple μ of the μ_v is the minimal polynomial of T^g on $\underline{N}_w / \mathfrak{a}_0 \underline{N}_w$. Note that T^g operates on $\underline{N}_w / \mathfrak{a}_0 \underline{N}_w$ as the Frobenius π , hence $\mu = \text{mipo}_{\pi|_{\underline{\mathcal{F}}}}$ and $F = Q(\pi) = Q[T^g]/(\mu)$, where we write *mipo* for the minimal polynomial. By the semisimplicity of π (and Proposition 6.8) μ has no multiple factors in $Q_w[T^g]$. Since the μ_v are powers of irreducible polynomials by [Jac, Theorem 3.20] we conclude that all μ_v are themselves irreducible in $Q_w[T^g]$. Again [Jac, Theorem 3.20] implies that $\mu_v \neq \mu_{v'}$ since $\underline{N}_v \not\cong \underline{N}_{v'}$ and

$$\mu = \text{mipo}_{\pi|_{\underline{\mathcal{F}}}} = \prod_{v \in I} \mu_v \quad \text{inside } Q_w[T^g].$$

Thus $F \otimes_Q Q_w = Q_w[T^g]/(\mu) = \prod_{v \in I} Q_w[T^g]/(\mu_v) = \prod_{v|w} F_v$. So I is the set of places of F dividing w and $F_v = Q_w[T^g]/(\mu_v)$ is the completion of F at v , justifying our notation. Let π_v be the image of π in F_v . Its minimal polynomial over Q_w is μ_v . This implies that $E \otimes_Q Q_w$ decomposes further

$$E \otimes_Q Q_w = \bigoplus_{v \in I} \text{End}_R(\underline{N}_v^{\oplus n_v}) = \bigoplus_{v \in I} E \otimes_F F_v$$

and $E \otimes_F F_v \cong \text{End}_R(\underline{N}_v^{\oplus n_v})$.

Now fix a place v above w and consider the diagram of field extensions

$$\begin{array}{c}
 & & & & \mathbb{F}_v \mathbb{F}_s & & \\
 & & & & \swarrow^{g/h} & & \\
 \mathbb{F}_v & & & & \mathbb{F}_w \mathbb{F}_s & & \\
 & & & & \swarrow^{g/h} & & \\
 & & & & \mathbb{F}_w \mathbb{F}_s \cap \mathbb{F}_v & & \\
 & & & & \downarrow i & & \\
 & & & & \mathbb{F}_w(\mathbb{F}_v \cap \mathbb{F}_s) & & \\
 & & & & \downarrow i & & \\
 & & & & \mathbb{F}_v \cap \mathbb{F}_s & & \\
 & & & & \swarrow^{g/h} & & \\
 \mathbb{F}_w & & & & \mathbb{F}_s & & \\
 & & & & \swarrow^{g/h} & & \\
 & & & & \mathbb{F}_w \cap \mathbb{F}_s & & \\
 & & & & \downarrow i & & \\
 \mathbb{F}_w & & & & \mathbb{F}_w \cap \mathbb{F}_s & & \\
 & & & & \downarrow f & & \\
 \mathbb{F}_w & & & & \mathbb{F}_w \cap \mathbb{F}_s & & \\
 & & & & \downarrow f & & \\
 \mathbb{F}_q & & & & \mathbb{F}_q & & \\
 & & & & \downarrow f & & \\
 & & & & \mathbb{F}_q & &
 \end{array}$$

Let $h := [\mathbb{F}_v \cap \mathbb{F}_s : \mathbb{F}_q] = \gcd([\mathbb{F}_v : \mathbb{F}_q], g)$. Let $i := [\mathbb{F}_w : \mathbb{F}_q]$. From the formulas

$$\begin{aligned}
 [\mathbb{F}_w \mathbb{F}_s : \mathbb{F}_w] &= [\mathbb{F}_s : \mathbb{F}_q] = g, \\
 [\mathbb{F}_w(\mathbb{F}_v \cap \mathbb{F}_s) : \mathbb{F}_w] &= [\mathbb{F}_v \cap \mathbb{F}_s : \mathbb{F}_q] = h, \\
 [\mathbb{F}_w \mathbb{F}_s : (\mathbb{F}_w \mathbb{F}_s \cap \mathbb{F}_v)] &= [\mathbb{F}_v \mathbb{F}_s : \mathbb{F}_v] = [\mathbb{F}_s : \mathbb{F}_v \cap \mathbb{F}_s] = \frac{g}{h}, \quad \text{and} \\
 \mathbb{F}_w(\mathbb{F}_v \cap \mathbb{F}_s) &\subset \mathbb{F}_w \mathbb{F}_s \cap \mathbb{F}_v,
 \end{aligned}$$

we obtain $\mathbb{F}_w \mathbb{F}_s \cap \mathbb{F}_v = \mathbb{F}_w(\mathbb{F}_v \cap \mathbb{F}_s) = \mathbb{F}_q^{fhi}$. Let $F_{v,L}$ be the compositum of $Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s$ and F_v in an algebraic closure of Q_w . Note that $F_{v,L}$ is well defined since $\mathbb{F}_s/\mathbb{F}_q$ is Galois. Let $F_{v,L}[T']$ be the non-commutative polynomial ring with

$$T' \cdot (a \otimes b) = (a \otimes b^{q^{fhi}}) \cdot T' \quad \text{and} \quad T' \cdot x = x \cdot T'$$

for $a \in Q_w$, $b \in \mathbb{F}_s$, and $x \in F_v$ and set $\Delta_v = F_{v,L}[T'] / ((T')^{g/h} - \pi_v^i)$. Observe that the commutation rules of T' are well defined since $(Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s) \cap F_v$ has residue field $\mathbb{F}_w \mathbb{F}_s \cap \mathbb{F}_v = \mathbb{F}_q^{fhi}$ and is unramified over Q_w , because $Q_w \otimes_{\mathbb{F}_q} \mathbb{F}_s$ is. Moreover, the extension $F_{v,L}/F_v$ is unramified of degree $[\mathbb{F}_v \mathbb{F}_s : \mathbb{F}_v] = \frac{g}{h}$ and $\tilde{T} := (T')^{[F_v:\mathbb{F}_q]/fhi}$ is its Frobenius automorphism. Since $\tilde{T}^{g/h} = \pi_v^{[F_v:\mathbb{F}_q]/fh}$ in Δ_v , our Δ_v is just the cyclic algebra $(F_{v,L}/F_v, \tilde{T}, \pi_v^{[F_v:\mathbb{F}_q]/fh})$ and has Hasse invariant $\frac{[F_v:\mathbb{F}_q]}{[F_s:\mathbb{F}_q]} \cdot v(\pi_v)$; compare [Rei, p. 266]. We relate Δ_v to $E \otimes_F F_v$. Firstly by [Jac, Theorem 3.20] there exists a positive integer u such that $\underline{N}_v^{\oplus u} \cong R/R\mu_v(T^g)$. Therefore

$$M_u(E \otimes_F F_v) \cong M_u(\text{End}_R(\underline{N}_v^{\oplus un_v})) = \text{End}_R(\underline{N}_v^{\oplus un_v}) = M_{n_v}((R/R\mu_v(T^g))^{\text{op}}).$$

Secondly we choose integers m and n with $m > 0$ and $mi + ng = 1$. We claim that the morphism $R/R\mu_v(T^g) \rightarrow M_h(\Delta_v)$, which maps

$$a \otimes b \longmapsto \begin{pmatrix} a \otimes b & & & \\ & a \otimes b^{q^f} & & \\ & & \ddots & \\ & & & a \otimes b^{q^{f(h-1)}} \end{pmatrix} \quad \text{and} \quad T \longmapsto \pi_v^n \cdot \begin{pmatrix} 0 & & 1 & \\ & \ddots & & \\ & & & 1 \\ (T')^m & & & 0 \end{pmatrix}$$

for $a \in Q_w$ and $b \in \mathbb{F}_s$, is an isomorphism of F_v -algebras. It is well defined since it maps $T \cdot (a \otimes b)$ and $(a \otimes b^{q^f}) \cdot T$ to the same element because $(T')^m = (T')^{1/i}$ in $\text{Gal}(F_{v,L}/F_v)$, and it maps $T^g = (T^h)^{g/h}$ to

$\pi_v^{ng}(T')^{mg/h} \cdot \text{Id}_h = \pi_v \cdot \text{Id}_h$. Since $R\mu_v(T^g) \subset R$ is a maximal two sided ideal the morphism is injective. To prove surjectivity we compare the dimensions as Q_w -vector spaces. We compute

$$\begin{aligned} \dim_{F_v} M_h(\Delta_v) &= h^2 \cdot \left(\frac{g}{h}\right)^2 = g^2, \\ \dim_{Q_w \otimes_{\mathbb{F}_f} \mathbb{F}_s} (R/R\mu_v(T^g)) &= g \cdot \deg \mu_v = g \cdot [F_v : Q_w], \quad \text{and} \\ \dim_{Q_w} (R/R\mu_v(T^g)) &= g^2 \cdot [F_v : Q_w] = \dim_{Q_w} M_h(\Delta_v). \end{aligned}$$

Altogether $M_u(E \otimes_F F_v) \cong M_{h\nu_v}(\Delta_v^{\text{op}})$ and $\text{inv}_v E = -\text{inv}_v \Delta_v = -\frac{[\mathbb{F}_v : \mathbb{F}_q]}{[\mathbb{F}_s : \mathbb{F}_q]} \cdot v(\pi_v)$ as claimed.

It remains to convert this formula into the special form asserted for $v \nmid \varepsilon\infty$ or $v|\infty$. If $v|\infty$ and $\varepsilon \neq \infty$, let e_v be the ramification index of F_v/Q_∞ . Then we get from Theorem 7.8 the formula $q^{e \cdot \text{wt}(\underline{\mathcal{F}})} = |\pi|_\infty = q^{-v(\pi_v)/e_v}$, since the residue field of Q_∞ is \mathbb{F}_q . This implies as desired

$$-\frac{[\mathbb{F}_v : \mathbb{F}_q]}{[\mathbb{F}_s : \mathbb{F}_q]} \cdot v(\pi_v) = -\frac{[\mathbb{F}_v : \mathbb{F}_q] \cdot (-e_v e \cdot \text{wt}(\underline{\mathcal{F}}))}{e} = \text{wt}(\underline{\mathcal{F}}) \cdot [F_v : Q_\infty]$$

Finally if $w \neq \varepsilon, \infty$ is a place of Q , the local σ -shtuka $\underline{M}_w(\underline{\mathcal{F}})$ at w is étale. So $\mu = \text{mipo}_{\pi|\underline{\mathcal{F}}}$ has coefficients in A_w with constant term in A_w^\times . Therefore $v(\pi_v) = 0$ for all places v of F dividing w . \square

Example 9.4. Let $C = \mathbb{P}_{\mathbb{F}_q}^1$, $C \setminus \{\infty\} = \text{Spec } \mathbb{F}_q[t]$ and $L = \mathbb{F}_q$. Let d be a positive integer. Let $\mathcal{F}_i := \mathcal{O}(d[\frac{i}{2}] \cdot \infty) \oplus \mathcal{O}(d[\frac{i-1}{2}] \cdot \infty)$ for $i \in \mathbb{Z}$ and let $\tau := \begin{pmatrix} 0 & t^d \\ 1 & 0 \end{pmatrix}$. Then $\underline{\mathcal{F}} = (\mathcal{F}_i, H_i, \tau_i)$ is an abelian τ -sheaf of rank 2, dimension d , and characteristic $\varepsilon = V(t) \in \mathbb{P}^1$ over \mathbb{F}_q . Hence the Frobenius endomorphism π equals τ . If d is odd then $\underline{\mathcal{F}}$ is primitive (that means $(d, r) = 1$) and therefore simple by [BH1, Proposition 7.4]. In particular, π is semisimple. We have

$$\mu_\pi = \chi_v = x^2 - t^d = (x - \sqrt{t^d})(x + \sqrt{t^d})$$

which means that π_v is *not* absolutely semisimple in characteristic 2. Moreover, we calculate $r_{Q_v}(\chi_v, \chi_v) = 1 \cdot 1 \cdot 2 = 2$ whereas in the field extension $Q_v(\sqrt{t})/Q_v$ we have

$$r_{Q_v(\sqrt{t})}(\chi_v, \chi_v) = \begin{cases} 2 \cdot 2 \cdot 1 = 4 & \text{in characteristic 2,} \\ 1 \cdot 1 \cdot 1 + 1 \cdot 1 \cdot 1 = 2 & \text{in characteristic different from 2.} \end{cases}$$

Although the later has no further significance it illustrates the remark after Definition 6.3. By Theorem 9.1/3. we have $E = F = Q(\pi)$ commutative and $[E : Q] = 2 = r$. Moreover, $|\pi|_\infty = |\sqrt{t^d}|_\infty = q^{d/2}$ and χ_v is irreducible. But χ_v is not separable in characteristic 2.

If $d = 2n$ is even then the minimal polynomial of π is

$$\mu_\pi = \chi_v = x^2 - t^d = (x - t^{d/2})(x + t^{d/2}).$$

So π is semisimple if and only if $\text{char}(\mathbb{F}_q) \neq 2$. In this case $\underline{\mathcal{F}}$ is quasi-isogenous to the abelian τ -sheaf $\underline{\mathcal{F}}'$ with $\mathcal{F}'_i = \mathcal{O}_{C_L}(in \cdot \infty)^{\oplus 2}$ and $\tau'_i = \begin{pmatrix} -t^n & 0 \\ 0 & t^n \end{pmatrix}$. The quasi-isogeny $f : \underline{\mathcal{F}}' \rightarrow \underline{\mathcal{F}}$ is given by $f_{0,\eta} = \begin{pmatrix} -t^n & t^n \\ 1 & 1 \end{pmatrix} : \mathcal{F}'_{0,\eta} \xrightarrow{\sim} \mathcal{F}_{0,\eta}$. The abelian τ -sheaf $\underline{\mathcal{F}}'$ equals the direct sum $\underline{\mathcal{F}}^{(1)} \oplus \underline{\mathcal{F}}^{(2)}$ where $\mathcal{F}'_i^{(j)} = \mathcal{O}_{C_L}(in \cdot \infty)$ and $\tau_i^{(j)} = (-1)^j t^n$. Note that $\underline{\mathcal{F}}^{(1)}$ and $\underline{\mathcal{F}}^{(2)}$ are not isogenous over \mathbb{F}_q , since the equation $-t^n \cdot \sigma^*(g) = g \cdot t^n$ has no solution $g \in Q$ for $\text{char}(\mathbb{F}_q) \neq 2$. Therefore

$$Q \oplus Q = \bigoplus_{j=1}^2 Q\text{End}(\underline{\mathcal{F}}^{(j)}) \cong E = F = Q[x]/(x^2 - t^{2n}) \cong Q \oplus Q.$$

Now we consider the same abelian τ -sheaf over $L = \mathbb{F}_{q^2}$. This means $\pi = \tau^2 = t^d \in Q$ and therefore $\chi_v = (x - t^d)^2$. Thus π is semisimple. By Theorem 9.1/4 we have $F = Q(\pi) = Q$ and E is central

simple over Q with $[E : Q] = 4$ and $\text{inv}_\infty E = \text{inv}_\varepsilon E = \frac{d}{2}$. Moreover, $|\pi|_\infty = |t^d|_\infty = q^d$. In this case, π_v is absolutely semisimple. Note that if d is even and $\text{char}(\mathbb{F}_q) = 2$ this is another example for Theorem 6.15.

If d is odd then $\underline{\mathcal{F}}$ is still primitive, whence simple and E is a division algebra. If $d = 2n$ is even then the abelian τ -sheaves $\underline{\mathcal{F}}^{(1)}$ and $\underline{\mathcal{F}}^{(2)}$ defined above are isomorphic $\underline{\mathcal{F}}^{(1)} \xrightarrow{\sim} \underline{\mathcal{F}}^{(2)}$, $1 \mapsto \lambda$ where $\lambda \in \mathbb{F}_{q^2}$ satisfies $\lambda^{q-1} = -1$. Therefore $M_2(Q) = M_2(\text{QEnd}(\underline{\mathcal{F}}^{(1)})) \cong E$ in accordance with the Hasse invariants just computed.

Example 9.5. We compute another example which displays other phenomena. Let $C = \mathbb{P}_{\mathbb{F}_q}^1$ and let $C \setminus \{\infty\} = \text{Spec } \mathbb{F}_q[t]$. Let $\mathcal{F}_i = \mathcal{O}_{C_L}([\frac{i-1}{2}] \cdot \infty)^{\oplus 2} \oplus \mathcal{O}_{C_L}([\frac{i}{2}] \cdot \infty)^{\oplus 2}$, let Π_i be the natural inclusion, and let τ_i be given by the matrix

$$T := \begin{pmatrix} 0 & 0 & 0 & a \\ 0 & b & 1 & 0 \\ t & 0 & -b & 0 \\ 0 & t & 0 & 0 \end{pmatrix} \quad \text{with } a, b \in \mathbb{F}_q \setminus \{0\}.$$

Then $\underline{\mathcal{F}}$ is an abelian τ -sheaf of rank 4 and dimension 2 with $l = 2, k = 1$ and characteristic $\varepsilon = V(t) \in \mathbb{P}^1$. One checks that the minimal polynomial of the matrix T is $x^4 - b^2x^2 - at^2$ which is irreducible over Q if $\text{char}(\mathbb{F}_q) \neq 2$, since it has neither zeroes in $\mathbb{F}_q[t]$ nor quadratic factors in $Q[x]$. If $\text{char}(\mathbb{F}_q) = 2$ then the minimal polynomial is a square and $\underline{\mathcal{F}}$ is not semisimple.

For $L = \mathbb{F}_q$ and $2 \nmid q$ we obtain $\pi = \tau$ semisimple and $E = F = Q(\pi) = Q[x]/(x^4 - b^2x^2 - at^2)$.

For $L = \mathbb{F}_{q^2}$ we have $\pi = \tau^2$ and the minimal polynomial of π over Q is $x^2 - b^2x - at^2$, which is irreducible also in characteristic 2 since it has no zeroes in $\mathbb{F}_q[t]$. Hence π is semisimple, F is a field with $[F : Q] = 2$ and $[E : F] = 4$ by Corollary 6.6. This again illustrates Theorem 6.15. We compute the decomposition of ∞ and ε in F .

Decomposition of ε : Modulo t the polynomial $x^2 - b^2x - at^2$ has two zeroes $x = b^2$ and $x = 0$ in \mathbb{F}_q . So by Hensel's lemma $F \otimes_Q Q_\varepsilon \cong F_v \oplus F_{v'}$ splits with $F_v \cong F_{v'} \cong Q_\varepsilon$ and $v(\pi) = 0$ and $v'(\pi) = v'(at^2) = 2$. Thus the Hasse invariants of E are $\text{inv}_v E = \text{inv}_{v'} E = 0$.

Decomposition of ∞ : Set $y = \pi/t$. Then $y^2 - \frac{b^2}{t}y - a = 0$.

Case (a). If $2|q$ then $(y - a^{q/2})^2 - \frac{b^2}{t}(y - a^{q/2}) - \frac{b^2}{t}a^{q/2} = 0$, that is, ∞ ramifies in F , $F \otimes_Q Q_\infty = F_w$ with $w(\frac{\pi}{t} - a^{q/2}) = 1$ and $w(\frac{1}{t}) = 2 \cdot \infty(\frac{1}{t}) = 2$. So $[F_w : Q_\infty] = 2$ and $\text{inv}_w E = 0$.

Case (b). If $2 \nmid q$ and $\sqrt{a} \in \mathbb{F}_q$ then the polynomial $y^2 - \frac{b^2}{t}y - a$ has two zeroes $y = \pm\sqrt{a}$ modulo $\frac{1}{t}$. So by Hensel's lemma $F \otimes_Q Q_\infty \cong F_w \oplus F_{w'}$ splits with $[F_w : Q_\infty] = [F_{w'} : Q_\infty] = 1$. Thus the local Hasse invariants of E are $\text{inv}_w E = \text{inv}_{w'} E = \frac{1}{2}$. As was remarked in 9.2 such a distribution of the Hasse invariants can occur only if $d \geq 2$.

Case (c). If $2 \nmid q$ and $\sqrt{a} \notin \mathbb{F}_q$ then $y^2 - \frac{b^2}{t}y - a$ is irreducible modulo $\frac{1}{t}$ and ∞ is inert in F , $F \otimes_Q Q_\infty = F_w$ with $[F_w : Q_\infty] = 2$. Thus the Hasse invariant of E is $\text{inv}_w E = 0$.

In case (b) E is a division algebra and $\underline{\mathcal{F}}$ is simple. In cases (a) and (c) $E \cong M_2(F)$ and $\underline{\mathcal{F}}$ is quasi-isogenous to $(\underline{\mathcal{F}}')^{\oplus 2}$ for an abelian τ -sheaf $\underline{\mathcal{F}}'$ of rank 2, dimension 1 and $\text{QEnd}(\underline{\mathcal{F}}') = F$. This surprising result is due to the fact that $\underline{\mathcal{F}}'$, being of dimension 1, is associated with a Drinfeld module and thus of the form $\mathcal{F}'_i = \mathcal{O}_{C_L}([\frac{i}{2}] \cdot \infty) \oplus \mathcal{O}_{C_L}([\frac{i-1}{2}] \cdot \infty)$ with $\tau'_i = \begin{pmatrix} c & t \\ d & 0 \end{pmatrix}$ and $c, d \in \mathbb{F}_{q^2}$. Then $\pi' = (\tau')^2 = \begin{pmatrix} c^{q+1} + d^q t & ct \\ c^q d & dt \end{pmatrix}$ has minimal polynomial $x^2 - (c^{q+1} + (d + d^q)t)x + d^{q+1}t^2$ which must be equal to $x^2 - b^2x - at^2$. This is possible only if $d + d^q = 0$ and $d^{q+1} = -a$. So either $d \in \mathbb{F}_q$ and $2|q$ and we are in case (a), or $d \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, $d^q = -d$, and $a = d^2$. The later implies $2 \nmid q$ and $\sqrt{a} = d \notin \mathbb{F}_q$ and we are in case (c). If we choose $c = b$ in case (c) a quasi-isogeny $f : \underline{\mathcal{F}} \rightarrow (\underline{\mathcal{F}}')^{\oplus 2}$ over \mathbb{F}_{q^2} is given for instance by

$$\begin{pmatrix} d & a & -bd/t & 0 \\ 0 & 0 & -d & a \\ 0 & 0 & d/t & a/t \\ 1 & -d & 0 & bd/t \end{pmatrix}.$$

10 Kernel Ideals for Pure Anderson Motives

In this section we investigate which orders of E can arise as endomorphism rings $\text{End}(\underline{M})$ for pure Anderson motives \underline{M} . For this purpose we define for each right ideal of the endomorphism ring $\text{End}(\underline{M})$ an isogeny with target \underline{M} and discuss its properties. This generalizes Gekeler's results for Drinfeld modules [Gek, §3] and translates the theory of Waterhouse [Wat, §3] for abelian varieties to the function field case. These two sources are themselves the translation, respectively the higher dimensional generalization of Deuring's work on elliptic curves [Deu].

Let \underline{M} be a pure Anderson motive over L and abbreviate $R := \text{End}(\underline{M})$. Let $I \subset R$ be a right ideal which is an A -lattice in $E := R \otimes_A Q$. This is equivalent to saying that I contains an isogeny, since every lattice contains some isogeny $a \cdot \text{id}_{\underline{M}}$ for $a \in A$ and conversely the existence of an isogeny $f \in I$ implies that the lattice $f \cdot f^\vee \cdot R$ is contained in I .

Definition 10.1. 1. Let \underline{M}^I be the pure Anderson sub-motive of \underline{M} whose underlying A_L -module is $\sum_{g \in I} \text{im}(g)$. This is indeed a pure Anderson motive, since if $I = f_1 R + \dots + f_n R$ are arbitrary generators, then \underline{M}^I equals the image of the morphism

$$(f_1, \dots, f_n) : \underline{M} \oplus \dots \oplus \underline{M} \longrightarrow \underline{M}.$$

As I contains an isogeny, \underline{M}^I has the same rank as \underline{M} and the natural inclusion is an isogeny which we denote $f_I : \underline{M}^I \rightarrow \underline{M}$.

2. If $I = \{f \in R : \text{im}(f) \subset \underline{M}^I\}$ then I is called a kernel ideal for \underline{M} .

The later terminology is borrowed from Waterhouse [Wat, §3]. Since $\{f \in R : \text{im}(f) \subset \underline{M}^I\}$ is the right ideal annihilating coker f_I one should maybe use the name "cokernel ideal" instead.

Proposition 10.2. Let $I \subset R$ be a right ideal which is a lattice, and consider the right ideal $J := \{f \in R : \text{im}(f) \subset \underline{M}^I\} \subset R$ containing I . Then $\underline{M}^J = \underline{M}^I$. In particular, J is a kernel ideal for \underline{M} . We call J the kernel ideal for \underline{M} associated with I .

Proof. Obviously J is a right ideal and $\underline{M}^J \subset \underline{M}^I$ by definition of J . Conversely $\underline{M}^I \subset \underline{M}^J$ since $I \subset J$. \square

Lemma 10.3. 1. For any $g \in I$, $f_I^{-1} \circ g : \underline{M} \rightarrow \underline{M}^I$ is a morphism and $g = f_I \circ (f_I^{-1} \circ g)$.

2. If $I = gR$ is principal, g an isogeny, then $f_I^{-1} \circ g : \underline{M} \rightarrow \underline{M}^I$ is an isomorphism and I is a kernel ideal.

Proof. 1 is obvious since the image of g lies inside \underline{M}^I .

2. Clearly $f_I^{-1} \circ g$ is injective since g is an isogeny and surjective by construction, hence an isomorphism. To show that I is a kernel ideal let $f \in R$ satisfy $\text{im}(f) \subset \underline{M}^I$. Consider the diagram

$$\begin{array}{ccccc} \underline{M} & \xrightarrow{f_I^{-1} \circ f} & \underline{M}^I & \xrightarrow{f_I} & \underline{M} \\ \downarrow & \nearrow & & & \\ \underline{M} & & & & \end{array}$$

$h \downarrow$ $f_I^{-1} \circ g$

and let $h := (f_I^{-1} \circ g)^{-1} \circ (f_I^{-1} \circ f)$. Then $f = gh \in I$ as desired. \square

Example. If $a \in A$ and $I = aR$, then $\underline{M}^I = a\underline{M}$ and $\text{coker } f_I = \underline{M}/a\underline{M}$. More generally if $\mathfrak{a} \subset A$ is an ideal and $I = \mathfrak{a}R$ then $\underline{M}^I = \mathfrak{a}\underline{M}$ and $\text{coker } f_I = \underline{M}/\mathfrak{a}\underline{M}$.

Proposition 10.4. Let $I \subset R$ and $J \subset \text{End}(\underline{M}^I)$ be right ideals which are lattices in E . Then also the product $K := f_I \cdot J \cdot f_I^{-1} \cdot I$ is a right ideal of R and a lattice in E and $f_K^{-1} \circ f_I \circ f_J$ is an isomorphism of $(\underline{M}^I)^J$ with \underline{M}^K

$$(\underline{M}^I)^J \xrightarrow{f_J} \underline{M}^I \xrightarrow{f_I} \underline{M} \xleftarrow{f_K} \underline{M}^K.$$

Proof. If $f \in I$ and $g \in J$ then the morphism $f_I^{-1} \circ f : \underline{M} \rightarrow \underline{M}^I$ can be composed with $f_I \circ g$ to yield an element of R . Since I and J contain isogenies, K is a right ideal and contains an isogeny. Clearly the images of $f_I \circ f_J$ and f_K in \underline{M} coincide since they equal the sum $\sum_{i,j} f_I \circ g_j \circ f_I^{-1} \circ f_i(\underline{M})$ for sets of generators $\{f_i\}$ of I and $\{g_j\}$ of J . \square

Theorem 10.5. *Let $I, J \subset \text{End}(\underline{M}) =: R$ be right ideals which are lattices in $E := R \otimes_A Q$ and consider the following assertions:*

1. *I and J are isomorphic R -modules,*
2. *the pure Anderson motives \underline{M}^I and \underline{M}^J are isomorphic.*

Then 1 implies 2 and if moreover I and J are kernel ideals, also 2 implies 1.

Proof. $1 \Rightarrow 2$. Since I and J are lattices, the R -isomorphism $I \rightarrow J$ extends to an E -isomorphism of E and is thus given by left multiplication with a unit $g \in E^\times$, that is, $J = gI$. There is an $a \in A$ such that $ag \in I \subset R$. Then $\text{im}(ag) \subset \underline{M}^I$, that is, $f_I^{-1} \circ ag : \underline{M} \rightarrow \underline{M}^I$ is an isogeny.

Let K be the right ideal $f_I \cdot (f_I^{-1} \circ ag \circ f_I \cdot \text{End}(\underline{M}^I)) \cdot f_I^{-1} \cdot I$ of R . We claim that $\underline{M}^K \cong \underline{M}^{(ag)I}$. Namely, $\underline{M}^{(ag)I} \subset \underline{M}^K$ since $agI \subset K$. Conversely if $f \in I$, $h \in \text{End}(\underline{M}^I)$, and $m \in \underline{M}$, then we find $m' := f_I \circ h \circ f_I^{-1} \circ f(m) \in \underline{M}^I$, that is, $m' = \sum_i f_i(m_i)$ for suitable $f_i \in I$ and $m_i \in \underline{M}$. It follows that $ag(m') = \sum_i ag f_i(m_i) \in \underline{M}^{(ag)I}$ and therefore $\underline{M}^{(ag)I} = \underline{M}^K$.

Applying Lemma 10.3 and Proposition 10.4 now yields an isomorphisms $\underline{M}^I \cong \underline{M}^K = \underline{M}^{(ag)I}$. Likewise we obtain $\underline{M}^J \cong \underline{M}^{aJ}$ and the equality $aJ = agI$ then implies $\underline{M}^J \cong \underline{M}^I$ as desired.

$2 \Rightarrow 1$. Let I and J be kernel ideals and let $u : \underline{M}^I \rightarrow \underline{M}^J$ be an isomorphism. There is an $a \in A$ with $a\underline{M} \subset \underline{M}^I$. Therefore $g := f_J \circ u \circ (f_I^{-1} \circ a) : \underline{M} \rightarrow \underline{M}$ is an isogeny.

We claim that $gI = aJ$, that is, left multiplication by $a^{-1}g$ is an isomorphism of I with J . Let $f \in I$, then $h := f_J \circ u \circ (f_I^{-1} \circ f) \in R$ has $\text{im}(h) \subset \underline{M}^J$. So $h \in J$ since J is a kernel ideal, and $gf = ah \in aJ$, since a commutes with all morphisms. Conversely let $h \in J$, then $f := f_I \circ u^{-1} \circ (f_J^{-1} \circ h) \in R$ has $\text{im}(f) \subset \underline{M}^I$. So $f \in I$ since I is a kernel ideal, and $ah = gf \in gI$ as desired. \square

Proposition 10.6. *Let $I \subset R$ be a right ideal which is a lattice in E . Then $f_I \cdot \text{End}(\underline{M}^I) \cdot f_I^{-1}$ contains the left order $\mathcal{O} = \{f \in E : fI \subset I\}$ of I and equals it if I is a kernel ideal.*

Remark. Recall that $\text{End}(\underline{M}^I) \otimes_A Q$ is identified with E by mapping $h \in \text{End}(\underline{M}^I)$ to $f_I \circ h \circ f_I^{-1}$.

Proof. Let $f \in \mathcal{O}$ and $g \in I$. Then $fg \in I$ and $f_I^{-1} \circ f \circ f_I \circ (f_I^{-1} \circ g) = f_I^{-1} \circ fg$ is a morphism from \underline{M} to \underline{M}^I . If g varies, the images of $f_I^{-1} \circ g$ exhaust all of \underline{M}^I . Hence $f_I^{-1} \circ f \circ f_I$ is indeed an endomorphism of \underline{M}^I . Conversely let I be a kernel ideal and let $f = f_I \circ h \circ f_I^{-1} \in f_I \cdot \text{End}(\underline{M}^I) \cdot f_I^{-1}$. If $g \in I$ then $f \circ g = f_I \circ h \circ (f_I^{-1} \circ g) \in R$ has $\text{im}(f \circ g) \subset \underline{M}^I$. So $fg \in I$ as desired. \square

We will now draw conclusions about the endomorphism ring R similar to Waterhouse' results [Wat] on abelian varieties by simply translating his arguments.

Theorem 10.7. *Every maximal order in E occurs as the endomorphism ring $f \cdot \text{End}(\underline{M}') \cdot f^{-1} \subset E$ of a pure Anderson motive \underline{M}' isogenous to \underline{M} via an isogeny $f : \underline{M}' \rightarrow \underline{M}$.*

Proof. Let S be a maximal order of E . Then the lattice R contains aS for some $a \in A$. Consider the right ideal $I = aS \cdot R$ whose left order contains S . By Proposition 10.6, $f_I \cdot \text{End}(\underline{M}^I) \cdot f_I^{-1}$ contains the left order of I . Since S is maximal we find $S = f_I \cdot \text{End}(\underline{M}^I) \cdot f_I^{-1}$. \square

Theorem 10.8. *If E is semisimple and $\text{End}(\underline{M})$ is a maximal order in E , so is $f_I \cdot \text{End}(\underline{M}^I) \cdot f_I^{-1}$ for any right ideal $I \subset R$.*

Proof. By [Rei, Theorem 21.2] the left order of I is also maximal and then Proposition 10.6 yields the result. \square

From now on we assume that L is a finite field and we set $e := [L : \mathbb{F}_q]$. Let π be the Frobenius endomorphism of \underline{M} .

Proposition 10.9. *The order R in E contains π and $\deg(\pi)/\pi$.*

Proof. Clearly the isogeny π belongs to R . Let now $a \in \deg(\pi)$. Then a annihilates $\text{coker } \pi$ by 2.10 and so there is an isogeny $f : \underline{M} \rightarrow \underline{M}$ with $\pi \circ f = a$. The image a/π of f in E belongs to R . \square

Proposition 10.10. *If \underline{M} is a semisimple pure Anderson motive over a finite field and $\text{End}(\underline{M})$ is a maximal order in $E = \text{End}(\underline{M}) \otimes_A Q$, then every right ideal $I \subset \text{End}(\underline{M})$, which is a lattice, is a kernel ideal for \underline{M} , and $\deg(f_I) = N(I) := (N(f) : f \in I)$.*

Proof. (cf. [Wat, Theorem 3.15]) Let $f \in I$, then $f = f_I \circ f_I^{-1}f$ and $N(f) \in \deg(f) \subset \deg(f_I)$ by Lemma 2.9. Therefore $N(I) \subset \deg(f_I)$. Let R' be the left order of I . It is maximal by [Rei, Theorem 21.2]. For a suitable $a \in A$ the set $J' := \{x \in E : xI \subset aR\}$ is a right ideal in R' and a lattice in E and satisfies $J' \cdot I = aR$ by [Rei, Theorem 22.7]. Let $J := f_I^{-1}J'I \subset \text{End}(\underline{M}^I)$ be the induced right ideal of $\text{End}(\underline{M}^I) = f_I^{-1}R'f_I$; see 10.6. Then $\text{coker } f_I \circ f_J = \text{coker } f_{J'I} = \text{coker } a$ by Proposition 10.4. Therefore Theorem 7.3 and [Rei, 24.12 and 24.11] imply

$$N(a) \cdot A = N(J') \cdot N(I) \subset (\deg f_J)(\deg f_I) = \deg(a) = N(a) \cdot A.$$

By the above we must have $N(I) = \deg(f_I)$ since A is a Dedekind domain. If I were not a kernel ideal its associated kernel ideal would be a larger ideal with the same norm. But this is impossible by [Rei, 24.11]. \square

Like for abelian varieties there is a strong relation between the ideal theory of orders of E and the investigation of isomorphism classes of pure Anderson motives isogenous to \underline{M} . We content ourselves with the following result which is analogous to Waterhouse [Wat, Theorem 6.1]. The interested reader will find many other results without much difficulty.

Theorem 10.11. *Let \underline{M} be a simple pure Anderson motive of rank r and dimension d over the smallest possible field \mathbb{F}_q . Then*

1. $\text{End}(\underline{M})$ is commutative and $E := \text{End}(\underline{M}) \otimes_A Q = Q(\pi)$.
2. All orders R in $Q(\pi)$ containing π are endomorphism rings of pure Anderson motives isogenous to \underline{M} . Any such order automatically contains $N(\pi)/\pi = N_{Q(\pi)/Q}(\pi)/\pi$.
3. For each such R the isomorphism classes of pure Anderson motives isogenous to \underline{M} with endomorphism ring R correspond bijectively to the isomorphism classes of A -lattices in E with order R .

Proof. 1 follows from 6.11 and 9.3.

2. Let R be an order in $Q(\pi)$ containing π and let $v \neq \varepsilon$ be a maximal ideal of A . Since $[E : Q] = r$ and E_v is semisimple, there is by Lemma 7.2 an isomorphism $E_v \xrightarrow{\sim} V_v \underline{M}$ of (left) E_v -modules given by $f \mapsto f(x)$ for a suitable $x \in V_v \underline{M}$. It identifies $R_v := R \otimes_A A_v$ with a π -stable lattice $\Lambda_v = R_v \cdot x$ in $V_v \underline{M}$, which without loss of generality is contained in $T_v \underline{M}$. By Proposition 4.4 there is an isogeny $f : \underline{M}' \rightarrow \underline{M}$ of pure Anderson motives with $T_v f(T_v \underline{M}') = \Lambda_v$. By Theorem 4.2 we conclude

$$\text{End}(\underline{M}') \otimes_A A_v = \text{End}_{A_v[\pi]}(\Lambda_v) = R_v.$$

For $v = \varepsilon$ note that $Q_{\varepsilon, L} = Q_\varepsilon$ since $L = \mathbb{F}_q$. In particular $\mathbb{F}_\varepsilon = \mathbb{F}_q$. Since $\dim_{Q_\varepsilon} \underline{N}_\varepsilon(\underline{M}) = r = [E : Q]$, Theorem 3.7 together with Lemma 7.2 show that E_ε is isomorphic to $\underline{N}_\varepsilon(\underline{M})$ as left E_ε -modules. Since R contains π , the image of $R_\varepsilon := R \otimes_A A_\varepsilon$ in $\underline{N}_\varepsilon(\underline{M})$ is a local σ -substtuka \hat{M}' of $\underline{M}_\varepsilon(\underline{M})$ of the same rank. (If it is not contained in $\underline{M}_\varepsilon(\underline{M})$, multiply it with a suitable $a \in A$.) Then Proposition 3.9 yields an isogeny of pure Anderson motives $f : \underline{M}' \rightarrow \underline{M}$ such that $\underline{M}_\varepsilon(f)(\underline{M}_\varepsilon(\underline{M}')) = \hat{M}'$ and

$$\text{End}(\underline{M}') \otimes_A A_\varepsilon = \text{End}_{A_\varepsilon[\phi]}(\hat{M}') = R_\varepsilon$$

by Theorem 3.7. Since each of these operations only modifies $\text{End}(\underline{M})$ at the respective place v , this shows that we may modify \underline{M} at all places to obtain a pure Anderson motive \underline{M}' with $\text{End}(\underline{M}') = R$. Now the last statement follows from Proposition 10.9 and Theorem 7.3.

3. Let R be such an order. By what we proved in 2 there is a pure Anderson motive \tilde{M} for which all $T_v \tilde{M} \cong R_v$ and $\underline{M}_\varepsilon(\tilde{M}) \cong R_\varepsilon$. Let $I \subset R$ be a (right) ideal which is an A -lattice in E and consider the isogeny $f_I : \underline{M}^I \rightarrow \underline{M}$. Under the above isomorphisms $T_v f_I(T_v \underline{M}^I) \cong I \otimes_A A_v =: I_v$ and $\underline{M}_\varepsilon f_I(\underline{M}_\varepsilon \underline{M}^I) \cong I \otimes_A A_\varepsilon =: I_\varepsilon$. Conversely if $f : \underline{M}' \rightarrow \underline{M}$ is an isogeny then $\underline{M}_v f(\underline{M}_v \underline{M}')$ is a (left) R_v -module because $R = \text{End}(\underline{M})$, hence isomorphic to an R_v -ideal I_v . This shows that any isogeny $f : \underline{M}' \rightarrow \underline{M}$ is of the form $f_I : \underline{M}^I \rightarrow \underline{M}$.

If now $f \in R$ satisfies $\text{im}(f) \subset \underline{M}^I$ then $f \in I_\varepsilon$ and $f \in I_v$ for all v and therefore $f \in I$. This shows that every I is a kernel ideal for \underline{M} . By Proposition 10.6, $\text{End}(\underline{M}^I)$ is the (left) order of I . Since every lattice with order R in E is isomorphic to an ideal of R , we have

$$\begin{array}{c} \{A\text{-lattices in } E \text{ with order } R\} / \sim \\ \parallel \\ \{I \subset R \text{ Ideals with order } R\} / \sim \end{array} \xrightarrow{\sim} \{ \underline{M}^I \xrightarrow{f_I} \underline{M} \rightarrow \underline{M} \text{ with } \text{End}(\underline{M}^I) = R \} / \sim$$

and the assertion now follows from Theorem 10.5. \square

References

- [An1] G. Anderson: t -Motives, *Duke Math. J.* **53** (1986), 457–502.
- [An2] G. Anderson: On Tate Modules of Formal t -Modules, *Internat. Math. Res. Notices* **2** (1993), 41–52.
- [BH1] M. Bornhofen, U. Hartl: *Pure Anderson Motives and Abelian τ -Sheaves*, Preprint on [arXiv:0709.2809](https://arxiv.org/abs/0709.2809), version 1 from 09/18/2007.
- [BH2] M. Bornhofen, U. Hartl: *Pure Anderson Motives and Abelian τ -Sheaves over Finite Fields*, Preprint on [arXiv:math.NT/0609733](https://arxiv.org/abs/math/0609733), version 3 from 09/18/2007.
- [BS] A. Blum, U. Stuhler: Drinfeld Modules and Elliptic Sheaves, in: *Vector Bundles on Curves: New Directions*, S. Kumar, G. Laumon, U. Stuhler, M. S. Narasimhan, eds., pp. 110–188, Lecture Notes in Mathematics 1649, Springer-Verlag, Berlin, etc. 1991.
- [Boe] G. Böckle: Global L -functions over function fields. *Math. Ann.* **323** (2002), no. 4, 737–795.
- [Bou] N. Bourbaki: *Algèbre, chapitre 8*. Hermann, Paris (1958).
- [Deu] M. Deuring: Die Typen der Multiplikatorenringe elliptischer Funktionenkörper, *Abh. Math. Sem. Hansischen Univ.* **14** (1941), 197–272.
- [Dr1] V.G. Drinfeld: Elliptic Modules, *Math. USSR-Sb.* **23** (1976), 561–592.
- [Dr2] V.G. Drinfeld: Commutative Subrings of Certain Noncommutative Rings, *Funct. Anal. Appl.* **11** (1977), 9–12.
- [EGA] A. Grothendieck: *Éléments de Géométrie Algébrique*, Publ. Math. IHES **4**, **8**, **11**, **17**, **20**, **24**, **28**, **32**, Bures-Sur-Yvette, 1960–1967; see also Grundlehren **166**, Springer-Verlag, Berlin etc. 1971.
- [Gar] F. Gardeyn: A Galois criterion for good reduction of τ -sheaves, *J. Number Theory* **97** (2002), 447–471.
- [Gek] E.-U. Gekeler: On finite Drinfel'd modules, *J. Algebra* **141** (1991), no. 1, 187–203.
- [Gos] D. Goss: *Basic Structures of Function Field Arithmetic*. Springer-Verlag, Berlin, etc. (1996).
- [Ha1] U. Hartl: Uniformizing the Stacks of Abelian Sheaves, in *Number Fields and Function fields - Two Parallel Worlds, Papers from the 4th Conference held on Texel Island, April 2004*, G. van der Geer, B. Moonen, R. Schoof, Editors, Progress in Math. 239, Birkhäuser-Verlag, Basel 2005, pp. 167–222.
- [Ha2] U. Hartl: *Period Spaces for Hodge Structures in Equal Characteristic*, Preprint on [arXiv:math.NT/0511686](https://arxiv.org/abs/math/0511686).

- [Jac] N. Jacobson: *The Theory of Rings*, AMS Mathematical Surveys, vol. II, American Mathematical Society, New York, 1943.
- [Kat] N. Katz: p -adic properties of modular schemes and modular forms, *Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972)*, pp. 69–190. LNM **350**, Springer, Berlin, 1973.
- [Lan] S. Lang: *Abelian Varieties*. Springer, New York (1983).
- [Lau] G. Laumon: *Cohomology of Drinfeld Modular Varieties I*, Cambridge Studies in Advanced Mathematics **41**, Cambridge University Press, Cambridge, 1996.
- [PT] R. Pink, M. Traulsen: The isogeny conjecture for t -motives associated to direct sums of Drinfeld modules, *J. Number Theory* **117**, no. 2 (2006), 355–375.
- [Rei] I. Reiner: *Maximal Orders*. Academic Press, London, etc. (1975).
- [Tag] Y. Taguchi: The Tate conjecture for t -motives, *Proc. Amer. Math. Soc.* **123** (1995), no. 11, 3285–3287.
- [Tam] A. Tamagawa: Generalization of Anderson’s t -motives and Tate conjecture, in: *Moduli Spaces, Galois Representations and L -Functions*, Sūrikaiseikikenkyūsho Kōkyūroku, n° 884, Kyoto (1994), 154–159.
- [Tat] J. Tate: Endomorphisms of Abelian Varieties over Finite Fields, *Invent. Math.* **2** (1966), 134–144.
- [TW] Y. Taguchi, D. Wan: L -Functions of φ -Sheaves and Drinfeld Modules, *J. Amer. Math. Soc.* **9**, no. 3 (1996), 755–781.
- [Wat] W. C. Waterhouse: Abelian Varieties over Finite Fields, *Ann. Sci. École Norm. Sup. (4)*, **2** (1969), 521–560.
- [WM] W. Waterhouse, J. Milne: Abelian Varieties over Finite Fields, *1969 Number Theory Institute (Proc. Sympos. Pure Math., Vol. XX, Stony Brook, N.Y., 1969)*, pp. 53–64, Amer. Math. Soc., Providence, R.I. 1971.

Matthias Bornhofen
Am Rösslewald 4
D – 79874 Breitnau
Germany

Urs Hartl
Mathematisches Institut
Universität Münster
Einsteinstr. 62
D – 48149 Münster
Germany
<http://www.math.uni-muenster.de/u/urs.hartl/>